

AMENABILITY OF VECTOR-VALUED GROUP ALGEBRAS

A. GHAFARI AND S. JAVADI*

ABSTRACT. The purpose of carrying out this work is to develop the amenability notations for vector-valued group algebras. We prove that $L^1(G, A)$ is approximately weakly amenable where A is a unital separable Banach algebra. We give the necessary, and sufficient conditions for the existence of a left invariant mean on $L^\infty(G, A^*)$, $LUC(G, A^*)$, $WAP(G, A^*)$, and $C_0(G, A^*)$.

1. INTRODUCTION

In 1972, B. E. Johnson proved that a locally compact group G is amenable if and only if $L^1(G)$ is amenable [7]. The concept of Johnson's amenability for Banach algebras has been a main stream in the theory of Banach algebras. Here we develop the concept of Johnson's amenability for vector-valued Banach algebras.

Let G be a locally compact group with a fixed left Haar measure m and A be a unital separable Banach algebra. Let $L^1(G, A)$ be the set of all measurable vector-valued (equivalence classes of) functions $f : G \rightarrow A$ such that $\|f\|_1 = \int_G \|f(t)\| dm(t) < \infty$. Equipped with the norm $\|\cdot\|_1$ and the convolution product $*$ specified by:

$$f * g(x) = \int f(t)g(t^{-1}x)dm(t) \quad (f, g \in L^1(G, A)),$$

$L^1(G, A)$ is a Banach algebra. We prove the analogues of the classical results on amenability of Banach algebras. We show that G is amenable

MSC(2010): Primary: 65F05; Secondary: 46L05, 11Y50.

Keywords: Amenability, Banach algebras, Derivation, Group algebra, Invariant mean.

Received: 19 April 2014, Revised: 26 December 2015.

*Corresponding author.

if and only if $L^1(G, A)$ is amenable for each unital separable Banach algebra A . The symbol $M(G, A)$ stands for the space of regular A -valued Borel measures of bounded variation on G . The space $L^1(G, A)$ is a closed two-sided ideal of $M(G, A)$.

Let $L^\infty(G, A^*)$ be the set of all functions f of G into A^* that are scalarwise measurable and $N_\infty(\|f\|) = \text{loc ess sup}_{t \in G} (\|f(t)\|) < \infty$. From now on, A will be a separable Banach algebra. By Theorem 8.18.2 in [5], the dual of $L^1(G, A)$ may be identified with $L^\infty(G, A^*)$. Note that the dual of $L^1(G, A)$ is in general not $L^\infty(G, A^*)$. We show that every continuous derivation, from $L^1(G, A)$ into $L^\infty(G, A^*)$ is approximately inner, that is, of the form

$$D(a) = \lim_{\alpha} (F_{\alpha} \cdot a - a \cdot F_{\alpha})$$

for some $\{F_{\alpha}\}_{\alpha \in I} \in L^\infty(G, A^*)$.

Let $C(G, A^*)$ be the space of bounded continuous functions from G into A^* , let $C_0(G, A^*)$ be the continuous functions from G into A^* vanishing at infinity and let $C_{00}(G, A^*)$ be the continuous functions from G into A^* with compact support under the norm $\|f\| = \sup_{t \in G} \|f(t)\|$. For $f \in L^\infty(G, A^*)$, set $L_x f(t) = f(xt)$ ($x, t \in G$). Then f is called left uniformly continuous, if the map $x \mapsto L_x f$ from G into $L^\infty(G, A^*)$ is continuous with respect to $N_\infty(\|f\|)$ on $L^\infty(G, A^*)$. The set of uniformly continuous functions is denoted by $LUC(G, A^*)$. A function $f \in C(G, A^*)$ is called weakly almost periodic if the set $\{L_x f : x \in G\}$ is relatively compact in the weak-topology on $C(G, A^*)$. The set of weakly almost periodic functions are denoted by $WAP(G, A^*)$. In the case $A = \mathbb{C}$, the complex field, these spaces are denoted by $L^1(G)$, $M(G)$, $C(G)$, $C_0(G)$, $C_{00}(G)$, $LUC(G)$ and $WAP(G)$. For general terms in vector-valued functions, we follow [5].

The left invariant means on spaces of vector-valued functions were first considered by Dixmier in [3]. In this work, we set up a relation between a vector-valued mean and a scalar-valued mean, by which we will be able to translate many important results developed in the classic theory. We also present some of the properties of left invariant means on $LUC(G, A^*)$, $WAP(G, A^*)$ and $C_0(G, A^*)$. Our references for the vector-valued integration theory are [1], [2] and [5].

2. MAIN RESULTS

Definition 2.1. Let A be a Banach algebra and X be a subspace of $L^\infty(G, A^*)$. A map $M : L^\infty(G, A^*) \rightarrow A^*$ is called a mean on X if

- (i) M is linear;
- (ii) For each $f \in X$, $M(f)$ belongs to the set $\overline{\text{co}\{f(x) : x \in G\}}$, where

closure is taken in the weak*-topology, and coX denotes the convex hull of a set X .

If X is also left invariant, then M is called left invariant if $M(L_a f) = M(f)$ for each $a \in G$ and $f \in X$. Dixmier showed in [3] that if m is a left invariant mean on $L^\infty(G)$. Then m induces a left invariant mean M on $L^\infty(G, A^*)$ such that $\langle M(f), a \rangle = m(\langle f(\cdot), a \rangle)$ for each $a \in A$, here $\langle f(\cdot), a \rangle$ denotes the functions $x \mapsto \langle f(x), a \rangle$.

Let A and B be two Banach algebras, and let A be a closed ideal in B . For each $a \in A$, put $\rho_a(b) := \|ba\| + \|ab\|$ ($b \in B$), then ρ_a is a seminorm. The topology defined on B by these seminorms is called the strict topology. We write B^{*st} for the dual of B with respect to the strict topology.

Theorem 2.2. *Let G be a locally compact group. Then G is amenable if and only if $L^1(G, A)$ is amenable for each unital separable Banach algebra A .*

Proof. Suppose that $L^1(G, A)$ is amenable for each unital separable Banach algebra A . Consider $A = \mathbb{C}$. Then G is amenable [7]. Conversely, let X be a Banach $L^1(G, A)$ -bimodule, and let $D : L^1(G, A) \rightarrow X^*$ be a continuous derivation. By Proposition 8.1 in [4], $L^1(G, A)$ has a bounded approximate identity. Then there is no loss of generality if we suppose that X is pseudo-unital. By Proposition 2.1.6 in [12], there is a unique $\bar{D} : M(G, A) \rightarrow X^*$ that extends D and is continuous with respect to the strict topology on $M(G, A)$, and the weak*-topology on X^* .

We can embed G into $M(G, A)$. The map $\delta_x : G \rightarrow M(G, A)$ given by $\delta_x(H) = \chi_H(x)e_A$ for each $x \in G$ and $H \subseteq G$ is the required embedding. We claim that $\Delta = \{\delta_x : x \in G\}$ is dense in $M(G, A)$ with respect to the strict topology. We assume to the contrary that μ is not in $\bar{\Delta}$ where closure is taken in the strict topology, thus there are some $f \in M(G, A)^{*st}$, such that $\langle f, \mu \rangle = 1$, $\langle f, \delta_x \rangle = 0$ for each $x \in G$. By Proposition 23.18 and Proposition 23.33 in [2], the map $x \mapsto \delta_x$ is continuous with respect to the strict topology. This implies that $\langle f, \mu \rangle = \int \langle f, \delta_x \rangle d\mu(x) = 0$ [11].

It suffices to show that $\bar{\Delta} |_\Delta$ is inner. Because, if $\mu \in M(G, A)$, there will be a net $\{\delta_{x_\alpha}\}_{\alpha \in I}$ in Δ with $\delta_{x_\alpha} \rightarrow \mu$ in the strict topology. So $\bar{D}(\delta_{x_\alpha}) \rightarrow \bar{D}(\mu)$ in the weak*-topology. But, there is $\beta^* \in X^*$ such that $\bar{D}(\mu) = w_k^* - \lim_\alpha (\beta^* \cdot \delta_{x_\alpha} - \delta_{x_\alpha} \cdot \beta^*)$. Now let $\beta \in X$. Since X is pseudo-unital Banach $L^1(G, A)$ -bimodule, then there are $f_1, f_2 \in L^1(G, A)$

and $\hat{\beta} \in X$ such that $\beta = f_1 \cdot \hat{\beta} \cdot f_2$. Hence:

$$\begin{aligned} \langle \beta^* \cdot \delta_{x_\alpha} - \delta_{x_\alpha} \cdot \beta^*, \beta \rangle &= \langle \beta^* \cdot \delta_{x_\alpha} - \delta_{x_\alpha} \cdot \beta^*, f_1 \cdot \hat{\beta} \cdot f_2 \rangle \\ &\rightarrow \langle \beta^* \cdot \mu - \mu \cdot \beta^*, \beta \rangle. \end{aligned}$$

Consequently:

$$\overline{D}(\mu) = \beta^* \cdot \mu - \mu \cdot \beta^*.$$

Consider the constant function $1 \in L^\infty(G)$. There is a function $f_0 : G \rightarrow A^*$ by $f_0(x) = \langle 1, x \rangle a^*$ ($a^* \in A^*$, $x \in G$). More generally, each function $g \in L^\infty(G)$ has the form $\langle f(\cdot), a \rangle$ with $a \neq 0$, $a^* \in A^*$, $\langle a^*, a \rangle = 1$ and $f(x) = \langle g, x \rangle a^*$. For each $\beta \in X$, define $\langle \Lambda_\beta, \mu \rangle = \langle \overline{D}(\mu) \mu, \beta \rangle$ ($\mu \in M(G, A)$). Put $\Lambda_{\hat{\beta}} = \Lambda_\beta|_\Delta$. Now define $\langle \rho_\beta, x \rangle = \langle \Lambda_{\hat{\beta}}, \delta_x \rangle = \langle \overline{D}(\delta_x) \cdot \delta_x, \beta \rangle$ ($x \in G$). Since $\rho_\beta \in L^\infty(G)$, then there are some $f_\beta \in L^\infty(G, A^*)$ such that $f_\beta(x) = \rho_\beta(x) a^*$ ($a^* \in A^*$, $x \in G$). By assumption, there is $m \in L^\infty(G)^*$ such that $\langle m, 1 \rangle = 1$, $\langle m, L_x f \rangle = \langle m, f \rangle$ for each $f \in L^\infty(G)$ and $x \in G$. Thus m induces a left invariant mean M on $L^\infty(G, A^*)$. Now put $\langle \lambda, \beta \rangle = \langle M, f_\beta \rangle$. The space X becomes a Banach Δ -module via $\delta_x \square \beta = \beta$, $\beta \square \delta_x = \delta_x \cdot \beta$. δ_x ($x \in G$, $\beta \in X$). Then dual actions on X^* are given by $\beta^* \square \delta_x = \beta^*$, $\delta_x \square \beta^* = \delta_x \cdot \beta^*$.

Define $\overline{D}_0(\delta_x) = \overline{D}(\delta_x) \cdot \delta_x$. It is routinely checked that \overline{D}_0 is a derivation, and \overline{D} is inner if and only if \overline{D}_0 is inner for this new module structure. It remains to be shown that \overline{D}_0 is inner. For $\delta_{x_0} \in M(G, A)$,

$$\begin{aligned} \langle \delta_{x_0} \square \lambda - \lambda \square \delta_{x_0}, \beta \rangle &= \langle \lambda, \beta \square \delta_{x_0} - \delta_{x_0} \square \beta \rangle \\ &= \langle \lambda, \beta \square \delta_{x_0} - \beta \rangle = \langle M, f_{\beta \square \delta_{x_0} - \beta} \rangle. \end{aligned}$$

Furthermore, we have

$$\begin{aligned} \langle f_{\beta \square \delta_{x_0} - \beta}, x \rangle &= \langle \rho_{\beta \square \delta_{x_0} - \beta}, x \rangle a^* = \overline{D}(\delta_x) \cdot \delta_x (\beta \square \delta_{x_0} - \beta) a^* \\ &= \overline{D}_0(\delta_x) (\beta \square \delta_{x_0} - \beta) a^* = \langle \delta_{x_0} \square \overline{D}_0(\delta_x) - \overline{D}_0(\delta_x), \beta \rangle a^* \\ &= \langle \overline{D}_0(\delta_{x_0} \delta_x) - \overline{D}_0(\delta_{x_0}) \square \delta_x - \overline{D}_0(\delta_x), \beta \rangle a^* \\ &= \langle \overline{D}_0(\delta_{x_0} \delta_x), \beta \rangle a^* - \langle \overline{D}_0(\delta_{x_0}), \beta \rangle a^* - \rho_\beta(x) a^* \\ &= \langle \rho_\beta, x_0 x \rangle a^* - \langle \overline{D}_0(\delta_{x_0}), \beta \rangle \langle 1, x \rangle a^* - \rho_\beta(x) a^* \\ &= f_\beta(x_0 x) - \langle \overline{D}_0(\delta_{x_0}), \beta \rangle f_0(x) - f_\beta(x), \end{aligned}$$

then $\langle \delta_{x_0} \square \lambda - \lambda \square \delta_{x_0}, \beta \rangle = -\langle \overline{D}_0(\delta_{x_0}), \beta \rangle$. It follows that $\overline{D}_0(\delta_{x_0}) = \delta_{x_0} \square \lambda_0 - \lambda_0 \square \delta_{x_0}$ where $\lambda_0 = -\lambda$. Thus \overline{D}_0 is inner. \square

Theorem 2.3. *Let G be a locally compact group. Then the following statements are equivalent:*

- (i) G is amenable.

- (ii) For every unital separable Banach algebra A , there exists a bounded net $\{\psi_\alpha\}_{\alpha \in I} \subseteq L^1(G, A)$ such that $\|\delta_x * \psi_\alpha - \psi_\alpha\|_1 \rightarrow 0$ whenever $x \in G$.
- (iii) For every unital separable Banach algebra A , there exists a bounded net $\{\psi_\alpha\}_{\alpha \in I} \subseteq L^1(G, A)$ such that for every compact set $K \subseteq G$, $\|\psi * \psi_\alpha - \psi_\alpha\|_1 \rightarrow 0$ uniformly for all $\psi \in L^1(G, A)$ with $\int_{G \setminus K} \|\psi(t)\| dm(t) = 0$.

Proof. Consider $A = \mathbb{C}$. Then Theorem 6.7 in [10] yields (ii) \implies (i), (iii) \implies (i).

(i) \implies (ii) Suppose that G is amenable. Let m be a left invariant mean on $L^\infty(G)$, and M be an induced mean on $L^\infty(G, A^*)$. Choose $a \in A$ such that, $\|a\| = 1$. Define $\langle \Gamma_a, f \rangle = \langle M(f), a \rangle$ for each $f \in L^\infty(G, A^*)$. Regards Γ_a as an element of $L^\infty(G, A^*)^*$. The rest of the proof is essentially the same as the Lemma 6.3 in [10].

(i) \implies (iii) This is just a re-statement of Proposition 6.7 in [10]. \square

Let $\{e_\alpha\}_\alpha$ be a bounded approximate identity for $L^1(G)$ and e_A be an identity in A . By Proposition 8.1 in [4], $\{e_\alpha \otimes e_A\}_\alpha$ is a bounded approximate identity for $L^1(G) \hat{\otimes} A$ where $\hat{\otimes}$ denotes the completion of usual tensor product of Banach spaces with respect to the projective tensor norm. We consider $\{e_\alpha \otimes e_A\}_\alpha$ as an element in $(L^1(G) \hat{\otimes} A)^{**}$ and $F \in (L^1(G) \hat{\otimes} A)^*$. Using exactly the same notation as in [6], we put $\langle (e_\alpha \otimes e_A), F \rangle = \int Fd(e_\alpha \otimes e_A)$. Given a dual Banach space X^* and $F \in B(L^1(G), A; X^*)$, we define $\int Fd(e_\alpha \otimes e_A) \in X^*$ by

$$\left\langle \int Fd(e_\alpha \otimes e_A), x \right\rangle = \int \langle F(f, a), x \rangle d(e_\alpha \otimes e_A)(f, a),$$

where $f \in L^1(G)$, $a \in A$ and $x \in X$.

Theorem 2.4. *Let G be a locally compact group and let A be a unital separable Banach algebra. Then $L^1(G, A)$ is approximately weakly amenable.*

Proof. Let $D : L^1(G, A) \rightarrow L^1(G, A)^*$ be a continuous derivation. It is well-known that the space $L^1(G, A)$ is isometrically isomorphic to $L^1(G) \hat{\otimes} A$. Therefore we define $F : L^1(G) \times A \rightarrow L^1(G, A)^*$ by $F(f, a) = D(f \otimes a)$. Put, $g_\alpha = \int F(f, a)d(e_\alpha \otimes e_A)(f, a)$. We know that $L^1(G, A)^* \cong L^\infty(G, A^*)$. Then for each $F(f, a) \in L^1(G, A)^*$, its image under isometry onto $L^\infty(G, A^*)$ is a map whose values at $x \in G$ is $F(f, a)(x) = \langle D(f \otimes a), x \rangle$. We put $\overline{F}_x : L^1(G) \times A \rightarrow A^*$ with $\overline{F}_x(f, a) = F(f, a)(x)$, $f \in L^1(G)$, $a \in A$, and $x \in G$. Note that $\overline{F}_x \in B(L^1(G), A; A^*)$. By the above argument, we define

$\int \overline{F}_x(f, a)d(e_\alpha \otimes e_A)(f, a) \in A^*$ by

$$\left\langle \int \overline{F}_x(f, a)d(e_\alpha \otimes e_A)(f, a), c \right\rangle = \int \langle \overline{F}_x(f, a), c \rangle d(e_\alpha \otimes e_A)(f, a), (c \in A).$$

The map $x \mapsto g_\alpha(x) = \int F(f, a)(x)d(e_\alpha \otimes e_A)(f, a)$ is a scalarwise measurable function, and $N_\infty(\|g_\alpha(x)\|) < \infty$ for each α . Then, by Theorem 8.18.2 in [5], there is a map κ_{g_α} in $B(A, L^\infty(G))$ such that, $\langle \kappa_{g_\alpha}(a), f \rangle = \int f(x)\langle g_\alpha(x), a \rangle dm(x)$ for each $f \in L^1(G)$, and $a \in A$, where κ_{g_α} is defined by $\kappa_{g_\alpha}(a) = \langle g_\alpha(x), a \rangle$.

Using the same notation as in [6], we have $e_\alpha \otimes e_A = \int f \otimes g d(e_\alpha \otimes e_A)(f, g)$. Moreover, $e_\alpha \otimes e_A$ is a bounded approximate identity for $L^1(G) \hat{\otimes} A$. Therefore, for each $F : L^1(G) \times A \rightarrow L^1(G, A)^*$, $f, g \in L^1(G)$, and $a, b \in A$, we have

$$\begin{aligned} \lim_\alpha \int F(fg, ab)d(e_\alpha \otimes e_A)(f, a) &= \lim_\alpha \left\langle \int (fg \otimes ab)d(e_\alpha \otimes e_A)(f, a), F \right\rangle \\ &= \lim_\alpha \left\langle \int (gf \otimes ba)d(e_\alpha \otimes e_A)(f, a), F \right\rangle \\ &= \lim_\alpha \int F(gf, ba)d(e_\alpha \otimes e_A)(f, a). \end{aligned}$$

Hence

$$\begin{aligned} &\lim_\alpha (g \otimes b) \langle f, \kappa_{g_\alpha}(\acute{a}) \rangle \\ &= \lim_\alpha \int \acute{f}(x) \langle (g \otimes b) \cdot g_\alpha(x), \acute{a} \rangle dm(x) \\ &= \lim_\alpha \int \acute{f}(x) \left\langle \int (g \otimes b) \cdot D(f \otimes a)(x) d(e_\alpha \otimes e_A)(f, a), \acute{a} \right\rangle dm(x) \\ &= \lim_\alpha \int \acute{f}(x) \left\langle \int D(gf \otimes ba)(x) d(e_\alpha \otimes e_A)(f, a), \acute{a} \right\rangle dm(x) \\ &= \lim_\alpha \int \acute{f}(x) \left\langle \int D(g \otimes b) \cdot (f \otimes a)(x) d(e_\alpha \otimes e_A)(f, a), \acute{a} \right\rangle dm(x) \\ &= \int \acute{f}(x) \lim_\alpha \left\langle \int F(gf, ba)(x) d(e_\alpha \otimes e_A)(f, a), \acute{a} \right\rangle dm(x) \\ &= \int \acute{f}(x) \lim_\alpha \left\langle D(g \otimes b) \int (f \otimes a)(x) d(e_\alpha \otimes e_A)(f, a), \acute{a} \right\rangle dm(x) \\ &= \int \acute{f}(x) \lim_\alpha \left\langle \int F(fg, ab)(x) d(e_\alpha \otimes e_A)(f, a), \acute{a} \right\rangle dm(x) \\ &= \int \acute{f}(x) \langle D(g \otimes b)(x), \acute{a} \rangle dm(x) \end{aligned}$$

$$\begin{aligned}
 &= \int \acute{f}(x) \lim_{\alpha} \langle \int F(f, a)(x) d(e_{\alpha} \otimes e_A)(f, a), \acute{a} \rangle dm(x) (g \otimes b) \\
 &- \int \acute{f}(x) \langle D(g \otimes b)(x), \acute{a} \rangle dm(x) \\
 &= \lim_{\alpha} \int \acute{f}(x) \langle g_{\alpha}(x), \acute{a} \rangle dm(x) (g \otimes b) - \int \acute{f}(x) \langle D(g \otimes b)(x), \acute{a} \rangle dm(x) \\
 &= \lim_{\alpha} \langle \acute{f}, \kappa_{g_{\alpha}}(\acute{a}) \rangle (g \otimes b) - \langle \acute{f}, \kappa_{D(g \otimes b)}(\acute{a}) \rangle
 \end{aligned}$$

for all $g \otimes b \in L^1(G) \otimes A$, $\acute{a} \in A$ and $\acute{f} \in L^1(G, A)$. Consequently,

$$\begin{aligned}
 \lim_{\alpha} ((g \otimes b) \kappa_{g_{\alpha}}(\acute{a}) - \kappa_{g_{\alpha}}(\acute{a})(g \otimes b)) &= -\kappa_{D(g \otimes b)}(\acute{a}) \\
 \lim_{\alpha} \langle (g \otimes b) \cdot g_{\alpha}(x), \acute{a} \rangle - \langle g_{\alpha}(x) \cdot (g \otimes b), \acute{a} \rangle &= -\langle D(g \otimes b)(x), \acute{a} \rangle
 \end{aligned}$$

for all $g \otimes b \in L^1(G) \otimes A$ and $\acute{a} \in A$. It follows that

$$D(g \otimes b) = \lim_{\alpha} ((g \otimes b) \cdot g'_{\alpha} - g'_{\alpha} \cdot (g \otimes b))$$

for all $g \otimes b \in L^1(G) \otimes A$, where $g'_{\alpha} = -g_{\alpha}$. □

It is known that G is amenable if and only if $LUC(G)$ has a left invariant mean. It is interesting to have a direct proof of this fact. We present a vector version of this characterization.

Theorem 2.5. *Let G be a locally compact group, and let A be a unital separable Banach algebra. Then:*

- (i) $L^{\infty}(G, A^*)L^1(G, A) = LUC(G, A^*)$.
- (ii) G is amenable if and only if $LUC(G, A^*)$ has a left invariant mean.

Proof. (i) Let $f \in LUC(G, A^*)$. Let $\{U_{\alpha}\}$ be a net of neighborhoods of e directed downwards. Let $\{\nu_{\alpha}\}_{\alpha \in I}$ be an approximate identity of norm 1 in $L^1(G, A)$ such that $supp \nu_{\alpha} \subseteq U_{\alpha}$. Given $\epsilon > 0$, there exists $\alpha_0 \in I$ such that for each $\alpha \geq \alpha_0$ and $y \in U_{\alpha}$, $N_{\infty}(\|f_y - f\|) < \epsilon$. Then

$$\begin{aligned}
 &|\langle f \cdot \nu_{\alpha}, \mu \rangle - \langle f, \mu \rangle| \\
 &= |\langle f, \nu_{\alpha} * \mu \rangle - \langle f, \mu \rangle| \\
 &= \left| \int \langle f(t), \nu_{\alpha} * \mu(t) \rangle dm(t) - \int \langle f(t), \mu(t) \rangle dm(t) \right| \\
 &= \left| \int \langle f(t), \int \mu(y^{-1}t) d\nu_{\alpha}(y) \rangle dm(t) - \int \langle f(t), \mu(t) \rangle dm(t) \right| \\
 &= \left| \int \langle f(yt), \int \mu(t) d\nu_{\alpha}(y) \rangle dm(t) - \int \langle f(t), \mu(t) \rangle dm(t) \right| \\
 &\leq N_{\infty}(\|f_y - f\|) \|\mu\| \|\nu_{\alpha}\|_1.
 \end{aligned}$$

On the other hand, $f.\nu_\alpha \in L^\infty(G, A^*)L^1(G, A)$, and $L^\infty(G, A^*)L^1(G, A)$ is a Banach space. It follows that $f \in L^\infty(G, A^*)L^1(G, A)$. It is easy to see that $L^\infty(G, A^*)L^1(G, A) \subseteq LUC(G, A^*)$, and so $L^\infty(G, A^*)L^1(G, A) = LUC(G, A^*)$.

(ii) This follows from (i) and the proof of Lemma 3.2 in [9]. \square

Let A be a Banach algebra. Recall that a functional $f \in A^*$ for which $\{f.a : \|a\| \leq 1, a \in A\}$ is relatively compact in the weak-topology of A^* is said to be weakly almost periodic. The set of weakly almost periodic functionals on A is denoted by $WAP(A)$ (see [8]). It is known that $WAP(G) = WAP(L^1(G))$ [13]. Note that for $f \in L^\infty(G, A^*)$ and $\mu \in M(G, A)$, we define $\langle f\mu, \nu \rangle = \langle f, \mu * \nu \rangle$ for every $\nu \in L^1(G, A)$.

Theorem 2.6. *Let G be a locally compact group, and A be a unital separable Banach algebra. Then:*

- (i) *If $f \in L^\infty(G, A^*)$, then $f \in WAP(L^1(G, A))$ if and only if $\{f\delta_x : x \in G\}$ is relatively weakly compact in $L^\infty(G, A^*)$.*
- (ii) *$WAP(L^1(G, A)) = WAP(G, A^*)$.*
- (iii) *$WAP(L^1(G, A))$ has a left invariant mean.*

Proof. (i) Let $f \in WAP(L^1(G, A))$. It is known that $C_0(G, A)^* = M(G, A^*)$ [2]. Now consider $\delta_x \in M(G, A^*)$. By the Hahn Banach Theorem, we may assume that $m \in L^\infty(G, A^*)^*$ is an extension of δ_x with norm one. Then there is a net $\{\mu_\alpha\}_{\alpha \in I}$ in $L^1(G, A)$ with $\|\mu_\alpha\| \leq 1$ such that $\mu_\alpha \rightarrow m$ in the weak*-topology. Hence, for every $\phi \in L^1(G, A)$,

$$\langle \phi.f, \mu_\alpha \rangle \rightarrow \langle \phi.f, m \rangle.$$

Since $\{f.\mu : \|\mu\| \leq 1, \mu \in L^1(G, A)\}$ is relatively weakly compact, then there is an element $g \in L^\infty(G, A^*)$ and a subnet $\{\mu_\beta\}_{\beta \in I}$ of $\{\mu_\alpha\}_{\alpha \in I}$ such that $f.\mu_\beta \rightarrow g$ in the weak-topology. On the other hand, $\langle m, \phi.f \rangle = \langle \delta_x, \phi.f \rangle = \langle f\delta_x, \phi \rangle$, and so $g = f\delta_x$. Thus the set $\{f\delta_x : x \in G\}$ is contained in $\overline{\{f.\mu : \|\mu\| \leq 1, \mu \in L^1(G, A)\}}$, where closure is taken in the weak-topology, and the compactness of $\overline{\{f\delta_x : x \in G\}}$ follows from the compactness of $\overline{\{f.\mu : \|\mu\| \leq 1\}}$.

Conversely, suppose that $f \in L^\infty(G, A^*)$ and $\mu \in L^1(G, A)$ with $\|\mu\| \leq 1$, and $\{f\delta_x : x \in G\}$ is relatively weakly compact. By the Krein-Smulian Theorem, $co\{f\delta_x : x \in G\}$ is relatively weakly compact. We claim that

$$f\mu \in \overline{co\{f\delta_x : x \in G\}},$$

where closure is taken in the weak-topology. We assume to the contrary that $f\mu$ is not in $\overline{co\{f\delta_x : x \in G\}}$. By the Hahn Banach Theorem,

there exists $F \in L^\infty(G, A^*)^*$, such that

$$Re|\langle F, f\mu \rangle| \geq \gamma_1 > \gamma_2 > Re|\langle F, f\delta_x \rangle|,$$

where $\gamma_1, \gamma_2 \in \mathbb{R}$, and $x \in G$. By Theorem 8.14.8 in [5], the integral $\int f\delta_x d\mu$ belongs to $L^\infty(G, A^*)$. Moreover, for any $\nu \in L^1(G, A)$, $\langle \int f\delta_x d\mu, \nu \rangle = \int \langle f.\delta_x, \nu \rangle d\mu = \int \langle f, \delta_x * \nu \rangle d\mu = \langle f, \mu * \nu \rangle = \langle f\mu, \nu \rangle$ (see Chapter 3 in [11]). By Theorem 3.28 in [11], we have

$$|\langle F, f\mu \rangle| = \left| \int \langle F, f\delta_x \rangle d\mu \right| \leq \int |\langle F, f\delta_x \rangle| d|\mu|(x) < Re\langle F, f\mu \rangle.$$

This is a contradiction, and so $\{f.\mu : \|\mu\| \leq 1, \mu \in L^1(G, A)\}$ is contained in the closure of $co\{f\delta_x : x \in G\}$, and the compactness of $\overline{\{f.\mu : \|\mu\| \leq 1, \mu \in L^1(G, A)\}}$ follows from the compactness of $co\{f\delta_x : x \in G\}$. Consequently, $f \in WAP(L^1(G, A))$, and the proof is complete.

(ii) Let $f \in WAP(G, A^*)$. Then the set $\{L_x f : x \in G\}$ is relatively weakly compact in $C(G, A^*)$. Note that for each $x \in G$, we have $\langle f\delta_x, \mu \rangle = \langle f, \delta_x * \mu \rangle = \int \langle f(t), \int \mu(y^{-1}t) d\delta_x(t) \rangle dm(t) = \langle L_x f, \mu \rangle$. Then, by the Eberlien-Smulian Theorem, $\{f\delta_x : x \in G\}$ is relatively weakly compact in $L^\infty(G, A^*)$, and from (i), $f \in WAP(L^1(G, A))$.

Conversely, let $f \in WAP(L^1(G, A))$. The map $x \mapsto f\delta_x$ is continuous with respect to the weak*-topology. From (i), $\{f\delta_x : x \in G\}$ is relatively weakly compact. Then $x \rightarrow f\delta_x$ is continuous with respect to the weak-topology. Now, let $\{U_\alpha\}$ be a net of neighborhood of e directed downwards. Let $\{\nu_\alpha\}_{\alpha \in I}$ be an approximate identity of norm 1 in $L^1(G, A)$ such that $supp\nu_\alpha \subseteq U_\alpha$. Given $\epsilon > 0$ and $F \in L^\infty(G, A^*)^*$, there exists an α_0 such that for each $\alpha \geq \alpha_0$ and $x \in U_\alpha$, $|\langle F, f\delta_x \rangle - \langle F, f \rangle| < \epsilon$, and so $|\int \langle F, f\delta_x \rangle d\nu_\alpha - \langle F, f \rangle| < \epsilon$. By Theorem 8.14.8 in [5] and the Krein-Smulian Theorem, $\int f\delta_x d\nu_\alpha \in L^\infty(G, A^*)$. Moreover, for any $\mu \in L^1(G, A)$, $\langle \int f\delta_x d\nu_\alpha(x), \mu \rangle = \langle f\nu_\alpha, \mu \rangle$. Therefore, for each $\alpha \geq \alpha_0$,

$$\begin{aligned} |\langle F, f\nu_\alpha \rangle - \langle F, f \rangle| &= \left| \langle F, \int f\delta_x d\nu_\alpha(x) \rangle - \langle F, f \rangle \right| \\ &= \left| \int \langle F, f\delta_x \rangle d\nu_\alpha - \langle F, f \rangle \right| < \epsilon. \end{aligned}$$

So $f\nu_\alpha \rightarrow f$ in the weak-topology. An argument similar to that in the proof of Lemma 6.3 in [10] shows that we can find a bounded net $\{\nu_\alpha\}_{\alpha \in I}$ consisting of convex combination of elements in $\{\nu_\alpha\}_{\alpha \in I}$ such that $f\nu_\alpha \rightarrow f$ in the norm topology. In addition, $L^\infty(G, A^*)L^1(G, A)$ is a Banach space. Then from Theorem 2.5, $f \in C(G, A^*)$. Since

$\{f\delta_x : x \in G\}$ is relatively weakly compact, then by the Eberlian-Smulian Theorem, $\{L_x f : x \in G\}$ is relatively weakly compact in $C(G, A^*)$. Hence, $f \in WAP(G, A^*)$.

(iii) It is an immediate consequence of Theorem 4.3 in [14] and (ii). \square

In the next theorem, we present an interesting property about a left invariant mean when considered on $C_0(G, A^*)$. Analogous to the scalar function case, we can easily obtain the following theorem.

Theorem 2.7. *Let G be a non-compact amenable group and let $f \in C_0(G, A^*)$. If M is left invariant mean on $L^\infty(G, A^*)$, then $|M(f)| = 0$.*

Proof. Let M be a left invariant mean on $L^\infty(G, A^*)$ and $f \in L^\infty(G, A^*)$. Then the set $\{f(t) : \|f(t)\| \leq \|f\|_\infty\}$ is weak*-closed in A^* and $M(f) \in \overline{\{f(x) : x \in G\}}$, where closure is taken in the weak*-topology. It follows that $|M(f)| \leq \|f\|_\infty$. Using the Urysohn Lemma, it is easy to see that $\overline{C_{00}(G, A^*)}^{\|\cdot\|} = C_0(G, A^*)$. Then it is enough to prove that the result in the case where $f \in C_{00}(G, A^*)$. Let $K = \text{supp} f$. There exists an infinite sequence $\{a_n\}_{n \in \mathbb{N}}$ in G such that $a_0 = e$, and $(a_i K) \cap (a_j K) = \emptyset$, whenever $i, j \in \mathbb{N}, i \neq j$ [10]. Put $g_n = \sum_{i=0}^n L_{a_i} f$ ($n \in \mathbb{N}$). For any $n \in \mathbb{N}$,

$$|M(g_n)| = |nM(f)| \leq \|g_n\|_\infty = \|f\|_\infty.$$

Consequently, $M(f) = 0$. \square

Acknowledgments

The authors would like to thank the referee for his/her careful reading of the paper and many valuable suggestions.

REFERENCES

1. J. Diestel and J. J. Uhl, Jr., Vector measures, Amer. Math. Soc. 1977.
2. N. Dinculeanu, Integration on locally compact spaces, Noorhoff International Publishing, Leyden 1974.
3. J. Dixmier, Les moyennes invariantes dans les semi-groupes et leur applications, *Acta Sci. Math. (Szege)* **12** (1950), 213-227.
4. R. S. Doran, J. Wichmann, Approximate identities and factorization in Banach modules, Lecture Notes in Mathematics, 768, Springer Verlag, 1979.
5. R. E. Edwards, Functional analysis, Theory and applications, Holt, Rinehart and Winston, New York, 1965.
6. E. Effros, Amenability and Virtual Diagonals for von Neumann algebras, *J. Funct. Anal.* **78** (1988), 137-153.

7. B. E. Johnson, Cohomology in Banach algebras, *Mem. Amer. Math. Soc.* **127** (1972).
8. A. Ghaffari, Strongly and weakly almost periodic linear maps on semigroup algebras, *Semigroup Forum* **76** (2008), 95-106.
9. A. Ghaffari and S. Javadi, φ -means of some Banach subspaces on a Banach algebras, *Proceedings of the Romanian Academy* **13** (2012), 302-309.
10. J. P. Pier, Amenable locally compact groups, John Wiley And Sons New York, 1984.
11. W. Rudin, Functional analysis, McGraw Hill New York, 1991.
12. V. Runde, Lectures on Amenability, Lecture Notes in Mathematics 1774, Springer Verlag, 2002.
13. A. Ulgar, Continuity of weakly almost periodic functionals on $L^1(G)$, *Quart. J. Math. Oxford Ser.* **237** (1986), 495-497.
14. C. Zhang, Vector-valued means and weakly almost periodic functions, *Intenat. J. Math.* **17** (1994), 227-238.

A. Ghaffari

Department of Mathematics, University of Semnan, P.O.Box 35195-363, Semnan, Iran.

Email: aghaffari@semnan.ac.ir

S. Javadi

Faculty of Engineering- East Guilan, University of Guilan, P.O.Box 44891-63157, Rudsar, Iran.

Email: s.javadi62@gmail.com

AMENABILITY OF VECTOR VALUED GROUP ALGEBRAS

JAVADI S. AND GHAFARI A.

میانگین پذیری جبرهای باناخ برداری مقدار

علی غفاری - سمانه جوادی
دانشگاه سمنان - دانشکده ریاضی دانشگاه گیلان - دانشکده فنی شرق گیلان

هدف این مقاله توسعه مفهوم میانگین پذیری جبرهای گروهی برداری مقدار است. نشان می دهیم اگر A جبر باناخ جدایی پذیر یکدار باشد، آنگاه جبر گروهی $L^1(G, A)$ میانگین پذیر ضعیف تقریبی است. همچنین شرایط لازم و کافی برای وجود میانگین پایای چپ روی $LUC(G, A^*)$, $WAP(G, A^*)$ و $C_0(G, A^*)$ را بررسی می کنیم.

کلمات کلیدی: میانگین پذیری، جبر باناخ، مشتق، جبر گروهی، میانگین پایا.