

Analysis of *IL-33* Gene Polymorphisms (rs1157505C/G and rs11792633C/T) and the Risk of Tuberculosis in Southeastern Iran

Tayebeh Baranzehi¹, Shahrbanou Karimi-Davani² and Dor Mohammad Kordi-Tamandani^{1*}

¹ Department of Biology, University of Sistan and Baluchestan, Zahedan, Iran

² South Iranian Genetic Medical Center, Shiraz University of Medical Sciences, Shiraz, Iran

ARTICLE INFO

Article history:

Received 20 December 2019

Accepted 22 January 2020

Available online 02 February 2020

Keywords:

Bacillus

IL-33

Mycobacterium

rs1157505 C/G

rs11792633 C/T

Tuberculosis

Corresponding author:

✉ D.M. Kordi-Tamandani

dor_kordi@science.usb.ac.ir

p-ISSN 2423-4257

e-ISSN 2588-2589

ABSTRACT

Tuberculosis is a vagarious infectious disease that generally affects the lungs. Accordingly, in some cases, it can also affect the liver and kidney. Host genetic may affect tuberculosis caused by bacillus *Mycobacterium tuberculosis*. The main risk factors for the disease are a weakened immune system because of diabetes, some cancers, HIV/AIDS, severe kidney disease, cancer treatment, and malnutrition. *IL-33* is involved in the activation of eosinophils, mast cells, basophils, and natural killer cells and the maturation of T helper type2 cells. The developments of CD4 (+) TH1 and CD8 (+) T cell responses are involved in protection against TB, *IL-33* promotes the development of these cells. The purpose of the present research was to investigate the association between *Mycobacterium tuberculosis* infection and *IL-33* gene polymorphisms (rs1157505C/G and rs11792633C/T) with tuberculosis in the cases and controls from the area of high tuberculosis prevalence in Iran. In this study, 100 patients with tuberculosis disease and 91 healthy controls were included. Polymorphisms of the *IL-33* gene were genotyped using T-ARMS-PCR. The analysis of the haplotype combinations among *IL-33* polymorphisms demonstrated that the magnitude of the association was higher for the combined CC/CT genotypes. The CT genotype related to C/T polymorphism of the *IL-33* gene increased the risk of tuberculosis. The combined CG+GG genotypes related to C/G polymorphism of the *IL-33* gene also increased the risk of tuberculosis, but the difference was not statistically significant. In conclusion, *IL-33* gene polymorphisms may be considered as important contributors to tuberculosis in Iran.

© 2015 UMZ. All rights reserved.

Please cite this paper as: Baranzehi T, Karimi-Davani S, Kordi-Tamandani DM. 2020. Analysis of *IL-33* gene polymorphisms (rs1157505C/G and rs11792633C/T) and the risk of tuberculosis in Southeastern Iran. *J Genet Resour* 6(1): 54-59. doi: 10.22080/jgr.2020.17777.1166.

Introduction

Mycobacterium tuberculosis is the main cause of tuberculosis, a disease that affects different organs and spreads through breathing between different people (Flynn and Chan, 2001; Moosazadeh *et al.*, 2014). Tuberculosis usually influences the lungs, but may also affect other parts of the body EPTB (Raja, 2004). It was established that the people infected with HIV are more likely vulnerable to TB due to the weakness of the immune system (Getahun *et al.*, 2010). The prevalence of Tuberculosis is higher

in adult males than females (Zumla *et al.*, 2013). Tuberculosis is one of the most important health problems in Iran. It has been previously reported that the human host exerted a complex response to TB. Immune system cells play a pivotal role in *Mycobacterium tuberculosis*. Zumla *et al.* also reported the association between TB, and the strong pro-inflammatory immune responses, and the harmful activity of pro-inflammatory cytokines. In humans, 'hyperinflammatory' reaction is a sign of TB progression (Zumla *et al.*, 2013). Interleukin 33, a 30 kDa protein, is the latest member of the IL-1 superfamily. The

Production and expression of IL-33 from immune cells such as mast cells, macrophages, and dendritic cells resulted in the production of cytokines and chemokines, including IL-13, IL-8, and IL-6. Non-immune cells, including smooth muscle, endothelial, epithelial cells, and fibroblasts, also produced this cytokine (Yagami *et al.*, 2010). IL-33 is mostly expressed in the Skin, gut, and lung, and expression is enriched in the brain and spinal cord (Schmitz *et al.*, 2005). Its gene is located on the short arm of chromosome 9 (9p24). The protein domains at the N- terminal part of IL-33 make this cytokine as a nuclear factor to regulate the transcription of genes. The IL-33 suppresses transcription by binding to H2A-H2B dimers at the surface of the nucleosomes (Carriere *et al.*, 2007). Also, it was thought that it could function as an ‘alarmin’ released following cell necrosis to alerting the immune system about the stress or tissue damage (Moussion *et al.*, 2008). Moreover, it exerts its biological effects by binding to the ST2 receptor, a member of the interleukin-1 receptor family with two isoforms as follows: transmembrane (ST2L) and soluble (sST2), which interact with il-33 that is a single ligand for this receptor. Accordingly, ST2 is a part of the Toll-like receptor (TLR)/IL1R superfamily. IL-33 induces the production of cytokine by Th2, thus can elevate the pathogenesis of Th2-related diseases such as anaphylaxis, asthma, and atopic dermatitis (Ensaf *et al.*, 2014). IL- 33 can also induce inflammatory airway response by stimulating lung ILC2s (Cayrol, 2014). Depending on the disease and model, the effects of IL-33 can be pro-inflammatory or anti-inflammatory (Miller, 2011). We aimed to determine the role of IL-33 gene polymorphisms (rs1157505C/G and rs11792633C/T) with the risk of tuberculosis.

Materials and Methods

Study subjects

Study subjects' baseline characteristics are shown in table 1. Two IL-33 polymorphisms were successfully genotyped in 100 tuberculosis patients (mean age= 38.83±20.31 years) and 91 controls (mean age= 38.56±16.22 years). These samples were collected from Ali-ibn Abi Talib Hospital,

Zahedan, from 2012 to 2015. Detection of TB disease was performed by physical examination, medical history, test to diagnosis *M. tuberculosis* infection, and Chest X-ray The ethical committee at the Medical University of Zahedan approved this study.

Blood samples of cases and controls were collected in tubes containing EDTA. DNA was extracted from 500µl of blood samples using total DNA Extraction Kit tests 50 (Cat. No. EX6001.Sinaclon, Iran). The Tetra Arms technique, a fast and straightforward detection technique, was used to identify the genetic diversity of the gene under study.

PCR experiments were performed in a total volume of 25 µL. The PCR mixture consisted of 50 ng of genomic DNA, 10 pM of each primer, and premix master mix (including *Taq*, dNTP, PCR buffer, and Mg²⁺). The PCR reactions were subjected to initial heating at 95°C for 5 min, followed by 35 cycles at 95°C for the 30s, annealing temperature at 63°C (C/G SNP), 61°C(C/T) for 20s, extension at 72°C for 30s and final extension 72°C for 10 min. The PCR products were electrophoresed using a 2% agarose gel, stained with DNA green viewer, and visualized under ultraviolet illumination. The sequences and annealing temperature of primers are shown in Table 2. Statistical analysis was performed using SPSS statistical software ver. 20 and Epical version 3.2. The association between polymorphisms and susceptibility to tuberculosis was examined by estimating odds ratios (OR) and 95% confidence intervals (95% CI). Categorical data were analyzed by Pearson's χ^2 .

Results

No significant differences in allele frequencies of rs1157505C/G and rs11792633C/T were observed between cases and controls groups, with OR= 0.8, 95% CI= 0.468-1.3, $p= 0.42$, and OR= 1.06, 95% CI=0.69-1.64, $p=0.77$, respectively (Table 3). In this study, we investigated two SNPs in the IL-33 gene, (rs1157505C/G and rs11792633C/T) and discovered one genotype of rs11792633C/T SNP of IL-33 gene, CT genotype, which was significantly associated with tuberculosis at univariate analysis, with OR=1.94, 95% CI=1.025-3.66, p -value=0.042 (Table 4).

Table 1. The socio-demographic characteristics of the case and control groups.

Variables	Cases (N=100)	Controls (N=91)	P-value
Sex			
Females	54	55	0.37
Males	46	36	
Age	38.83±20.31	38.56±16.22	0.92
BMI	18.83±3.95	20.6±4.74	0.006
Educational level			
Illiteracy	70	18	<0.001
Primary	17	49	
School			
Guidance	8	19	
High school	2	2	
AD	3	2	
BA	-	-	
≥ MA	-	1	
Job-status			
Unemployed	7	5	0.13
Housekeeper	46	46	
Worker	12	2	
Farmer	2	0	
Rancher	1	1	
Self-employment	23	30	
Employee & Retired	3	4	
Student	6	3	

Table 2. The sequences and annealing temperatures of the primers.

SNPs	Primer name	Sequence (5'→3')	Product size (bp)	Annealing temp (°C)
IL-33rs1157505 C/G	Fi(C)	5'-GCCTCCAATTCCTGGGCTCAAGCAATCTTC-3'	139(C)	63
	Ri(G):	5'-CCAGCTGCTTGGGAAGCTGAGATCGC-3'	109(G)	
	Fo	5'-ATCTTGCTCTGTAGCGCAGGCTGGAGCA-3'	192	
	Ro	5'-TTCCAGCTGGCAACAAGCAAGACCC-3'		
IL-33rs11792633 C/T	Fi(C)	5'-CCCAGAGTCCACACTCAGTATTAGGCAGGC-3'	176(C)	61
	Ri(T)	5'-TAGTCAGCATCACATGGGAACGTGATCGA-3'	193(T)	
	Fo	5'-TGCTTGCTCTACTAGATGCTAGCCCCACA-3'	310	
	Ro	5'-GCATGAGTTTTGGTGGAAACATTCAAACCA-3'		

Table 3. Genotype and allele frequency of *IL-33* gene C/G and C/T polymorphisms in Tuberculosis patients and healthy controls.

Genotype	Cases (%) N= 100	Controls (%) N= 91	OR	CI (95%)	P-value
A) Genotypes and allele Frequency of IL33 C/G					
CC	76(76%)	73(80.23%)	-	-	Ref
CG	11(11%)	8(8.79%)	0.75	0.288-1.99	0.57
GG	13(13%)	10(10.98%)	0.8	0.331-1.94	0.62
CG+GG	24(24%)	18(19.78%)	1.28	0.64-2.55	0.48
C	163(81.5%)	154(84.61%)	1	-	Ref
G	37(18.5%)	28(15.39%)	0.8	0.468-1.37	0.42
B) Genotypes and allele Frequency of IL33 C/T					
CC	56(56%)	43(47.25%)	-	-	Ref
CT	27(27%)	38(41.75%)	1.94	1.025-3.66	0.042
TT	17(17%)	10(11%)	0.78	0.325-1.87	0.57
CT+TT	44(44%)	48(52.74%)	0.71	0.398-1.24	0.23
C	139(69.5%)	124(68.14%)	1	-	Ref
T	61(30.5%)	58(31.86%)	1.06	0.69-1.64	0.77

Moreover, the rs11792633C/T polymorphism has no significant association with tuberculosis in the same genotype after adjustment for binary logistic regression (OR=2.2, 95% CI= 0.875-5.47, $p=0.09$), which shown in table 4. As given in Table 5, the CC+CT genotype of IL-33 gene

significantly increased the risk of tuberculosis (OR= 0.48, 95% CI= 0.236-1.01, $p=0.054$). The analysis of gene-gene interaction between IL-33(rs1157505C/G and rs11792633C/T) did not find significant data (Table 5).

Table 4. Association of rs1157505C/G and rs11792633C/T variations and Tuberculosis after adjustment for binary logistic regression.

Genotype	Unadjusted			Adjusted*		
	OR	95%CI	P-value	OR	95%CI	P-value
IL33 C/G						
CC (ref)	1	-	-	1	-	-
CG	0.75	0.288-1.99	0.57	0.37	0.067-2.003	0.25
GG	0.8	0.331-1.94	0.62	0.91	0.224-3.65	0.88
CG+GG	1.28	0.64-2.55	0.48	2.379E9	-	0.99
IL33 C/T						
CC (ref)	-	-	-	-	-	-
CT	1.94	1.025-3.66	0.042	2.2	0.875-5.47	0.09
TT	0.78	0.325-1.87	0.57	0.4	0.1-1.49	0.17
CT+TT	0.71	0.398-1.24	0.23	0.84	-	1

Table 5. Association of different genotypic combinations of *IL-33* gene and Tuberculosis.

Genotype	Cases (%)	Controls (%)	OR	95%CI	P-value
CC/CC	44(44%)	34(37.36%)	-	-	Ref
CC/CT	19(19%)	30(32.96%)	0.48	0.236-1.01	0.054
CC/TT	13(13%)	9(9.89%)	1.11	0.427-2.91	0.82
GC/CC	3(3%)	4(4.39%)	0.58	0.121-2.76	0.49
GC/CT	6(6%)	3(3.29%)	1.54	0.36-6.63	0.55
GC/TT	2(2%)	1(1.09%)	1.54	0.134-17.76	0.72
GG/CC	10(10%)	5(5.49%)	1.54	0.483-4.94	0.46
GG/CT	1(1%)	5(5.49%)	0.155	0.017-1.38	0.09
GG/TT	2(2%)	0	-	-	0.9

Discussion

The present study analyzed for the first time for SNPs of *IL-33* encoding gene among Iranian tuberculosis patients. Clinical evidence showed that the host response to *M. tuberculosis* plays a vital role in the clinical signs of people exposed to this microorganism (Schluger and Rom 1998). Tissue damage, inflammation, and infection caused by an environmental agent triggers the release of *il-33* from the cells. This stimulation has been linked to the induction of *IL-33*. For example, a pulmonary challenge with LPS containing house dust mite can lead to an epithelial-dependent release of *IL-33* protein in the lungs of mice (Smith, 2010). *IL-33* plays an important role in the development and pathogenesis of tuberculosis by reducing macrophage-derived foam cell formation, support of Th2 cells, and endothelial cell function (Kurowska-Stolarska *et al.*, 2011). In the asthmatics and idiopathic pulmonary fibrosis patients (Tajima *et al.*, 2003), the concentration of sST2 is elevated, that is showing that various human diseases are associated with the *IL-33*-sST2 signaling pathway (Oshikawa *et al.*, 2001). The up-regulation of soluble receptor in pathological conditions provides indirect evidence for activation of pathway related to disease; however, the possible roles of *IL-33*

gene in tuberculosis remain poorly understood up to now. Many research projects have indicated that the variation in the *IL-33* gene is related to Allergic rhinitis (Sakashita *et al.*, 2008) and asthma (Gudbjartsson *et al.*, 2009) disorders. According to a study performed by Azazi *et al.*, high levels of *il-33* and its receptor sST2 in the serum of patients with bronchial asthma, indicate their importance as therapeutic targets in asthma (Azazi *et al.*, 2014). Moreover, the results of a study by Lee *et al.* have shown that there is an association among the elevated pleural *IL-33* levels and TB pleurisy (Lee *et al.*, 2013). *IFN- γ* and *TNF*, which locally increased in the pleural space of patients with the tuberculous pleural effusion, can induce *IL-33* expression. Findings of the study by Lee *et al.* indicated that *IL-33* is more relevant to the TPE pathophysiology than other types of pleural effusions (Lee *et al.*, 2013). It is still unclear how genetic diversity may influence the *IL-33* function to increase or decrease the risk of TB. However, the present study had several limitations. In the present study, which was a case-control, all the contributors were Iranian, and confounding variables including ethnicity were excluded. Indeed, the subjects should be from diverse racial and ethnic backgrounds that may influence the results of the study. So, no test was performed to determine the practical

consequences of genetic variability in the cases, such as the levels of IL-33 in tissue or plasma, and also the relation among polymorphisms in the *IL-33* gene and tuberculosis are conjectural. Finally, the results of the current study are based on a few numbers of cases and must be viewed as preliminary. Therefore, further studies on large-scale and diverse populations are needed to clarify the exact association among *IL-33* gene polymorphisms and the development of TB.

Conclusion

The study concluded that genetic variation of rs11792633C/T in the *IL-33* gene maybe have a central role in developing tuberculosis in Southeastern Iran and supplied more contribution toward new chance to consider prevention, pathogenesis, and treatment of tuberculosis.

Conflicts of interest

The authors declare that they have no conflict of interest.

References

- Azazi EA, Elshora AE, Tantawy EA, Elsayd MA. 2014. Serum levels of Interleukin-33 and its soluble receptor ST2 in asthmatic patients. *Egypt J Chest Dis Tuberc* 63(2): 279-284.
- Blok DC, Kager LM, Hoogendijk AJ, Lede IO, Rahman W, Afroz R, de Jong MD. 2015. Expression of inhibitory regulators of innate immunity in patients with active tuberculosis. *BMC Infect Dis* 15(1): 98. doi: 10.1186/s12879-015-0833-z.
- Carriere V, Roussel L, Ortega N, Lacorre DA, Americh L, Aguilar L, Girard JP. 2007. IL-33, the IL-1-like cytokine ligand for ST2 receptor, is a chromatin-associated nuclear factor in vivo. *Proc Natl Acad Sci* 104: 282-287.
- Cayrol C, Girard JP. 2014. IL-33: an alarmin cytokine with crucial roles in innate immunity, inflammation and allergy. *Curr Opin Immunol* 31: 31-37.
- Flynn JL, Chan J. 2001. Tuberculosis: latency and reactivation. *Infect Immun* 69: 4195-4201.
- Getahun H, Gunneberg C, Granich R, Nunn P. 2010. HIV infection-associated tuberculosis: the epidemiology and the response. *Clin Infect Dis* 50 Suppl 3: S201-207.
- Gudbjartsson DF, Bjornsdottir US, Halapi E, Helgadóttir A, Sulem P, Jonsdóttir GM, Williams C. 2009. Sequence variants affecting eosinophil numbers associate with asthma and myocardial infarction. *Nat Genet* 41: 342-347.
- Kurowska-Stolarska M, Hueber A, Stolarski B, McInnes IB. 2011. Interleukin-33: a novel mediator with a role in distinct disease pathologies. *J Intern Med* 269: 29-35
- Lee KS, Kim HR, Kwak S, Choi KH, Cho JH, Lee YJ, Park DS. 2013. Association between elevated pleural interleukin-33 levels and tuberculous pleurisy. *Ann Lab Med* 33: 45-51.
- Miller AM. 2011. Role of IL-33 in inflammation and disease. doi: 10.1186/1476-9255-8-22
- Moosazadeh M, Khanjani N, Bahrampour A, Nasehi M. 2014. Does tuberculosis have a seasonal pattern among migrant population entering Iran? *Int J Health Policy Manag* 2: 181-185.
- Moussion C, Ortega N, Girard JP. 2008. The IL-1-like cytokine IL-33 is constitutively expressed in the nucleus of endothelial cells and epithelial cells in vivo: a novel 'alarmin'? *PLoS One* 3: e3331.
- Oshikawa K, Kuroiwa K, Tago K, Iwahana H, Yanagisawa KEN, Ohno S, Sugiyama Y. 2001. Elevated soluble ST2 protein levels in sera of patients with asthma with an acute exacerbation. *Am J Respir Crit Care Med* 164: 277-281.
- Raja A. 2004. Immunology of tuberculosis. *Indian J Med Res* 120: 213-232.
- Sakashita M, Yoshimoto T, Hirota T, Harada M, Okubo K, Osawa Y, Tamari M. 2008. Association of serum interleukin-33 level and the interleukin-33 genetic variant with Japanese cedar pollinosis. *Clin Exp Allergy* 38: 1875-1881.
- Schluger NW, Rom WN. 1998. The host immune response to tuberculosis. *Am J Respir Crit Care Med*. 157(3): 679-691
- Schmitz J, Owyang A, Oldham E, Song Y, Murphy E, McClanahan TK, Gorman DM. 2005. IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines. *Immunity* 23: 479-490.

- Smith DE. 2010. IL-33: a tissue derived cytokine pathway involved in allergic inflammation and asthma. *Clin Exp Allergy* 40: 200-208.
- Tajima S, Oshikawa K, Tominaga SI, Sugiyama Y. 2003. The increase in serum soluble ST2 protein upon acute exacerbation of idiopathic pulmonary fibrosis. *Chest* 124: 1206-1214.
- Yagami A, Orihara K, Morita H, Futamura K, Hashimoto N, Matsumoto K, Matsuda A. 2010. IL-33 mediates inflammatory responses in human lung tissue cells. *J Immunol* 185: 5743-5750.
- Zumla A, George A, Sharma V, Herbert N. 2013. WHO's 2013 global report on tuberculosis: successes, threats, and opportunities. *Lancet* 382: 1765-1767.
- Zumla A, Rao M, Parida SK, Keshavjee S, Cassell G, Wallis R, Maeurer M. 2015. Inflammation and tuberculosis: host-directed therapies. *J Intern Med* 277: 373-387.

بررسی پلی مورفیسم‌های ژن IL-33 (rs11792633C/T و rs1157505C/G) و خطر توبرکلوزیس در جنوب شرق ایران

طیبه باران‌زهی^۱، شهربانو کریمی دوانی^۲، دُر محمد کردی تمندانی^۱

^۱گروه زیست‌شناسی، دانشگاه سیستان و بلوچستان، زاهدان، ایران

^۲مرکز جامع ژنتیک جنوب کشور، دانشگاه علوم پزشکی شیراز، شیراز، ایران

چکیده

سل یک بیماری عفونی مهیم است که به طور کلی ریه‌ها را تحت تأثیر قرار می‌دهد. در برخی موارد می‌تواند بر کبد و کلیه تأثیرگذار باشد. ژنتیک میزبان ممکن است، بر سل ناشی از باسیلوس مایکوباکتریوم توبرکلوزیس، تأثیر بگذارد. عمده فاکتورهای خطر این بیماری، ضعف سیستم ایمنی بدن به دلیل HIV / AIDS، دیابت، بیماری شدید کلیوی، برخی سرطانها، درمان سرطان و سوء تغذیه می‌باشند. اینترلوکین-۳۳ در فعال سازی mast cell ها، ائوزینوفیل ها، بازوفیل ها و سلول های کشنده طبیعی و بلوغ سلول های Th2 نقش دارد. محافظت در برابر سل به رشد TH1 و پاسخ های سلول T CD8(+) مرتبط است. IL-33، در حال حاضر با نمو هر دو نوع سلولهای ایمنی T CD8(+) و CD4(+) مرتبط است. هدف از این مطالعه بررسی، ارتباط بین عفونت مایکوباکتریوم توبرکلوزیس و پلی مورفیسم های ژن اینترلوکین-۳۳ (rs1157505C / G & rs11792633C / T) با سل، در نمونه های بیمار و سالم، در ناحیه با شیوع بالای این بیماری در ایران است. در این مطالعه ۱۰۰ بیمار مبتلا به بیماری سل و ۹۱ فرد سالم وارد مطالعه شدند. پلی مورفیسم ژن IL-33 توسط تکنیک T-ARMS-PCR ژنوتایپ شد. آنالیز ترکیب های دیپلو تیپ بین پلی مورفیسم های IL-33 نشان داد که، میزان ارتباط برای ژنوتیپهای CC / CT ترکیب شده بیشتر بود. هر چند، از لحاظ آماری تفاوتها معنی دار نبود. ژنوتیپ CT مربوط به پلی مورفیسم C / T ژن IL-33 خطر ابتلا به سل را افزایش داد. ژنوتیپهای ترکیبی CG + GG مربوط به پلی مورفیسم C / G ژن IL-33 همچنین خطر ابتلا به سل را افزایش داد، اما این اختلاف از نظر آماری معنی دار نبود. در نتیجه، پلی مورفیسم ژن IL-33 ممکن است به عنوان عامل مهمی در ایجاد سل در ایران در نظر گرفته شود.

واژگان کلیدی: باسیلوس، IL-33، rs1157505 C/G، rs11792633 C/T؛ توبرکلوزیس

Please cite this paper as: Baranzehi T, Karimi-Davani S, Kordi-Tamandani DM. 2020. Analysis of IL-33 gene polymorphisms (rs1157505C/G and rs11792633C/T) and the risk of tuberculosis in Southeastern Iran. *J Genet Resour* 6(1): 54-59. doi: 10.22080/jgr.2020.17777.1166.