



Research Paper

## GENUS OF COMMUTING CONJUGACY CLASS GRAPH OF CERTAIN FINITE GROUPS

PARTHAJIT BHOWAL AND RAJAT KANTI NATH\*

ABSTRACT. For a non-abelian group  $G$ , its commuting conjugacy class graph  $CCC(G)$  is a simple undirected graph whose vertex set is the set of conjugacy classes of the non-central elements of  $G$  and two distinct vertices  $x^G$  and  $y^G$  are adjacent if there exists some elements  $x' \in x^G$  and  $y' \in y^G$  such that  $x'y' = y'x'$ . In this paper we compute the genus of  $CCC(G)$  for six well-known classes of non-abelian two-generated groups (viz.  $D_{2n}, SD_{8n}, Q_{4m}, V_{8n}, U_{(n,m)}$  and  $G(p, m, n)$ ) and determine whether  $CCC(G)$  for these groups are planar, toroidal, double-toroidal or triple-toroidal.

### 1. INTRODUCTION

The commuting conjugacy class graph of a non-abelian group  $G$  is a simple undirected graph, denoted by  $CCC(G)$ , whose vertex set is the set of conjugacy classes of the non-central elements of  $G$  and two distinct vertices  $x^G$  and  $y^G$  are adjacent if there exist some elements

DOI: 10.29252/as.2021.2444

MSC(2010): Primary: 20D99, 05C25, 05C10, 05E15.

Keywords: Commuting conjugacy class graph, Genus, Finite group.

Received: 30 January 2021, Accepted: 24 October 2021.

\*Corresponding author

$x' \in x^G$  and  $y' \in y^G$  such that  $x'y' = y'x'$ . This graph extends the notion of commuting graph of a finite group introduced by Brauer and Fowler [5], in 1955. Commuting graphs of finite algebraic structures, its extensions, generalizations and their complements remain active topic of research over the years. In 2009, Herzog, Longobardi and Maj [8] initiated the study of commuting conjugacy class graph of groups. In 2016, finite groups having triangle-free commuting conjugacy class graph were characterized by Mohammadian et al. [9]. Ashrafi and Salahshour have also considered commuting conjugacy class graph of finite groups in their recent work [10], where they obtain structures of  $CCC(G)$  for the following groups:

$$\begin{aligned}
 D_{2n} &= \langle \alpha, \beta : \alpha^n = \beta^2 = 1, \beta\alpha\beta = \alpha^{-1} \rangle \text{ for } n \geq 3, \\
 SD_{8n} &= \langle \alpha, \beta : \alpha^{4n} = \beta^2 = 1, \beta\alpha\beta = \alpha^{2n-1} \rangle \text{ for } n \geq 2, \\
 Q_{4m} &= \langle \alpha, \beta : \alpha^{2m} = 1, \alpha^m = \beta^2, \beta^{-1}\alpha\beta = \alpha^{-1} \rangle \text{ for } m \geq 2, \\
 V_{8n} &= \langle \alpha, \beta : \alpha^{2n} = \beta^4 = 1, \beta\alpha = \alpha^{-1}\beta^{-1}, \beta^{-1}\alpha = \alpha^{-1}\beta \rangle \text{ for } n \geq 2, \\
 U_{(n,m)} &= \langle \alpha, \beta : \alpha^{2n} = \beta^m = 1, \alpha^{-1}\beta\alpha = \beta^{-1} \rangle \text{ for } m \geq 2 \text{ and } n \geq 2 \text{ and} \\
 G(p, m, n) &= \langle \alpha, \beta : \alpha^{p^m} = \beta^{p^n} = [\alpha, \beta]^p = 1, [\alpha, [\alpha, \beta]] = [\beta, [\alpha, \beta]] = 1 \rangle,
 \end{aligned}$$

where  $p$  is any prime,  $m \geq 1$  and  $n \geq 1$ .

Continuing the works of Ashrafi and Salahshour [10], in [2, 3] Bhowal and Nath have obtained various spectra and energies of commuting conjugacy class graphs of finite groups. In this paper we compute genus of commuting conjugacy class graph of the above mentioned groups and determine whether  $CCC(G)$  for those groups are planar, toroidal, double-toroidal or triple-toroidal. The genus  $\gamma(\mathcal{G})$  of a graph  $\mathcal{G}$  is the smallest integer  $k \geq 0$  such that  $\mathcal{G}$  can be embedded on the surface obtained by attaching  $k$  handles to a sphere. A graph  $\mathcal{G}$  is called planar, toroidal, double-toroidal or triple-toroidal if  $\mathcal{G}$  has genus 0, 1, 2 or 3 respectively. Results on genus of commuting graphs of finite groups, including its various extensions, can be found in [1, 4, 6, 7]. However, genus of commuting conjugacy class graph of finite groups are not yet studied.

## 2. GENUS OF $CCC(G)$

Let  $K_n$  be the complete graph on  $n$  vertices and  $mK_n$  the disjoint union of  $m$  copies of  $K_n$ . It is well known that  $\gamma(K_n) = 0$  if  $n = 1, 2$ . If  $n \geq 3$  then, by [12, Theorem 6-38], we have

$$(1) \quad \gamma(K_n) = \left\lceil \frac{(n-3)(n-4)}{12} \right\rceil,$$

where  $\lceil x \rceil$  denotes the smallest integer greater than or equal to  $x$  for any real number  $x$ . By [11, Corollary 2], we also have the following lemma.

**Lemma 2.1.** *If  $\mathcal{G} = m_1K_{n_1} \sqcup m_2K_{n_2}$  then  $\gamma(\mathcal{G}) = m_1\gamma(K_{n_1}) + m_2\gamma(K_{n_2})$ .*

Now we compute genus of commuting conjugacy class graph of the groups  $D_{2n}, SD_{8n}, Q_{4m}, V_{8n}, U_{(n,m)}$  and  $G(p, m, n)$  one by one and check their planarity, toroidality etc.

**Theorem 2.2.** *Let  $G = D_{2n}$ . Then*

- (1)  $CCC(G)$  is planar if and only if  $3 \leq n \leq 10$ .
- (2)  $CCC(G)$  is toroidal if and only if  $11 \leq n \leq 16$ .
- (3)  $CCC(G)$  is double-toroidal if and only if  $n = 17, 18$ .
- (4)  $CCC(G)$  is triple-toroidal if and only if  $n = 19, 20$ .
- (5)  $\gamma(CCC(G)) = \begin{cases} \left\lceil \frac{(n-7)(n-9)}{48} \right\rceil, & \text{if } n \text{ is odd and } n \geq 21 \\ \left\lceil \frac{(n-8)(n-10)}{48} \right\rceil, & \text{if } n \text{ is even and } n \geq 22. \end{cases}$

*Proof.* Consider the following cases.

**Case 1.**  $n$  is odd.

By [10, Proposition 2.1] we have  $CCC(G) = K_1 \sqcup K_{\frac{n-1}{2}}$ . Therefore, for  $n = 3$  and  $5$ , it follows that  $CCC(G) = 2K_1, K_1 \sqcup K_2$  respectively; and hence  $CCC(G)$  is planar. If  $n \geq 7$  then, by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = \gamma(K_{\frac{n-1}{2}}) = \left\lceil \frac{(n-7)(n-9)}{48} \right\rceil.$$

Clearly  $\gamma(CCC(G)) = 0$  if and only if  $n = 7$  or  $9$ . Also,  $\gamma(CCC(G)) = 1$  if  $n = 11, 13$  or  $15$ ;  $\gamma(CCC(G)) = 2$  if  $n = 17$ ;  $\gamma(CCC(G)) = 3$  if  $n = 19$ . For  $n \geq 21$  we have

$$\frac{(n-7)(n-9)}{48} \geq \frac{7}{2} = 3.5,$$

and so

$$\gamma(CCC(G)) = \left\lceil \frac{(n-7)(n-9)}{48} \right\rceil \geq 4.$$

Thus,  $CCC(G)$  is planar if and only if  $n = 3, 5, 7, 9$ ; toroidal if and only if  $n = 11, 13, 15$ ; double-toroidal if and only if  $n = 17$  and triple-toroidal if and only if  $n = 19$ .

**Case 2.**  $n$  is even.

By [10, Proposition 2.1] we have

$$CCC(G) = \begin{cases} 2K_1 \sqcup K_{\frac{n}{2}-1}, & \text{if } n \text{ and } \frac{n}{2} \text{ are even} \\ K_2 \sqcup K_{\frac{n}{2}-1}, & \text{if } n \text{ is even and } \frac{n}{2} \text{ is odd.} \end{cases}$$

Therefore, for  $n = 4$  and  $6$ , it follows that  $CCC(G) = 3K_1, 2K_2$  respectively; and hence  $CCC(G)$  is planar. If  $n \geq 8$  then, by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = \gamma(K_{\frac{n}{2}-1}) = \left\lceil \frac{(n-8)(n-10)}{48} \right\rceil.$$

Clearly  $\gamma(CCC(G)) = 0$  if and only if  $n = 8$  or  $10$ . Also,  $\gamma(CCC(G)) = 1$  if  $n = 12, 14$  or  $16$ ;  $\gamma(CCC(G)) = 2$  if  $n = 18$ ;  $\gamma(CCC(G)) = 3$  if  $n = 20$ . For  $n \geq 22$  we have

$$\frac{(n-8)(n-10)}{48} \geq \frac{7}{2} = 3.5,$$

and so

$$\gamma(CCC(G)) = \left\lceil \frac{(n-8)(n-10)}{48} \right\rceil \geq 4.$$

Thus,  $CCC(G)$  is planar if and only if  $n = 4, 6, 8, 10$ ; toroidal if and only if  $n = 12, 14, 16$ ; double-toroidal if and only if  $n = 18$  and triple-toroidal if and only if  $n = 20$ . Hence the result follows.  $\square$

**Theorem 2.3.** *Let  $G = SD_{8n}$ . Then*

- (1)  $CCC(G)$  is planar if and only if  $n = 2$  or  $3$ .
- (2)  $CCC(G)$  is toroidal if and only if  $n = 4$ .
- (3)  $CCC(G)$  is double-toroidal if and only if  $n = 5$ .
- (4)  $CCC(G)$  is not triple-toroidal.
- (5)  $\gamma(CCC(G)) = \begin{cases} \left\lceil \frac{(n-3)(2n-5)}{6} \right\rceil, & \text{if } n \text{ is odd and } n \geq 7 \\ \left\lceil \frac{(n-2)(2n-5)}{6} \right\rceil, & \text{if } n \text{ is even and } n \geq 6. \end{cases}$

*Proof.* Consider the following cases.

**Case 1.**  $n$  is odd.

By [10, Proposition 2.1] we have  $CCC(G) = K_4 \sqcup K_{2n-2}$ . For  $n \geq 3$ , by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = \gamma(K_4) + \gamma(K_{2n-2}) = \left\lceil \frac{(n-3)(2n-5)}{6} \right\rceil.$$

Clearly  $\gamma(CCC(G)) = 0$  if  $n = 3$ ;  $\gamma(CCC(G)) = 2$  if  $n = 5$ . For  $n \geq 7$  we have

$$\frac{(n-3)(2n-5)}{6} \geq 6,$$

and so

$$\gamma(CCC(G)) = \left\lceil \frac{(n-3)(2n-5)}{6} \right\rceil \geq 6.$$

Thus  $CCC(G)$  is planar if and only if  $n = 3$ ; double-toroidal if and only if  $n = 5$ .

**Case 2.**  $n$  is even.

By [10, Proposition 2.1] we have  $CCC(G) = 2K_1 \sqcup K_{2n-1}$ . For  $n \geq 2$ , by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = \gamma(K_{2n-1}) = \left\lceil \frac{(n-2)(2n-5)}{6} \right\rceil.$$

Clearly  $\gamma(CCC(G)) = 0$  if  $n = 2$ ;  $\gamma(CCC(G)) = 1$  if  $n = 4$ . For  $n \geq 6$  we have

$$\frac{(n-2)(2n-5)}{6} \geq \frac{14}{3},$$

and so

$$\gamma(CCC(G)) = \left\lceil \frac{(n-2)(2n-5)}{6} \right\rceil \geq 5.$$

Thus  $CCC(G)$  is planar if and only if  $n = 2$ ; toroidal if and only if  $n = 4$ . Hence the result follows.  $\square$

**Theorem 2.4.** *Let  $G = Q_{4m}$ . Then*

- (1)  $CCC(G)$  is planar if and only if  $m = 2, 3, 4$  or  $5$ .
- (2)  $CCC(G)$  is toroidal if and only if  $m = 6, 7$  or  $8$ .
- (3)  $CCC(G)$  is double-toroidal if and only if  $m = 9$ .
- (4)  $CCC(G)$  is triple-toroidal if and only if  $m = 10$ .
- (5)  $\gamma(CCC(G)) = \left\lceil \frac{(m-4)(m-5)}{12} \right\rceil$  for  $m \geq 11$ .

*Proof.* By [10, Proposition 2.1] we have

$$CCC(G) = \begin{cases} K_2 \sqcup K_{m-1}, & \text{if } m \text{ is odd} \\ 2K_1 \sqcup K_{m-1}, & \text{if } m \text{ is even.} \end{cases}$$

Therefore, for  $m = 2, 3$ , it follows that  $CCC(G) = 3K_1, 2K_2$  respectively; and hence  $CCC(G)$  is planar. If  $m \geq 4$  then, by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = \gamma(K_{m-1}) = \left\lceil \frac{(m-4)(m-5)}{12} \right\rceil.$$

Clearly  $\gamma(CCC(G)) = 0$  if and only if  $m = 4$  or  $5$ . Also,  $\gamma(CCC(G)) = 1$  if  $m = 6, 7$  or  $8$ ;  $\gamma(CCC(G)) = 2$  if  $m = 9$ ;  $\gamma(CCC(G)) = 3$  if  $m = 10$ . For  $m \geq 11$  we have

$$\frac{(m-4)(m-5)}{12} \geq \frac{7}{2} = 3.5,$$

and so

$$\gamma(CCC(G)) = \left\lceil \frac{(m-4)(m-5)}{12} \right\rceil \geq 4.$$

Thus,  $CCC(G)$  is planar if and only if  $m = 2, 3, 4, 5$ ; toroidal if and only if  $m = 6, 7, 8$ ; double-toroidal if and only if  $m = 9$  and triple-toroidal if and only if  $m = 10$ . Hence the result follows.

$\square$

**Theorem 2.5.** Let  $G = V_{8n}$ . Then

- (1)  $CCC(G)$  is planar if and only if  $n = 2$ .
- (2)  $CCC(G)$  is toroidal if and only if  $n = 3$  or  $4$ .
- (3)  $CCC(G)$  is not double-toroidal.
- (4)  $CCC(G)$  is triple-toroidal if and only if  $n = 5$ .
- (5)  $\gamma(CCC(G)) = \begin{cases} \left\lceil \frac{(n-2)(2n-5)}{6} \right\rceil, & \text{if } n \text{ is odd and } n \geq 7 \\ \left\lceil \frac{(n-3)(2n-5)}{6} \right\rceil, & \text{if } n \text{ is even and } n \geq 6. \end{cases}$

*Proof.* Consider the following cases.

**Case 1.**  $n$  is odd.

By [10, Proposition 2.1] we have  $CCC(G) = 2K_1 \sqcup K_{2n-1}$ . For  $n \geq 3$ , by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = \gamma(K_{2n-1}) = \left\lceil \frac{(n-2)(2n-5)}{6} \right\rceil.$$

Clearly  $\gamma(CCC(G)) = 1$  if  $n = 3$ ;  $\gamma(CCC(G)) = 3$  if  $n = 5$ . For  $n \geq 7$  we have

$$\frac{(n-2)(2n-5)}{6} \geq \frac{15}{2} = 7.5,$$

and so

$$\gamma(CCC(G)) = \left\lceil \frac{(n-2)(2n-5)}{6} \right\rceil \geq 8.$$

**Case 2.**  $n$  is even.

By [10, Proposition 2.1] we have  $CCC(G) = 2K_2 \sqcup K_{2n-2}$ . Therefore, for  $n = 2$  it follows that  $CCC(G) = 3K_2$ ; and hence  $CCC(G)$  is planar. If  $n \geq 4$  then, by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = \gamma(K_{2n-2}) = \left\lceil \frac{(n-3)(2n-5)}{6} \right\rceil.$$

Clearly  $\gamma(CCC(G)) = 1$  if  $n = 4$ . For  $n \geq 6$  we have

$$\frac{(n-3)(2n-5)}{6} \geq \frac{7}{2} = 3.5,$$

and so

$$\gamma(CCC(G)) = \left\lceil \frac{(n-3)(2n-5)}{6} \right\rceil \geq 4.$$

Thus  $CCC(G)$  is planar if and only if  $n = 2$ ; toroidal if and only if  $n = 4$ . Hence the result follows.  $\square$

**Theorem 2.6.** Let  $G = U_{(n,m)}$ . Then

- (1)  $CCC(G)$  is planar if and only if  $n = 2$  and  $m = 2, 3, 4, 5, 6$ ;  $n = 3$  and  $m = 2, 3, 4$ ; or  $n = 4$  and  $m = 2, 3, 4$ .
- (2)  $CCC(G)$  is toroidal if and only if  $n = 2$  and  $m = 7, 8$ ; or  $n = 3$  and  $m = 5, 6$ .

- (3)  $CCC(G)$  is double-toroidal if and only if  $n = 2$  and  $m = 9, 10$ ;  $n = 4$  and  $m = 5, 6$ ;  $n = 5$  and  $m = 2, 3$ ;  $n = 6$  and  $m = 2, 3$ ; or  $n = 7$  and  $m = 2, 3$ .
- (4)  $CCC(G)$  is triple-toroidal if and only if  $n = 3$  and  $m = 7, 8$ ;  $n = 5$  and  $m = 4$ ;  $n = 6$  and  $m = 4$ ; or  $n = 7$  and  $m = 4$ .

$$(5) \gamma(CCC(G)) = \begin{cases} \left\lceil \frac{(mn-n-6)(mn-n-8)}{48} \right\rceil, & \text{if } n = 2, m \text{ is odd and } m \geq 11 \\ \left\lceil \frac{(mn-2n-6)(mn-2n-8)}{48} \right\rceil, & \text{if } n = 2, m \text{ is even and } m \geq 12 \\ \left\lceil \frac{(mn-n-6)(mn-n-8)}{48} \right\rceil + \left\lceil \frac{(n-3)(n-4)}{12} \right\rceil, & \text{if } n = 3, m \text{ is odd and } m \geq 9; \\ & n = 4, m \geq 7; n = 5, m \geq 5; \\ & n = 6, m \geq 5; n = 7, m \geq 5; \\ & \text{or } n \geq 8, m \geq 3 \\ \left\lceil \frac{(mn-2n-6)(mn-2n-8)}{48} \right\rceil + 2 \left\lceil \frac{(n-3)(n-4)}{12} \right\rceil, & \text{if } n = 3, m \text{ is even and } m \geq 10; \\ & n = 4, m \geq 8; n = 5, m \geq 6; \\ & n = 6, m \geq 6; n = 7, m \geq 6; \\ & \text{or } n \geq 8, m \geq 2 \end{cases}$$

*Proof.* Consider the following cases.

**Case 1.**  $m$  is odd.

By [10, Proposition 2.3] we have  $CCC(G) = K_{\frac{n(m-1)}{2}} \sqcup K_n$ .

**Sub case 1.1**  $n = 2$ .

If  $n = 2$  then we have  $CCC(G) = K_{m-1} \sqcup K_2$ . Therefore, for  $m = 3$  it follows that  $CCC(G) = 2K_2$ ; and hence  $CCC(G)$  is planar. For  $m \geq 5$ , by Lemma 2.1, we have

$$\gamma(CCC(G)) = \gamma(K_{m-1}) = \left\lceil \frac{(m-4)(m-5)}{12} \right\rceil.$$

Clearly  $\gamma(CCC(G)) = 0$  if  $m = 5$ ;  $\gamma(CCC(G)) = 1$  if  $m = 7$ ;  $\gamma(CCC(G)) = 2$  if  $m = 9$ . For  $m \geq 11$  we have

$$\frac{(m-4)(m-5)}{12} \geq \frac{7}{2} = 3.5,$$

and so

$$\gamma(CCC(G)) = \left\lceil \frac{(m-4)(m-5)}{12} \right\rceil \geq 4.$$

Thus  $CCC(G)$  is planar if and only if  $m = 3, 5$ ; toroidal if and only if  $m = 7$ ; double-toroidal if and only if  $m = 9$ .

**Sub case 1.2**  $n \geq 3$ .

If  $n \geq 3$  then we have  $CCC(G) = K_{\frac{n(m-1)}{2}} \sqcup K_n$ . By Lemma 2.1, we have

$$\gamma(CCC(G)) = \gamma(K_{\frac{n(m-1)}{2}}) + \gamma(K_n) = \left\lceil \frac{(mn - n - 6)(mn - n - 8)}{48} \right\rceil + \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil.$$

Clearly  $\gamma(CCC(G)) = 0$  if  $n = 3, m = 3$  or  $n = 4, m = 3$ .  $\gamma(CCC(G)) = 1$  if  $n = 3, m = 5$ ;  $\gamma(CCC(G)) = 2$  if  $n = 4, m = 5$  or  $n = 5, m = 3$  or  $n = 6, m = 3$  or  $n = 7, m = 3$ ;  $\gamma(CCC(G)) = 3$  if  $n = 3, m = 7$ . If  $n = 3$  and  $m \geq 9$  then

$$\frac{(mn - n - 6)(mn - n - 8)}{48} = \frac{(m - 3)(3m - 11)}{16} \geq 6.$$

Therefore

$$\gamma(CCC(G)) = \left\lceil \frac{(mn - n - 6)(mn - n - 8)}{48} \right\rceil + \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 6.$$

If  $n = 4$  and  $m \geq 7$  then

$$\frac{(mn - n - 6)(mn - n - 8)}{48} = \frac{(2m - 5)(m - 3)}{6} \geq 6.$$

Therefore

$$\gamma(CCC(G)) = \left\lceil \frac{(mn - n - 6)(mn - n - 8)}{48} \right\rceil + \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 6.$$

If  $n = 5$  and  $m \geq 5$  then

$$\frac{(mn - n - 6)(mn - n - 8)}{48} = \frac{(5m - 11)(5m - 13)}{48} \geq \frac{7}{2} = 3.5 \quad \text{and} \quad \frac{(n - 3)(n - 4)}{12} = \frac{1}{6}.$$

Therefore

$$\gamma(CCC(G)) = \left\lceil \frac{(mn - n - 6)(mn - n - 8)}{48} \right\rceil + \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 5.$$

If  $n = 6$  and  $m \geq 5$  then

$$\frac{(mn - n - 6)(mn - n - 8)}{48} = \frac{(m - 2)(3m - 7)}{4} \geq 6 \quad \text{and} \quad \frac{(n - 3)(n - 4)}{12} = \frac{1}{2}.$$

Therefore

$$\gamma(CCC(G)) = \left\lceil \frac{(mn - n - 6)(mn - n - 8)}{48} \right\rceil + \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 7.$$

If  $n = 7$  and  $m \geq 5$  then

$$\frac{(mn - n - 6)(mn - n - 8)}{48} = \frac{(7m - 13)(7m - 15)}{48} \geq \frac{55}{6} \quad \text{and} \quad \frac{(n - 3)(n - 4)}{12} = 1.$$

Therefore

$$\gamma(CCC(G)) = \left\lceil \frac{(mn - n - 6)(mn - n - 8)}{48} \right\rceil + \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 11.$$

If  $n \geq 8$  and  $m \geq 3$  then

$$\frac{(mn - n - 6)(mn - n - 8)}{48} \geq \frac{(8(m - 1) - 6)(8(m - 1) - 7)}{48} \geq \frac{15}{8} \quad \text{and} \quad \frac{(n - 3)(n - 4)}{12} = \frac{5}{3}.$$



Therefore

$$\gamma(\mathcal{CCC}(G)) = \left\lceil \frac{(mn - n - 6)(mn - n - 8)}{48} \right\rceil + \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 4.$$

Thus  $\mathcal{CCC}(G)$  is planar if and only if  $n = 3, m = 3$  or  $n = 4, m = 3$ ; toroidal if and only if  $n = 3, m = 5$ ; double-toroidal if and only if  $n = 4, m = 5$  or  $n = 5, m = 3$  or  $n = 6, m = 3$  or  $n = 7, m = 3$ ; triple-toroidal if and only if  $n = 3, m = 7$ .

**Case 2.**  $m$  is even.

By [10, Proposition 2.3] we have  $\mathcal{CCC}(G) = K_{\frac{n(m-2)}{2}} \sqcup 2K_n$ .

**Sub case 2.1**  $n = 2$ .

If  $n = 2$  then we have  $\mathcal{CCC}(G) = K_{m-2} \sqcup 2K_2$ . Therefore, for  $m = 2, 4$  it follows that  $\mathcal{CCC}(G) = 2K_2$  and  $3K_2$ ; and hence  $\mathcal{CCC}(G)$  is planar. For  $m \geq 6$ , by Lemma 2.1, we have

$$\gamma(\mathcal{CCC}(G)) = \gamma(K_{m-2}) = \left\lceil \frac{(m - 5)(m - 6)}{12} \right\rceil.$$

Clearly  $\gamma(\mathcal{CCC}(G)) = 0$  if  $m = 6$ ;  $\gamma(\mathcal{CCC}(G)) = 1$  if  $m = 8$ ;  $\gamma(\mathcal{CCC}(G)) = 2$  if  $m = 10$ . For  $m \geq 12$  we have

$$\frac{(m - 5)(m - 6)}{12} \geq \frac{7}{2} = 3.5$$

and so

$$\gamma(\mathcal{CCC}(G)) = \left\lceil \frac{(m - 4)(m - 5)}{12} \right\rceil \geq 4.$$

Thus  $\mathcal{CCC}(G)$  is planar if and only if  $m = 2, 4, 6$ ; toroidal if and only if  $m = 8$ ; double-toroidal if and only if  $m = 10$ .

**Sub case 2.2**  $n \geq 3$ .

If  $n \geq 3$  then we have  $\mathcal{CCC}(G) = K_{\frac{n(m-2)}{2}} \sqcup 2K_n$ . By Lemma 2.1, we have

$$\gamma(\mathcal{CCC}(G)) = \gamma(K_{\frac{n(m-2)}{2}}) + \gamma(2K_n) = \left\lceil \frac{(mn - 2n - 6)(mn - 2n - 8)}{48} \right\rceil + 2 \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil.$$

Clearly  $\gamma(\mathcal{CCC}(G)) = 0$  if  $n = 3, m = 2, 4$  or  $n = 4, m = 2, 4$ .  $\gamma(\mathcal{CCC}(G)) = 1$  if  $n = 3, m = 6$ ;  $\gamma(\mathcal{CCC}(G)) = 2$  if  $n = 4, m = 6$  or  $n = 5, m = 2$  or  $n = 6, m = 2$  or  $n = 7, m = 2$ ;  $\gamma(\mathcal{CCC}(G)) = 3$  if  $n = 3, m = 8$  or  $n = 5, m = 4$  or  $n = 6, m = 4$  or  $n = 7, m = 4$ . If  $n = 3$  and  $m \geq 10$  then

$$\frac{(mn - 2n - 6)(mn - 2n - 8)}{48} = \frac{(m - 4)(3m - 14)}{16} \geq 6.$$

Therefore

$$\gamma(\mathcal{CCC}(G)) = \left\lceil \frac{(mn - 2n - 6)(mn - 2n - 8)}{48} \right\rceil + 2 \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 6.$$

If  $n = 4$  and  $m \geq 8$  then

$$\frac{(mn - 2n - 6)(mn - 2n - 8)}{48} = \frac{(m - 4)(2m - 7)}{6} \geq 6.$$

Therefore

$$\gamma(\mathcal{CCC}(G)) = \left\lceil \frac{(mn - 2n - 6)(mn - 2n - 8)}{48} \right\rceil + 2 \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 6.$$

If  $n = 5$  and  $m \geq 6$  then

$$\frac{(mn - 2n - 6)(mn - 2n - 8)}{48} = \frac{(5m - 16)(5m - 18)}{48} \geq \frac{7}{2} = 3.5 \quad \text{and} \quad \frac{(n - 3)(n - 4)}{12} = \frac{1}{6}.$$

Therefore

$$\gamma(\mathcal{CCC}(G)) = \left\lceil \frac{(mn - 2n - 6)(mn - 2n - 8)}{48} \right\rceil + 2 \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 6.$$

If  $n = 6$  and  $m \geq 6$  then

$$\frac{(mn - 2n - 6)(mn - 2n - 8)}{48} = \frac{(m - 3)(3m - 10)}{4} \geq 6 \quad \text{and} \quad \frac{(n - 3)(n - 4)}{12} = \frac{1}{6}.$$

Therefore

$$\gamma(\mathcal{CCC}(G)) = \left\lceil \frac{(mn - 2n - 6)(mn - 2n - 8)}{48} \right\rceil + 2 \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 8.$$

If  $n = 7$  and  $m \geq 6$  then

$$\frac{(mn - 2n - 6)(mn - 2n - 8)}{48} = \frac{(7m - 20)(7m - 22)}{48} \geq \frac{55}{6} \quad \text{and} \quad \frac{(n - 3)(n - 4)}{12} = 1.$$

Therefore

$$\gamma(\mathcal{CCC}(G)) = \left\lceil \frac{(mn - 2n - 6)(mn - 2n - 8)}{48} \right\rceil + 2 \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 12.$$

If  $n \geq 8$  and  $m \geq 2$  then

$$\frac{(n - 3)(n - 4)}{12} \geq \frac{5}{3} \quad \text{and} \quad \left\lceil \frac{(mn - 2n - 6)(mn - 2n - 8)}{48} \right\rceil \geq 0.$$

Therefore

$$\gamma(\mathcal{CCC}(G)) = \left\lceil \frac{(mn - 2n - 6)(mn - 2n - 8)}{48} \right\rceil + 2 \left\lceil \frac{(n - 3)(n - 4)}{12} \right\rceil \geq 4.$$

Thus  $\mathcal{CCC}(G)$  is planar if and only if  $n = 3, m = 2, 4$  or  $n = 4, m = 2, 4$ ; toroidal if and only if  $n = 3, m = 6$ ; double-toroidal if and only if  $n = 4, m = 6$  or  $n = 5, m = 2$  or  $n = 6, m = 2$  or  $n = 7, m = 2$ ; triple-toroidal if and only if  $n = 3, m = 8$  or  $n = 5, m = 4$  or  $n = 6, m = 4$  or  $n = 7, m = 4$ . Hence the result follows.  $\square$

**Theorem 2.7.** *Let  $G = G(p, m, n)$ . Then*

- (1)  $\mathcal{CCC}(G)$  is planar if and only if  $n = 1, m = 1, p = 2, 3, 5$ ;  $n = 1, m = 2, p = 2$ ;  $n = 1, m = 3, p = 2$ ;  $n = 2, m = 1, p = 2$ ;  $n = 2, m = 2, p = 2$ ; or  $n = 3, m = 1, p = 2$ .
- (2)  $\mathcal{CCC}(G)$  is not toroidal.
- (3)  $\mathcal{CCC}(G)$  is double-toroidal if and only if  $n = 2, m = 1, p = 3$ .
- (4)  $\mathcal{CCC}(G)$  is not triple-toroidal.

$$(5) \gamma(CCC(G)) = \left\{ \begin{array}{ll} (p+1) \left\lceil \frac{(p-4)(p-5)}{12} \right\rceil, & \text{if } n=1, m=1, p \geq 7 \\ (p+1) \left\lceil \frac{(p^2-p-3)(p^2-p-4)}{12} \right\rceil, & \text{if } n=1, m=2, p \geq 3 \\ (p+1) \left\lceil \frac{(p^3-p^2-3)(p^3-p^2-4)}{12} \right\rceil, & \text{if } n=1, m=3, p \geq 3 \\ (p+1) \left\lceil \frac{(p^m-p^{m-1}-3)(p^m-p^{m-1}-4)}{12} \right\rceil, & \text{if } n=1, m \geq 3, p \geq 2 \\ (p^2-p) \left\lceil \frac{(p-4)(p-5)}{12} \right\rceil + 2 \left\lceil \frac{(p^2-p-3)(p^2-p-4)}{12} \right\rceil, & \text{if } n=2, m=1, p \geq 5 \\ (p^2-p) \left\lceil \frac{(p^2-p-3)(p^2-p-4)}{12} \right\rceil \\ + 2 \left\lceil \frac{(p^3-p^2-3)(p^3-p^2-4)}{12} \right\rceil, & \text{if } n=2, m=2, p \geq 3 \\ (p^2-p) \left\lceil \frac{(p^{m-1}(p-1)-3)(p^{m-1}(p-1)-4)}{12} \right\rceil \\ + 2 \left\lceil \frac{(p^m(p-1)-3)(p^m(p-1)-4)}{12} \right\rceil, & \text{if } n=2, m \geq 3, p \geq 2 \\ 36, & \text{if } n=3, m=1, p=3 \\ p^2(p-1) \left\lceil \frac{(p-4)(p-5)}{12} \right\rceil + 2 \left\lceil \frac{(p^3-p^2-3)(p^3-p^2-4)}{12} \right\rceil, & \text{if } n=3, m=1, p \geq 5 \\ 4, & \text{if } n=3, m=2, p=2 \\ p^2(p-1) \left\lceil \frac{(p^m-p^{m-1}-3)(p^m-p^{m-1}-4)}{12} \right\rceil \\ + 2 \left\lceil \frac{(p^{m+2}-p^{m+1}-3)(p^{m+2}-p^{m+1}-4)}{12} \right\rceil, & \text{if } n=3, m=2, p \geq 3 \\ & \text{or } n=3, m \geq 3, p \geq 2 \\ 2 \left\lceil \frac{(p^{n-1}(p^m-p^{m-1})-3)(p^{n-1}(p^m-p^{m-1})-4)}{12} \right\rceil, & \text{if } n \geq 4, m \geq 1, p \geq 2 \\ & \text{and } p^m - p^{m-1} \leq 4 \\ (p^n - p^{n-1}) \left\lceil \frac{(p^m-p^{m-1}-3)(p^m-p^{m-1}-4)}{12} \right\rceil \\ + 2 \left\lceil \frac{(p^{n-1}(p^m-p^{m-1})-3)(p^{n-1}(p^m-p^{m-1})-4)}{12} \right\rceil, & \text{if } n \geq 4, m \geq 1, p \geq 2 \\ & \text{and } p^m - p^{m-1} \geq 5. \end{array} \right.$$

*Proof.* By [10, Proposition 2.6] we have

$$CCC(G) = (p^n - p^{n-1})K_{p^{m-n}(p^n-p^{n-1})} \sqcup K_{p^{n-1}(p^m-p^{m-1})} \sqcup K_{p^{m-1}(p^n-p^{n-1})}.$$

Consider the following cases.

**Case 1.**  $n = 1$ .

We have  $CCC(G) = (p+1)K_{p^{m-1}(p-1)}$ . For  $m = 1$  and  $p = 2, 3$ , it follows that  $CCC(G) = 2K_1$  or  $3K_2$  which is planar. If  $m = 1$  and  $p \geq 5$ , by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = (p+1)\gamma(K_{p-1}) = (p+1) \left\lceil \frac{(p-4)(p-5)}{12} \right\rceil.$$

Clearly  $\gamma(CCC(G)) = 0$  for  $p = 5$ . If  $p \geq 7$  then

$$\frac{(p-4)(p-5)}{12} \geq \frac{1}{2}$$

and so

$$\gamma(CCC(G)) = (p+1) \left\lceil \frac{(p-4)(p-5)}{12} \right\rceil \geq 8.$$

If  $m = 2$  and  $p = 2$  then  $\gamma(CCC(G)) = 3\gamma(K_2) = 0$ . For  $m = 2$  and  $p \geq 3$ , by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = (p+1)\gamma(K_{p(p-1)}) = (p+1) \left\lceil \frac{(p^2-p-3)(p^2-p-4)}{12} \right\rceil.$$

If  $p \geq 3$  then

$$\frac{(p^2-p-3)(p^2-p-4)}{12} \geq \frac{1}{2}$$

and so

$$\gamma(CCC(G)) = (p+1) \left\lceil \frac{(p^2-p-3)(p^2-p-4)}{12} \right\rceil \geq 4.$$

If  $m = 3$  then  $\gamma(CCC(G)) = (p+1)\gamma(K_{p^2(p-1)})$ . Therefore, if  $m = 3$  and  $p \geq 2$  then by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = (p+1)\gamma(K_{p^2(p-1)}) = (p+1) \left\lceil \frac{(p^3-p^2-3)(p^3-p^2-4)}{12} \right\rceil.$$

Clearly if  $m = 3$  and  $p = 2$  then  $\gamma(CCC(G)) = 0$ . If  $p \geq 3$  then

$$\frac{(p^3-p^2-3)(p^3-p^2-4)}{12} \geq \frac{35}{2}$$

and so

$$\gamma(CCC(G)) = (p+1) \left\lceil \frac{(p^3-p^2-3)(p^3-p^2-4)}{12} \right\rceil \geq 72.$$

If  $m \geq 4$  and  $p \geq 2$  then  $\gamma(CCC(G)) = (p+1)\gamma(K_{p^{m-1}(p-1)})$ . Therefore, by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = (p+1)\gamma(K_{p^{m-1}(p-1)}) = (p+1) \left\lceil \frac{(p^m-p^{m-1}-3)(p^m-p^{m-1}-4)}{12} \right\rceil.$$

We have

$$\frac{(p^m-p^{m-1}-3)(p^m-p^{m-1}-4)}{12} \geq \frac{20}{12}$$

and so

$$\gamma(CCC(G)) = (p+1) \left\lceil \frac{(p^m-p^{m-1}-3)(p^m-p^{m-1}-4)}{12} \right\rceil \geq 6.$$

Therefore,  $CCC(G)$  is planar if and only if  $n = 1, m = 1, p = 2, 3, 5$ ;  $n = 1, m = 2, p = 2$ ; or  $n = 1, m = 3, p = 2$ . Also, in this case,  $CCC(G)$  is neither toroidal, double-toroidal nor triple-toroidal.

**Case 2.**  $n = 2$ .

We have  $CCC(G) = (p^2 - p)K_{p^{m-1}(p-1)} \sqcup 2K_{p^m(p-1)}$ . For  $m = 1$  and  $p = 2$ , it follows that  $CCC(G) = 2K_1 \sqcup 2K_2$  which is planar. If  $m = 1$  and  $p = 3$  then, by Lemma 2.1 and (1), we have

$$\gamma(CCC(G)) = 2\gamma(K_6) = 2.$$

If  $m = 1$  and  $p \geq 5$ , by Lemma 2.1 and (1), we have

$$\begin{aligned} \gamma(CCC(G)) &= (p^2 - p)\gamma(K_{p-1}) + 2\gamma(K_{p(p-1)}) \\ &= (p^2 - p) \left\lceil \frac{(p-4)(p-5)}{12} \right\rceil + 2 \left\lceil \frac{(p^2 - p - 3)(p^2 - p - 4)}{12} \right\rceil. \end{aligned}$$

Since  $p \geq 5$  then

$$\frac{(p^2 - p - 3)(p^2 - p - 4)}{12} \geq \frac{68}{3}$$

and so

$$\gamma(CCC(G)) \geq 2 \left\lceil \frac{(p^2 - p - 3)(p^2 - p - 4)}{12} \right\rceil \geq 46.$$

If  $m = 2$  and  $p \geq 2$  then  $CCC(G) = (p^2 - p)K_{p(p-1)} \sqcup 2K_{p^2(p-1)}$ . Therefore, if  $p = 2$  then  $CCC(G) = 2K_2 \sqcup 2K_4$  hence by (1) we have  $\gamma(CCC(G)) = 2\gamma(K_4) = 0$ . If  $p \geq 3$ , by Lemma 2.1 and (1), we have

$$\begin{aligned} \gamma(CCC(G)) &= (p^2 - p)\gamma(K_{p(p-1)}) + 2\gamma(K_{p^2(p-1)}) \\ &= (p^2 - p) \left\lceil \frac{(p^2 - p - 3)(p^2 - p - 4)}{12} \right\rceil + 2 \left\lceil \frac{(p^3 - p^2 - 3)(p^3 - p^2 - 4)}{12} \right\rceil. \end{aligned}$$

Also,  $\frac{(p^3 - p^2 - 3)(p^3 - p^2 - 4)}{12} \geq \frac{35}{2}$  and so

$$\gamma(CCC(G)) \geq 2 \left\lceil \frac{(p^3 - p^2 - 3)(p^3 - p^2 - 4)}{12} \right\rceil \geq 36.$$

If  $m \geq 3$  and  $p \geq 2$  then

$$\begin{aligned} \gamma(CCC(G)) &= (p^2 - p)\gamma(K_{p^{m-1}(p-1)}) + 2\gamma(K_{p^m(p-1)}) \\ &= (p^2 - p) \left\lceil \frac{(p^{m-1}(p-1) - 3)(p^{m-1}(p-1) - 4)}{12} \right\rceil + 2 \left\lceil \frac{(p^m(p-1) - 3)(p^m(p-1) - 4)}{12} \right\rceil \geq 4. \end{aligned}$$

Therefore,  $CCC(G)$  is planar if and only if  $n = 2, m = 1, p = 2$ ;  $n = 2, m = 2, p = 2$ ; or  $n = 3, m = 1, p = 2$  and double-toroidal if and only if  $n = 2, m = 1, p = 3$ . In this case,  $CCC(G)$  is neither toroidal nor triple-toroidal.

**Case 3.**  $n = 3$ .

We have  $CCC(G) = p^2(p-1)K_{p^{m-1}(p-1)} \sqcup 2K_{p^{m+1}(p-1)}$ . If  $m = 1$  and  $p = 2$  then  $CCC(G) = 4K_1 \sqcup 2K_4$ , and so by Lemma 2.1 and (1)  $\gamma(CCC(G)) = 2\gamma(K_4) = 0$ . For  $p = 3$  we have  $CCC(G) = 18K_2 \sqcup 2K_{18}$ . Therefore, by 2.1 and (1) we have  $\gamma(CCC(G)) = 2\gamma(K_{18}) = 36$ . For  $p \geq 5$ , by Lemma 2.1 and (1) we have

$$\begin{aligned} \gamma(CCC(G)) &= p^2(p-1)\gamma(K_{p-1}) + 2\gamma(K_{p^2(p-1)}) \\ &= p^2(p-1) \left\lceil \frac{(p-4)(p-5)}{12} \right\rceil + 2 \left\lceil \frac{(p^3-p^2-3)(p^3-p^2-4)}{12} \right\rceil > 36. \end{aligned}$$

If  $m = 2$  and  $p = 2$  then we have  $CCC(G) = 4K_2 \sqcup 2K_8$ . By Lemma 2.1 and (1) we have

$$\gamma(CCC(G)) = 2\gamma(K_8) = 4.$$

If  $m = 2$  and  $p \geq 3$  or  $m \geq 3$  and  $p \geq 2$  then we have  $CCC(G) = p^2(p-1)K_{p^{m-1}(p-1)} \sqcup 2K_{p^{m+1}(p-1)}$ . By Lemma 2.1 and (1) we have

$$\begin{aligned} \gamma(CCC(G)) &= p^2(p-1)\gamma(K_{p^{m-1}(p-1)}) + 2\gamma(K_{p^{m+1}(p-1)}) \\ &= p^2(p-1) \left\lceil \frac{(p^{m-1}(p-1)-3)(p^{m-1}(p-1)-4)}{12} \right\rceil \\ &\quad + 2 \left\lceil \frac{(p^{m+1}(p-1)-3)(p^{m+1}(p-1)-4)}{12} \right\rceil. \end{aligned}$$

We have

$$\frac{(p^{m+1}(p-1)-3)(p^{m+1}(p-1)-4)}{12} \geq \frac{5}{3}$$

and so

$$\gamma(CCC(G)) \geq 2 \left\lceil \frac{(p^{m+1}(p-1)-3)(p^{m+1}(p-1)-4)}{12} \right\rceil \geq 4.$$

Therefore,  $CCC(G)$  is planar if and only if  $n = 3, m = 1, p = 2$ . Also, in this case,  $CCC(G)$  is neither toroidal, double-toroidal nor triple-toroidal.

**Case 4.**  $n \geq 4$ .

We have  $CCC(G) = (p^n - p^{n-1})K_{p^m - p^{m-1}} \sqcup 2K_{p^{n-1}(p^m - p^{m-1})}$ . Therefore, by Lemma 2.1, we have

$$(2) \quad \gamma(CCC(G)) = (p^n - p^{n-1})\gamma(K_{p^m - p^{m-1}}) + 2\gamma(K_{p^{n-1}(p^m - p^{m-1})})$$

For  $m \geq 1$  and  $p \geq 2$  we have

$$\gamma(K_{p^{n-1}(p^m - p^{m-1})}) \geq \gamma(K_{p^{n-1}}) \geq \gamma(K_8) = 2,$$

noting that  $K_8$  and  $K_{p^{n-1}}$  are subgraphs of  $K_{p^{n-1}}$  and  $K_{p^{n-1}(p^m - p^{m-1})}$  respectively. Therefore

$$\gamma(CCC(G)) \geq 2\gamma(K_{p^{n-1}(p^m - p^{m-1})}) \geq 4.$$

Further, if  $p^m - p^{m-1} \leq 4$  then, by (2) and (1), we have

$$\begin{aligned} \gamma(\mathcal{CCC}(G)) &= 2\gamma(K_{p^{n-1}(p^m - p^{m-1})}) \\ &= 2 \left\lceil \frac{(p^{n-1}(p^m - p^{m-1}) - 3)(p^{n-1}(p^m - p^{m-1}) - 4)}{12} \right\rceil. \end{aligned}$$

If  $p^m - p^{m-1} \geq 5$  then, by (2) and (1), we have

$$\begin{aligned} \gamma(\mathcal{CCC}(G)) &= (p^n - p^{n-1}) \left\lceil \frac{(p^m - p^{m-1} - 3)(p^m - p^{m-1} - 4)}{12} \right\rceil + \\ &2 \left\lceil \frac{(p^{n-1}(p^m - p^{m-1}) - 3)(p^{n-1}(p^m - p^{m-1}) - 4)}{12} \right\rceil. \end{aligned}$$

Hence the result follows.  $\square$

We conclude this paper with the following characterization of  $\mathcal{CCC}(G)$  for the groups listed in Section 1.

**Corollary 2.8.** *Let  $G = D_{2n}, SD_{8n}, Q_{4m}, V_{8n}, U_{(n,m)}$  or  $G(p, m, n)$ . Then*

- (1)  $\mathcal{CCC}(G)$  is planar if and only if  $G = D_6, D_8, D_{10}, D_{12}, D_{14}, D_{16}, D_{18}, D_{20}, SD_{16}, SD_{24}, Q_8, Q_{12}, Q_{16}, Q_{20}, V_{16}, U_{(2,2)}, U_{(2,3)}, U_{(2,4)}, U_{(2,5)}, U_{(2,6)}, U_{(3,2)}, U_{(3,3)}, U_{(3,4)}, U_{(4,2)}, U_{(4,3)}, U_{(4,4)}, G(2, 1, 1), G(3, 1, 1), G(5, 1, 1), G(2, 2, 1), G(2, 3, 1), G(2, 1, 2), G(2, 2, 2)$  or  $G(2, 1, 3)$ .
- (2)  $\mathcal{CCC}(G)$  is toroidal if and only if  $G = D_{22}, D_{24}, D_{26}, D_{28}, D_{30}, D_{32}, SD_{32}, Q_{24}, Q_{28}, Q_{32}, V_{24}, V_{32}, U_{(2,7)}, U_{(2,8)}, U_{(3,5)}$  or  $U_{(3,6)}$ .
- (3)  $\mathcal{CCC}(G)$  is double-toroidal if and only if  $G = D_{34}, D_{36}, SD_{40}, Q_{36}, U_{(2,9)}, U_{(2,10)}, U_{(4,5)}, U_{(4,6)}, U_{(5,2)}, U_{(5,3)}, U_{(6,2)}, U_{(6,3)}, U_{(7,2)}, U_{(7,3)}$  or  $G(3, 1, 2)$ .
- (4)  $\mathcal{CCC}(G)$  is triple-toroidal if and only if  $G = D_{38}, D_{40}, Q_{40}, V_{40}, U_{(3,7)}, U_{(3,8)}, U_{(5,4)}, U_{(6,4)}$  or  $U_{(7,4)}$ .

### 3. ACKNOWLEDGMENT

The authors would like to thank the referee for his/her valuable comments and suggestions. The first author is thankful to Council of Scientific and Industrial Research for the fellowship (File No. 09/796(0094)/2019-EMR-I).

### REFERENCES

- [1] M. Afkhami, M. Farrokhi D. G. and K. Khashyarmansh, *Planar, toroidal, and projective commuting and non-commuting graphs*, Comm. Algebra, **43** No. 7 (2015) 2964-2970.
- [2] P. Bhowal and R. K. Nath, *Spectral aspects of commuting conjugacy class graph of finite groups*, Alg. Struc. Appl., **8** No. 2 (2021) 67-118.

- [3] P. Bhowal and R. K. Nath, *Spectrum and energies of commuting conjugacy class graph of a finite group*, available at <https://arxiv.org/abs/2003.07142>.
- [4] P. Bhowal, D. Nongsiang and R. K. Nath, *Solvable graphs of finite groups*, Hacet. J. Math. Stat., **49** No. 6 (2020) 1955-1964.
- [5] R. Brauer and K. A. Fowler, *On groups of even order*, Ann. Math., **62** No. 2 (1955) 565-583.
- [6] A. K. Das and D. Nongsiang, *On the genus of the commuting graphs of finite non-abelian groups*, Int. Electron. J. Algebra, **19** (2016) 91-109.
- [7] A. K. Das and D. Nongsiang, *On the genus of the nilpotent graphs of finite groups* Comm. Algebra, **43** No. 1 (2015) 5282-5290.
- [8] M. Herzog, M. Longobardi and M. Maj, *On a commuting graph on conjugacy classes of groups*, Comm. Algebra, **37** No. 10 (2009) 3369-3387.
- [9] A. Mohammadian, A. Erfanian, D. G. M. Farrokhi and B. Wilkens, *Triangle-free commuting conjugacy class graphs*, J. Group Theory, **19** (2016) 1049-1061.
- [10] M. A. Salahshour and A. R. Ashrafi, *Commuting conjugacy class graph of finite CA-groups*, Khayyam J. Math., **6** No. 1 (2020) 108-118.
- [11] J. Battle, F. Harary, Y. Kodama and J. W. T. Youngs, *Additivity of the genus of a graph*, Bull. Amer. Math. Soc., **68** No. 6 (1962) 565-568.
- [12] A. T. White, *Graphs, Groups and Surfaces*, North-Holland Mathematics Studies, No. 8., American Elsevier Publishing Co., Inc., New York, 1973.

**Parthajit Bhowal**

Department of Mathematical Sciences

Tezpur University, Napaam-784028, Sonitpur, Assam, India.

Department of Mathematics, Cachar College, Silchar-788001, Assam, India.

bhowal.parthajit8@gmail.com

**Rajat Kanti Nath**

Department of Mathematical Sciences

Tezpur University, Napaam-784028, Sonitpur, Assam, India.

rajatkantinath@yahoo.com