



Study of relationship between related yield traits using correlation and regression in wild barley (*Hordeum murinum*)

Hossein Mehripour^{1*}, Abolfazl Rashidi², Mehdi Zahravi³ and Asa Ebrahimi⁴

¹M. Sc. Plant Breeding, Science and Research branch, Islamic Azad University, Tehran, Iran

²Assistant Professor of plant Breeding, Islamic Azad University, Shahre Rey Branch

³National Plant Gene Bank of Iran, Karaj, Iran

⁴Assistant Professor of Biotechnology, Science and Research branch, Islamic Azad University, Tehran, Iran

ABSTRACT

Genotypic variation is useful to breeders when selecting genotypes to improve particular traits. In order to study of genetic diversity 20 genotypes of *Hordeum murinum* studied in the form of randomized complete block design the experimental field of Islamic Azad University Shahre Rey Branch with 3 replication in the year 2012. Notes traits was conducted including days to heading, days to maturity, plant height, straw weight, harvest index, grain number per panicle, 10- panicle weight, 100- grain weight, biomass and grain yield. The results of analysis of variance showed that the significant differences were observed among genotypes for all morphological traits, except days to maturity and harvest index. Mean of traits classified using Duncan's multiple range test ($p = 0.05$). Line of 12-1092 produced the highest Plant height, days to heading, 10- panicle weight straw weight, biomass and grain yield. Correlation coefficients showed the grain yield has a positive at $p = 0.05$ and significant with biomass ($r=0.97$), number of nodes on the stem ($r=0.72$), 10- panicle weight (gr) and plant height. Stepwise regression analysis, grain yield as the dependent variable (Y) and other traits evaluated was considered as an independent variable (X). Results showed that the biomass and grain yield (with biomass) with 93.7% and 98.8% explained the maximum grain yield variation, respectively.

Key word; Barley, Correlation, Grain yield, *Hordeum murinum* and Regression.

INTRODUCTION

Barley is an important crop in the world, ranking fifth in the world production that is used for animals, malt, and human food (Khodabandeh, 2002). Its importance derives from the ability to grow and produce in marginal environments, which are often characterized by drought, low temperature and salinity (Hayes *et al*, 2003; Baum *et al*, 2003). Wild barley possess high genetic variation in several useful characters including earliness, biomass yield, protein content, resistance against powdery mildew and leaf rust (Nevo, 1992). *Hordeum* L. is a widely distributed genus of the tribe Triticeae of the Poaceae (Graminae) family. There are about 45 species and subspecies, most of which represent weedy annual or perennial grasses, found throughout the temperate zones of both northern and southern hemispheres (Morrell,

2003). The large genetic variability present in the wild cereals is an invaluable resource for cereal crop improvement. Assessment of the genetic diversity in a crop species is fundamental to its improvement. Genetic diversity among and within plant species is in danger of being reduced. In wild species genetic diversity may be lost because of severe reduction in population size, whereas in domesticated crops genetic diversity may be lost because of the narrow genetic base in many breeding programs (Kling *et al*, 2003; Cao *et al*, 1998). Estimates of genetic diversity can be based on different types of data. Phonological and morphological quantitative traits have frequently been used for studying genetic diversity in barley (Chand *et al*, 2008). Neyestani *et al* (2005) in study of 10 barley cultivars estimated that the correlation between the numbers of grains per spike with grain yield was positive. The purpose of this experiment was to determine the proportion of traits in determination of grain yield.

MATERIALS AND METHODS

In order to study of genetic diversity 20 genotypes of *Hordeum murinum* studied in the form of randomized complete block design the experimental field of Islamic Azad University Shahre Rey branch with 3 replication in the year 2012. Notes traits was conducted including plant height (cm), days to heading, number of grain per panicle, days to maturity, 10- panicle weight (gr), 100- grain weight (gr), harvest index, straw weight (gr), biomass (biological yield) (gr) and grain yield (gr.m^{-2}). Plants selected for sampling randomly within each block were experimental plots. To calculate the correlation coefficient of the mean traits were statistically analyzed for each experimental unit, also in order to evaluate the effect of reducing the number of independent variables and the fitted regression was used model, stepwise regression. Finally statistical calculations were performed using software SAS and SPSS₁₆.

RESULTS AND DISCUSSION

The results of analysis of variance (Table 1) showed that the significant differences were observed among genotypes for all morphological traits, except days to maturity and harvest index. Drikvand *et al* (2012) in Study of Genetic diversity among Barley Genotypes showed significant differences were observed among for morphological traits. Mean of traits classified using Duncan's multiple range test ($p = 0.05$). Line of 12-1092 produced the highest plant height (gr), days to heading, 10-panicle weight (gr) straw weight, biomass (gr) and grain yield (gr.m^{-2}). Number of grain in panicle and 100- grain weight was greatest in 1-826 and 6-986 lines, respectively (Table 2). These results showed that different genetic systems involved in controlling traits, which emphasized on important of study of these traits (Maktoobian *et al*, 2013; Khajavi *et al*, 2014). Among descriptive parameters studied (Table 3) the highest coefficient of variation was the grain yield (49.01%), biomass (40.71%) and straw weight (39.63%), thus, range of traits is widely in genotypes. The lowest coefficient of variation was the days to maturity (1.32%) and days to heading (1.52%). Therefore, these traits have not suitable diversity in crop improvement. Correlation coefficients (Table 4) showed the grain yield higher for the same amount of biomass and straw weight, the number of nodes in the stem, height and weight is the most 10- panicle weight. The grain yield has a positive at $p = 0.05$ and significant with biomass ($r=0.97$), number of nodes on the stem ($r=0.72$), 10- panicle weight (gr) and plant height (cm), also the increase in each of these characteristics, increase grain yield. plant height had a significant and positive correlation with the straw weight ($r=0.72$), biomass ($r=0.68$), number of grains per panicle ($r = 0.50$) and grain yield ($r=0.57$). Therefore, Plant height is perfect for select genotypes with high yield. Furthermore, the use of straw for livestock feed, selection this trait can be a significant impact increase straw (Baniya *et al.*, 1967; Zaheer., 2008). Kole (2006) reported a significant grain yield per plant has positive correlations with, number of tillers, number of spikelet's spike and 100- grain grains weight. According to the same reports and contradictory results, it is obvious that determining the correlation rate of yield. The significant coefficient in the successful regression equation indicating these attributes are to be effective in increasing yield. In other words, with the increase of this specification, yield will also increase. Afzali Far *et al* (2011) according to the stepwise regression analysis traits such as total grain yield, biomass and plant height introduced as an effective traits on the yield. Dadashi *et al* (2010) using

stepwise regression and at the 5% level three traits such as grain per spike, number of fertile tillers and seed weight introduced as an effective traits on the yield.

REFERENCES

- Afzali Far A., Zahravi M., Bihamta, M. (2011). Evaluation of tolerant genotypes to drought stresses the barley *spantanyum* iran in Karaj region. Journal of Agronomy and Plant Breeding, 7: 44-25.
- Baniya B, 1972. Crop Production in Dry Regions, Vol, 2. Leonard Hill, London.
- Baum M., Grando S., Backes G., Jahoor A., Sabbagh A., Ceccarelli, S. (2003). QTLs for agronomic traits in the Mediterranean environment identified in recombinant inbred lines of the cross 'Arta' x. spontaneum 41-1. Theor Appl Genet., 107, 1215–1225.
- Cao W G., Hucl P S G., Chibbar, R N. (1998). Genetic diversity within spelta and macha wheats based on RAPD analysis. Euphytica, 104: 181-189.
- Chand N., Vishwakarma S R., Verma O P, Kumar, M. (2008). Worth of genetic parameters to sort out new elite barley lines over heterogeneous environments. Barley Genet. Newslett. 38: 10-13.
- Dadashi M., Norinia, A A., MortezaAsghar M., Azizi Chakhrchaman, Sh. (2010). Evaluate the correlation a number of physiological and morphological characteristics of naked barley varieties with grain yield. Journal of Eco-physiology of crop plants and weeds. 15: 29-40.
- Drikvand R., Salahvarzi E., Salahvarzi A., Hossinpour, T. (2012). Study of Genetic Diversity among Rainfed Barley Genotypes Using ISJ Markers and Morphological Traits. Journal of Agricultural Science; 4: ISSN 1916-9752 E-ISSN 1916-9760.
- Golabady M., Arzani, A. (2003). Study of Genetic variation and Factor Analysis of Agronomic Traits in Durum wheat. Journal of Science Technology of Agriculture and Natural Resources, 1: 115-126.
- Hayes PM., Castro A., Marquez-Cedill L. A., Corey C., Henson B., Jones J., Kling D., Mathero., Matus I., Rossi C., Sato, K. (2002). Genetic diversity for quantitatively inherited agronomic and malting quality traits. In Diversity in Barley. R. von Bothmer and T. van Hintum (Eds.).
- Khajavi A, Aharizad S, Ahmadizadeh M.2014. Genetic diversity of promising lines of barley based on pheno-morphological traits in Ardabil area. International journal of Advanced Biological and Biomedical Research Volume 2, Issue 2, 2014: 456-462.
- khanghah A M., Yousef A., Moosavi, S S. (2014). Investigate the relationship between studied traits with grain yield using regression analysis and path analysis 34 barley lines and cultivars. International Journal of Plant, Animal and Environmental Sciences. 4: ISSN 2231-4490.
- Khodabandeh, N. (2003). Cereal crops. Seventh Edition. Tehran University Press.
- Kling J., Mather D., Matus I., Rossi C., Sato, K. 2003. Genetic diversity for quantitatively inherited agronomic and malting quality traits. In R. Von Bothmer et al. (ed.) Diversity in barley (*Hordeum vulgare*). Elsevier Science Publishers, Amsterdam.
- Kole, P C. (2006). Variability, Correlation and Regression Analysis in Third Somaclonal Generation of Barley. Barley Genetics Newsletter 36:44-47.

Maktoobian M., Soleymani A., Javanmard, H R. (2013). Drought Stress Effects on Seed Yield and Yield Components of Barley Cultivars. International Journal of Agronomy and Plant Production. 4: 3415-3418.

Morrell P L., Lundy K E., Clegg M T., (2003). Distinct geographic patterns of genetic diversity are maintained in wild barley (*Hordeum vulgare* ssp. *spontaneum*) despite migration. Proc. Natl. Acad. Sci U.S.A., 100:10812-10817.

Nevo, E. (1992). Origin, evolution, population genetic and resources for breeding of wild barley *Hordeum spontaneum*. In: Barley: Genetics. Biochemistry Molecular Biology and Biotechnology, (Ed.): PR Shewry, CAB international. Wallingford 19-143.

Neyestani, E., Mahmoodi, A A. and Rahimnia, F. (2005). Path analysis of grain yield and its components and estimation of heritability in barley. Journal of Agricultural. 2; 55-63.

Zaheer A., Ajmal, S. U., Munir, M., Zubair, M., Masood, M S (2008). Genetic diversity for morpho-genetic traits in barley germplasm. Pakistan Journal of Botany 40: 1217-1224.

Archive of SID

Table 1. Analysis of variance different traits in 20 barely genotypes (n=20).

| S.O.V | df | Plant height (cm) | Days to heading | Number of grain in panicle | Days to maturity | 10-Panicle weight | 100-Grain weight (gr) | Harvest index | straw weight (gr) | Biomass (gr) | Grain yield (gr.m ⁻²) |
|-------------|----|-------------------|-----------------|----------------------------|------------------|-------------------|-----------------------|---------------|-------------------|--------------|-----------------------------------|
| Genotype | 19 | 55.76** | 18.94** | 40.12** | 9.01 | 1.08** | 0.105** | 40.86 | 2899.65** | 6284.69** | 709.39* |
| Replication | 2 | 216.6** | 30.21** | 70.11** | 23.21* | 0.44 | 0.006 | 69.88 | 8135.39** | 17638.52** | 1867.82** |
| Error | 38 | 20.84 | 5.05 | 4.36 | 5.41 | 0.28 | 0.03 | 42.04 | 1054.36 | 2324.66 | 320.10 |

* and **; Significant at $p = 0.05$ and $p = 0.01$ levels, respectively.**Table 2.** Means of the estimated traits in 20 barely genotypes (n=20).

| Lines | Plant height (cm) | Days to heading | Number of grain in panicle | Days to maturity | 10-Panicle weight (gr) | 100-Grain weight (gr) | Harvest index | Straw weight (gr) | Biomass (gr) | Grain yield (gr.m ⁻²) |
|---------|-------------------|-----------------|----------------------------|------------------|------------------------|-----------------------|---------------|-------------------|--------------|-----------------------------------|
| 1-826 | 28.16af | 147.33be | 38a | 177.33ac | 3.01bf | 0.86df | 28.96ab | 86.27b | 122.03bc | 35.75bc |
| 2-828 | 27.66bf | 143.67de | 34ac | 175.33ac | 2.51df | 0.82f | 34.45ab | 84.93b | 130.8bc | 45.86bc |
| 3-833 | 24.16df | 148bd | 34.33ac | 178ab | 2.50df | 0.75f | 31.44ab | 63.25b | 91.44bc | 28.20bc |
| 4-943 | 31.33ae | 147.33be | 34.33ac | 175ac | 3.12be | 0.80f | 27.69ab | 93.57b | 130.28bc | 36.70bc |
| 5-985 | 26bf | 145.33be | 25e | 176.33ac | 3.21be | 1.27ab | 27.39ab | 48.28b | 68.36bc | 20.08bc |
| 6-986 | 20.66f | 145.33be | 24.33e | 178.33ab | 3.51bc | 1.36a | 27.62ab | 54.61b | 76.77bc | 22.15bc |
| 7-992 | 31.33ae | 143.67de | 31.33c | 174.33ac | 3.78b | 0.84ef | 35.28ab | 104.01b | 156.19b | 52.02bc |
| 8-1004 | 34.33ab | 144.33ce | 35.66ab | 173c | 2.42ef | 0.88cdf | 28.41ab | 106b | 146.01bc | 40bc |
| 9-1021 | 30.66ae | 148bd | 34.33ac | 173c | 3.4bd | 1.21ac | 30.58ab | 83.09b | 119.57bc | 36.47bc |
| 10-1085 | 24.66cf | 144.33ce | 27.33de | 175ac | 3.02bf | 1.18ae | 31.06ab | 62.24b | 91.34bc | 29.1bc |
| 11-1086 | 33.5ac | 143.33e | 34ac | 173c | 2.41ef | 0.89cf | 25.39b | 86.16b | 120.33bc | 34.16bc |
| 12-1092 | 36.66a | 153a | 32.66bc | 178.67a | 4.77a | 1.08af | 29.18ab | 190.34a | 276.68a | 86.34a |
| 13-1096 | 33.33ac | 148.33bc | 36.66ab | 176.67ac | 3.30be | 0.94bf | 30.76ab | 92.75b | 132.63bc | 39.88bc |
| 14-1146 | 31.66ad | 144.67be | 36.66ab | 175.33ac | 3.08be | 0.87cf | 30.46ab | 79.59b | 114.56bc | 34.97bc |
| 15-1171 | 22.33ef | 149.33ab | 30.66cd | 176ac | 3.77b | 1.19ad | 30.06ab | 51.74b | 77.05bc | 25.31bc |
| 16-1174 | 25.33bf | 149ab | 33bc | 175.33ac | 3.04bf | 0.90cf | 38.67a | 81.97b | 134.23bc | 52.25b |
| 17-1185 | 29.66ae | 149.67ab | 31.33c | 175.67ac | 2.16f | 0.77f | 24.9b | 43.22b | 59.58c | 16.36c |
| 18-1187 | 26bf | 146.67be | 33.66bc | 174bc | 2.78cf | 1.01bf | 33.91ab | 80.91b | 122.41bc | 41.5bc |
| 19-1199 | 26.33bf | 149.67ab | 34.66ac | 175.33ac | 3.04bf | 0.83f | 28.26ab | 76.88b | 108.09bc | 31.2bc |
| 20-1205 | 25.16cf | 148.33bc | 34.66ac | 177.67ab | 2.6df | 0.76f | 23.35b | 68.46b | 90.23bc | 21.76bc |

In each column, any two means having a common letter are not significantly different at $p = 0.05$ based on Duncan's multiple range test.

Table 3. Descriptive parameters of morphological traits in 20 barely genotypes (n=20).

| traits | Plant height (cm) | Days to heading | Number of grain in panicle | Days to maturity | 10-Panicle weight | 100-Grain weight (gr) | Harvest index | Straw weight (gr) | Biomass | Grain yield (gr.m ⁻²) |
|-------------------|-------------------|-----------------|----------------------------|------------------|-------------------|-----------------------|---------------|-------------------|---------|-----------------------------------|
| Mean | 28.45 | 147.08 | 32.83 | 175.67 | 3.07 | 0.96 | 29.89 | 81.92 | 118.43 | 36.5 |
| Minimum | 20.67 | 143.33 | 24.33 | 173 | 2.17 | 0.76 | 23.36 | 43.22 | 69.59 | 16.36 |
| Maximum | 36.67 | 153 | 38 | 178.67 | 4.77 | 1.36 | 38.68 | 190.34 | 276.68 | 86.34 |
| Rang of variation | 16 | 9.67 | 13.67 | 5.67 | 2.6 | 0.61 | 13.32 | 147.12 | 217.09 | 69.98 |
| SD | 431.38 | 2.51 | 3.65 | 1.73 | 0.6 | 0.18 | 3.69 | 310.88 | 457.69 | 153.77 |
| ‡ CV (%) | 16.04 | 1.52 | 6.35 | 1.32 | 14.98 | 17.74 | 21.68 | 39.63 | 40.71 | 49.01 |

‡ Coefficient of variation

Archive of SID

Table 4. Correlation of traits in 20 barely genotypes (n=20).

| | 10- Panicle weight | 100- Grain weight | Number of grain in panicle | Grain yield (gr.m ⁻²) | Biomass (gr) | Harvest index | Straw weight (gr) | Plant height(cm) | Days to heading | Days to maturity | Number of nodes on the stem |
|-----------------------------------|--------------------|-------------------|----------------------------|-----------------------------------|--------------|---------------|-------------------|------------------|-----------------|------------------|-----------------------------|
| 10- Panicle weight (gr) | 1 | | | | | | | | | | |
| 100- Grain weight(gr) | 0-.33 | 1 | | | | | | | | | |
| Number of Grain in panicle | 0-.24 | 0-.13 | 1 | | | | | | | | |
| Grain yield (gr.m ⁻²) | 0.58** | 0-.31 | 0.27 | 1 | | | | | | | |
| Biomass(gr) | 0.57** | 0-.31 | 0.33 | 0.97** | 1 | | | | | | |
| Harvest index | .20 | 0-.32 | 0.07 | 0.47* | 0.27 | 1 | | | | | |
| Straw weight(gr) | 0.55* | 0-.30 | 0.35 | 0.93** | 0.99** | 0.16 | 1 | | | | |
| Plant height (cm) | 0.14 | 0.04 | 0.50* | 0.57** | 0.68** | 0-.10 | 0.72** | 1 | | | |
| Days to heading | 0.40 | 0.24 | 0.20 | 0.26 | 0.28 | 0-.11 | 0.28 | 0.05 | 1 | | |
| Days to maturity | 0.34 | 0.005 | 0-.19 | 0.03 | 0.05 | 0-.19 | 0.06 | 0-.31 | 0.49* | 1 | |
| Number of nodes on the stem | 0.68** | 0-.07 | 0-.07 | 0.72** | 0.73** | 0.21 | 0.71** | 0.44 | 0.17 | 0.16 | 1 |

* and **; Significant at $p = 0.05$ and $p = 0.01$ levels, respectively.

Table 5. Stepwise regression for grain yield (dependent variable) and other traits (independent variable)

| stage | Cumulative coefficient | F | The final coefficient |
|-------|------------------------|----------|-----------------------|
| 1 | 93.7 | 282.32** | 0.32 |
| 2 | 98.8 | 769.90** | 0.95 |

| stage | trait | Width of source | The regression coefficient | | Cumulative coefficient |
|-------|---------------|-----------------|----------------------------|------|------------------------|
| | | | F | E | |
| 1 | Biomass | -2.07 | - | 0.32 | 93.7 |
| 2 | Harvest index | -28.24 | 0.95 | 0.30 | 98.8 |

$$y = -28.24 + 0.30 X_1 + 0.95 X_2$$