



# Journal of Human, Environment, and Health Promotion

Journal homepage: [www.zums.ac.ir/jhehp](http://www.zums.ac.ir/jhehp)



## Evaluation of the Prevalence of Thalassemia and Sickle Hemoglobin in Marriage Applicants Referring to the Genetic Disease Counseling Center in Kazeroun, Iran during 2014-2018

Mohsen Moghadami <sup>a</sup> | Masoud Sheybani <sup>b</sup> | Abbas Reziaean Zadeh <sup>c</sup> | Tayebbeh Rakhshani <sup>d\*</sup>

*a. Non Communicable Disease Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.*

*b. Community Education, School of Public Health, Shiraz University of Medical Sciences, Shiraz, Iran.*

*c. Departemant of Epidemiology, Faculty of Public Health, Shiraz Univercity of Medical Science, Shiraz, Iran.*

*d. Nutrition Research Center, Department of Public Health, School of Public Health, Shiraz University of Medical Sciences, Shiraz, Iran.*

**\*Corresponding author at:** Nutrition Research Center, Department of Public Health, School of Public Health, Shiraz University of Medical Sciences, Shiraz, Iran. Postal code: 734411639.  
E-mail address: [healthacademic2014@gmail.com](mailto:healthacademic2014@gmail.com)

### ARTICLE INFO

**Article type:**  
Original article

**Article history:**  
Received: 14 August 2019  
Revised: 12 October 2019  
Accepted: 23 October 2019

DOI: [10.29252/jhehp.5.4.7](https://doi.org/10.29252/jhehp.5.4.7)

**Keywords:**  
Prevalence  
Thalassemia  
Sickle cell anemia  
Marriage applicants

### ABSTRACT

**Background:** The present study aimed to determine the prevalence of thalassemia minor and sickle hemoglobin in marriage applicants referring to the Genetic Diseases Counseling Center of Kazeroun Health Center over 2014-2018.

**Methods:** This cross-sectional study was conducted in Kazeroun, Iran. The statistical population included all couples referring to the marriage counseling centers in Kazeroun for pre-marriage tests during 2014-2018. In this study, the registration data associated with the health records of the couples seeking marriage counseling were used.

**Results:** Over the years under scrutiny, 31114 applicants (15557 couples) referred to the health counseling center for marriage tests under the supervision of the health network. Among the marriage applicants, the highest prevalence of beta-thalassemia, alpha-thalassemia, anemia, sickle cell hemoglobin, and hemoglobinopathy was reported in 2014 (0.07), 2018 (0.15), 2014 (0.66), 2018 (0.9), and 2014 (0.05), respectively.

**Conclusion:** The results of the present study showed that the prevalence of thalassemia and sickle cell anemia in Kazeroun over the study years was similar to that in other parts of Iran over the same years.

## 1. Introduction

Thalassemia is a type of chronic hemolytic anemia caused by a defect in the production of one or more of the hemoglobin chains [1]. Thalassemia is inherited from parents to children [2,3]. The distortion of red blood cells into sickle shape is a disorder caused by a mutation in the beta gene; accordingly, amino acid valine substituted for glutamic amino acid [4,5] as blood fails to reach the tissues in position 6 of the beta chain. Sickle cell disease is either an HbSS homozygote or associated with HbS with other hemoglobin variants such as SC and SD or beta-thalassemia [6]. Thalassemia is diagnosed through iron deficiency by measuring hemoglobin A2, serum iron levels, TIBC, and

ferritin, since the hemoglobin A2 levels in these patients are higher than the normal subjects [7]. Thalassemia is more common in the Mediterranean and equatorial countries or in the areas near to the equator and in Africa [8]. In other words, the thalassemia belt spreads to the region of the Mediterranean Sea along the Arabian Peninsula, Turkey, Iran, India, and Southeast Asia, especially Thailand, Cambodia, and South China.

The prevalence of the thalassemia gene in these areas is between 2.5 and 15.5% [9]. In Iran, thalassemia is more common in areas around the Caspian Sea, the Persian Gulf, and the Oman Sea, i.e., Mazandaran, Gilan, Khuzestan, Fars, Bushehr, Hormozgan, Sistan-Baluchestan, and Kerman provinces [10].



This work is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported License which allows users to read, copy, distribute and make derivative works for non-commercial purposes from the material, as long as the author of the original work is cited properly.

In Iran, the prevalence of thalassemia genes varies from 3% to 4% by geographical region. It is estimated that approximately two million people carry the beta-thalassemia gene in Iran. Recognizing the carriers of this disease before marriage in areas with high prevalence rate, especially Fars as one of the ten most populated provinces in the country, and precluding marriage between carriers through raising awareness can prevent the birth of people with homozygous thalassemia (major), resulting in significant economic and social damages to the community [11]. Regarding the prevalence of heterozygote type, sickle cell anemia, especially in southern Iran, and the probable association between sickle cell disease and beta-thalassemia, it can lead to sickle cell thalassemia and severe socio-physical problems. Moreover, it should be noted that if the prevalence of sickle cell and  $\beta$ -thalassemia minor is equal in an area, the incidence of sickle cell syndromes is twice that of thalassemia major, and unfortunately, the implementation of thalassemia prevention programs not only fail to diminish the incidence of sickle cell syndromes but also relatively increase it [12,13]. Kazeroun is geographically known as one of the cities with a high prevalence of  $\beta$ -thalassemia and sickle cell (S). Therefore, the authors decided to investigate the prevalence of minor thalassemia and sickle cell hemoglobin in marriage applicants referring to the Genetic Diseases Counseling Center of Kazeroun Health Center, Shiraz University of Medical Sciences during 2014-2018.

## 2. Materials and Methods

### 2.1. Study Population

This cross-sectional study was carried out in Kazeroun during 2014-2018. All couples referring to the counseling center of Kazeroun for pre-marriage tests during 2014-2018 were included in this study. The inclusion criterion was applicants with a record in this case, and the exclusion criterion was the records with incomplete information.

### 2.2. Sampling Method

All the couples referring to the marriage counseling centers in Kazeroun for pre-marriage tests during 2014-2018 were included in the study based on the census.

### 2.3. Data Collection Instrument

In this study, data in couples' health records were used for marriage counseling. The marriage applicants were referred to screening laboratories for the detection of thalassemia following referral to the marriage registry office. After thalassemia tests, the results were sent to the counseling health center for interpreting. The counseling physician was responsible for interpreting the tests based on a national algorithm. Further experiments included chains examination, hemoglobin H examination, and DNA study. The results of the above experiments were sent to the academic advisor of the program for final interpretation. According to the diagnosis of the academic advisor, these

people were detected to be the carrier couples; hence, the necessary measures were taken based on the plan.

### 2.4. Data Analysis

Data were analyzed by SPSS V.20, and the normality of the data was assessed by the Kolmogorov-Smirnov test. After confirming the normality of the data, statistical analysis was performed using ANOVA and Chi-square tests.

## 3. Result and Discussion

During the 5 years of the study, 31114 applicants (15555 couples) referred to the health consulting center under scrutiny for marriage tests. The highest number of marriage applicants was in 2014 with a total of 7748 persons, and the lowest number was in 2018 with a total of 4872 persons. (Table 1).

According to Table 2, ANOVA results showed that there is a significant difference among the study years in terms of the average volume of red blood cells (80-96 normal), the mean of red blood cells, and red blood cells in the study years ( $P < 0.001$ ). However, there was no significant difference in hemoglobin levels among the study years ( $P = 0.16$ ).

Table 3 shows the prevalence of thalassemia, anemia, sickle cell hemoglobin, and hemoglobinopathy in different study years. In marriage applicants, the highest prevalence of beta-thalassemia, alpha thalassemia, anemia, sickle cell hemoglobin, and hemoglobinopathy was reported in 2014 (0.07), 2018 (0.15), 2014 (0.06), 2018 (0.09), and 2014 (0.05), respectively. In the total population, the highest prevalence of beta- and alpha-thalassemia was in 2014, the highest prevalence of anemia was in 2014, the highest incidence of sickle cell hemoglobin was in 2015, and the highest frequency of hemoglobinopathy was in 2014 (Table 3).

Chi-square test showed a significant difference between the study years in terms of the prevalence of thalassemia, anemia, sickle cell hemoglobin, and hemoglobinopathy ( $P < 0.001$ ). In 2014, thalassemia, anemia, and hemoglobinopathy had a higher rate than other study years while in 2015, sickle hemoglobin had a higher rate than the other study years (Table 4).

The aim of this study was to determine the prevalence of thalassemia and sickle hemoglobin in marriage applicants referring to the Genetic Disease Counseling Center of the Health Center of Kazeroun over 2014-2018. The results of the study showed that the prevalence of beta-thalassemia in marriage applicants was 39% in all study years (7% in 2014, 6.7% in 2015, 6.6% in 2016, 6.5% in 2017, and 6.5% in 2018).

The prevalence of thalassemia was 5.5% in a study conducted by Forouzan Nejad et al. in Ahwaz in 2018 [14].

**Table 1:** The number of marriage volunteers in study years

Year	Marriage applicants
2014	7748 (3874 couples)
2015	6732 (3366 couples)
2016	5972 (2986 couples)
2017	5790 (2895 couples)
2018	4872 (2436 couples)
2014-2018	31114 (15557 couples)

**Table 2:** Determining the mean and standard deviation of blood indices in marriage volunteers

Blood indices	2014	2015	2016	2017	2018	Pvalue
	Standard deviation ± mean	Standard deviation ± mean	Standard deviation ± mean	Standard deviation ± mean	Standard deviation ± mean	
RBC medium volume	85.35 ± 12.73	84.95 ± 10.33	84.41 ± 6.92	84.98 ± 10.78	84.00 ± 7.07	<0.001
Mean hemoglobin of the RBC	27.7 ± 3.49	27.64 ± 3.05	27.75 ± 3.07	27.86 ± 3.09	27.60 ± 3.09	<0.001
Hemoglobin	14.35 ± 3.80	14.37 ± 2.56	14.28 ± 1.71	14.36 ± 1.74	14.39 ± 1.74	<0.001
RBC	5.21 ± 0.64	5.22 ± 0.82	5.18 ± 0.62	5.19 ± 0.64	5.24 ± 0.65	< 0.001

In a study conducted in Pakistan in 2015, the prevalence of thalassemia was 6.5% [15]. In the study by Sherma et al. in India, the prevalence of beta-thalassemia was reported to be 78.2% [16], and in a study in China in 2017, it was 21.2% [17]. In another study, the prevalence of beta-thalassemia was reported to be 5.6% in Kashmir [18], 3.4% in Hong Kong, and 3.4% in Saudi Arabia [19]. These findings were somewhat consistent with the results yielded in the present study on the prevalence of thalassemia major over the years studied.

The results of the current study showed that the prevalence of alpha thalassemia in marriage applicants was 0.58 (58% in all study years: 1.4% in 2014, 1.5% in 2015, 1.4% in 2016, 1.4% in 2017, and 1.5% in 2018). In the study by Forouzan Nejad et al., the prevalence of α-thalassemia was 65.6% in Ahvaz [14]; in the study by Valizadeh et al., the prevalence of alpha thalassemia among marriage applicants was 53.3% [20]; in the study by Zandi et al., in Khuzestan, it was 9% [21], and in the southern regions of Iran, it was over 30% [22]. In the study by Alkara et al. in Emirate, the respective prevalence was 49% [23], and in the study by Borgrus et al. in Brazil, it was 42.8% [24]. Adeckel et al. in Kuwait reported an alpha thalassemia incidence of 46% [25], and White et al. reported the incidence rate of 39.8% in Amman [26]. In African regions, the α-thalassemia prevalence was reported to be 68% [27].

The results of the present study showed that the prevalence of anemia among marriage applicants was 0.24 (24% in the total study years: 6% in 2014, 4% in 2015, 5% in 2016, 4% in 2017, and 5% in 2018). In the study by Foruzan Nejad et al. in Ahvaz, the prevalence of anemia was reported to be 5.18% [14].

Based on the results of the present study, the prevalence of sickle cell hemoglobin in marriage applicants was 0.24 (24% for the total study years: 8% in 2014, 1% in 2015, 5% in 2016, 1% in 2017 and 9% in 2018). In the study by

Foruzan Nejad et al. in Ahvaz, the prevalence of sickle hemoglobin was 7.05% [14]. In the study by Canton et al. and in the Kagura region, the respective prevalence was 10% [28]. In the study conducted by Arias et al., the prevalence of sickle cell hemoglobin was 0.49% [29]. In another study by Genninki et al., the total number of patients with sickle cell anemia was 584, and the highest rate was 13 in 100,000 subjects [30].

The present study indicated that the prevalence of hemoglobinopathy in marriage applicants was 0.99 (9% for the total study years: 5% in 2014, 4% in 2015, 0.02% in 2016, 0.05% in 2017, and 0.00% in 2018).

The present study indicated that the prevalence of hemoglobinopathy in marriage applicants was 0.99 (9% for the total study years: 5% in 2014, 4% in 2015, 0.02% in 2016, 0.05% in 2017, and 0.00% in 2018).

Iran is located on the thalassemia belt, and the disease has become widespread due to the history of family marriages. Furthermore, in spite of implementing preventive strategies, preventing and controlling the disease in Iran encounter challenges due to the cultural multiplicity and numerous family marriages. When marriage between two people with thalassemia minor is precluded, these individuals are more likely to marry a person with a sickle cell compared with the previous condition. Accordingly, if the incidence of sickle cell trait is significant in an area where the thalassemia prevention programs are implementing, the problem of major thalassemia prevalence is likely to be replaced by the higher prevalence of hemoglobinopathies, including sickle cell syndrome, particularly sickle-cell-beta thalassemia [13].

### 3.1. Strengths

Large sample size and investigation of marriage applicants' data for several years were the strengths of the study.

**Table 3:** The prevalence of thalassemia, anemia, sickle cell hemoglobin, and hemoglopathy in different years of study (prevalence rate was reported per 1000 people)

Type of blood disease	Category	2014	2015	2016	2017	2018	Total
Thalassemia	In marriage applicants	Beta 0.07	0.067	0.066	0.065	0.065	0.39
	In total population	Alpha 0.14	0.15	0.14	0.14	0.15	0.58
		Beta 0.001	0.0014	0.001	0.0013	0.001	0.005
Anemia	In the marriage applicants	Beta 0.003	0.003	0.002	0.003	0.002	0.013
	In total population	Alpha 0.06	0.04	0.05	0.04	0.04	0.24
Sickle hemoglobin	In the marriage applicants	0.001	0.001	0.008	0.007	0.009	0.026
	In total population	0.08	0.01	0.05	0.01	0.09	0.24
Hemoglopathy	In the marriage applicants	0.001	0.001	0.001	0.001	0.001	0.001
	In total population	0.001	0.001	0.001	0.001	0.001	0.001
		0.05	0.04	0.002	0.005	0.001	0.09

**Table 4:** Determining the difference between the different study years based on thalassemia, anemia, sickle cell hemoglobin, and hemoglobinopathy

Type of blood disease	Category	2014	2015	2016	2017	2018	Pvalue
Thalassemia	Beta	551(7.1)	456(6.8)	397(6.6)	378(6.5)	317(6.5)	< 0.001
	Alpha	1099(14.2)	1029(15.3)	887(14.9)	859(14.8)	761(15.6)	
	normal	6098(78.7)	5247(77.9)	4688(78.5)	4553(78.6)	3794(77.9)	
Anemia	Yes	481(6.2)	333(4.9)	333(5.6)	278(4.8)	255(5.2)	< 0.001
	No	7246(93.8)	6399(95.1)	5639(94.4)	5512(95.2)	4617(94.8)	
Sickle hemoglobin	Yes	66(0.9)	73(1.1)	30(0.5)	8(0.1)	46(0.9)	< 0.001
	No	7748(99.1)	665(98.9)	5942(99.5)	5782(99.9)	4826(99.1)	
Hemoglobinopathy	Yes	44(0.6)	3(0.3)	17(0.3)	32(0.6)	4(0.1)	< 0.001
	No	7704(99.4)	67.9(99.7)	5955(99.7)	5758(99.4)	4868(99.9)	

### 3.2. Weaknesses

Non-generalizability of the study results and the incompleteness of some of the records were among the weaknesses of the present study.

## 4. Conclusion

The present study showed that the incidence of thalassemia and sickle cell anemia in the study years was similar to that in other parts of Iran. However, the implementation and continuation of the beta-thalassemia screening programs in this city can effectively prevent the birth of the newborns with major thalassemia and impose unwanted treatment costs.

## Authors' Contributions

A.R.Z., designed the article. T.R., and M.Sh., did write the article. T.R., was supervisor. M.M., edited the article. All authors read and approved the final manuscript.

## Conflict of Interest

The Authors declare that there is no conflict of interest.

## Acknowledgments

This paper was extracted from the M.Sc. thesis by Mostafa Eslahi. The thesis was approved and financially supported by the research vice-chancellor of Shiraz University of Medical Sciences (IR.SUMS.REC.1396.S794). The professors and officials of the school of health, colleagues, and the participants are also highly appreciated.

## References

- Hojjati MT, Einollahi N, Nabatchian F, Pourfathollah AA, Mahdavi MR. Allelespecific Oligonucleotide Polymerase Chainreaction for the Determination of Rh C/c and Rh E/e antigens in thalassaemic Patients. *Blood Transfus.* 2011; 9(3): 301-5.
- Sotesh H, Akhavan Niaki H, Kowsarian M, Aliasgharian A, Banihashemi A. Frequency of Beta-globin Gene Mutations in Beta-thalassemia Patients from East of Mazandaran. *J Mazand Univ Med Sci.* 2008; 18(67): 17-25.
- Ghotbi N, Tsukatani T. Evaluation of the National Health Policy of thalassaemia screening in the Islamic Republic of Iran. *East Mediterr Health J.* 2005; 11(3): 308-18.
- Olivieri NF. Medical progress: The  $\beta$ -Thalassemias. *N England J Med.* 1999; 341: 99-109.
- Finotti A, Breda L, Lederer CW, Bianchi N, Zuccato C, Kleanthous M, et al. Recent trends in the gene therapy of  $\beta$ -thalassemia. *J Blood Med.* 2015; 19(6): 69-85.
- Rachmilewitz EA, Giardina PJ. How I treat thalassemia. *Blood.* 2011; 118(13): 3479-88.
- Bordbar E, Taghipour M, Zucconi BE. Reliability of Different RBC Indices and Formulas in Discriminating between  $\beta$ -thalassemia Minor and Other Microcytic Hypochromic Cases. *Mediterr J Hematol Infect Dis.* 2015; 7(1): e2015022.
- Rahim F, Abromand M. Spectrum of  $\beta$ -thalassemia mutations in various ethnic regions of Iran. *Pak J Med Sci.* 2008; 24(3): 410-5.
- Miri M, Tabrizi NM, Hadipour DM, Sadeghian VF, Ahmadvand A, Yousefi D, et al. Thalassemia in Iran in Last Twenty Years: the Carrier Rates and the Births Trend. *Iran J Blood Cancer.* 2013; 6(1): 11-8.
- Valizadeh F, Mousavi A, Hashemi-Sotesh MB. Prevalence of Hemoglobinopathies in Premarriage Individuals Referred to Babolsar, Iran (2006-09). *J Gorgan Uni Med Sci.* 2012; 14(1): 106-12.
- Zandian Kh, Keikhaie B, Pedram M, Kianpoor Ghahfarokhi F. Prenatal Diagnosis and Frequency Determination of Alpha and Beta, Thalassemia, S, D, C, and H Hemoglobinopathies; Globin, Mutational Genes, Aanalysis among Voluntary Couples from Ahvaz. *Iran J Blood Cancer.* 2009; 1(3): 95-8.
- Tamaddoni A, Hadavi V, Nejad NH, Khosh Ain A, Siami R, Aghai Meibodi J, et al. Alpha-Thalassemia Mutation Analyses in Mazandaran Province, North Iran. *Hemoglobin.* 2009; 33(2): 115-23.
- Pooladi N, Hosseinpour Feizi MA, Haghi M, Azarfam P, Hosseinpour Feizi AA. Analysis of Beta Thalassemia Mutations Using the Single Strand Conformation Polymorphism (SSCP) Technique. *Sci J Kurdistan Univ Med Sci.* 2010; 15(3): 13-9.
- Nezhad FH, Nezhad KH, Choghakabodi PM, Keikhaei B. Prevalence and Genetic Analysis of  $\alpha$ - and  $\beta$ -Thalassemia and Sickle Cell Anemia in Southwest Iran. *J Epidemiol Glob Health.* 2018; 8(3): 189-95.
- Aziz M, Anwar M. Prevalence of Beta Thalassemia Trait in Quetta City, Cross Section Study. *JUMDC.* 2015; 6(4): 21-6.
- Sharma A, Vageriya V. Prevalence of Thalassemia in Reproductive Age Group Females in Central Gujarat-Literature Review. *Int J Nurs Educ.* 2017; 9(2): 71-4.
- Lai K, Huang G, Su L, He Y. The Prevalence of Thalassemia in Mainland China: Evidence from Epidemiological Surveys. *Sci Rep.* 2017; 7(1): 920.
- Ahmed MM, Salaria SM, Qamar S, Soaz M A, Bukhari M H, Qureshi A H. Incidence of  $\beta$ -Thalassemia Carriers in Muzaffarabad, Azad Kashmir. *Ann Punjab Med College.* 2016; 10(1): 11-9.
- Lau YL, Chan LC, Chan YY, Ha SY, Yeung CY, Waye JS, Chui DH. Prevalence and Genotypes of Alpha-and Beta-Thalassaemia Carriers in Hong Kong- Implications for Population Screening. *N Engl J Med.* 1997; 336: 1298-301.

20. Valizadeh F, Deylami A. Prevalence of Mutations of Alpha Globin Gene in Suspected Alpha Carrier Couples, Babolsar, 2006-2011. *J Mazandaran Univ Med Sci.* 2014; 23(109): 17-25.
21. Zandi Kh, Pedram M. Distribution of Alphathalassemia Mutations in Khuzestan Province: Genetics of the Third Millennium. *Hemoglobin.* 2007; 7(15): 33-42.
22. Neishabury ML, Abbasi Moheb L, Najmabadi H. Alpha-Thalassemia Deletion Analysis in Iran. *Arch Iran Med.* 2001; 4(4): 160-4.
23. El Kalla S, Baysal E. Alpha-Thalassemia in the United Arab Emirates. *Acta Haematol.* 1998; 100(1): 49-53.
24. Borges E, Wenning MR, Kimura EM, Gervásio SA, Sonati MF, Costa FF. High Prevalence of Alpha-Thalassemia among Individuals with Microcytosis and Hypochromia without Anemia. *Braz J Med Biol Res.* 2001; 34(6): 759-62.
25. Adekile AD, Gu LH, Baysal E, Haider MZ, Al Fuzae L, Aboobacker KC, et al. Molecular Characterization of Alphathalassemia Determinants, Beta-Thalassemia Alleles and Beta S Haplotypes among Kuwaiti Arabs. *Acta Haematol.* 1994; 92(4): 176-81.
26. White JM, Byrne M, Richards R, Buchanan TKatsoulis E, Weerasingh K. Red Cell Genetic abnormalities in Peninsular Arabsalphathalassemia. *J Med Genet.* 1986; 23(3): 245-51.
27. Fowkes FJ, Allen SJ, Allen A, Alpers MP, Weatherall DJ, Day KP. Increased Microerythrocyte Count Inhomozygous Alpha(+)- Thalassaemia Contributes Toprotection against Severe Malarial Anaemia. *PLoS Med.* 2008; 5(3): e56.
28. Canatan D. Thalassemias and Hemoglobinopathies in Turkey. *Hemoglobin.* 2014; 38(5): 305-7.
29. El Ariss AB, Younes M, Matar J, Berjaoui Z. Prevalence of Sickle Cell Trait in the Southern Suburb of Beirut, Lebanon. *Mediterr J Hematol Infect Dis.* 2016; 8(1): e2016015.
30. Hemminki K, Li X, Försti A, Sundquist J, Sundquist K. Thalassemia and Sickle Cell Anemia in Swedish Immigrants: Genetic Diseases have become Global. *Sage Open Med.* 2015; 3: 2050312115613097.

**How to cite:** Moghadami M, Sheybani M, Reziaean Zadeh A, Rakhshani T. Investigating the Prevalence of Thalassemia and Sickle Hemoglobin in Marriage Applicants Referring to the Genetic Disease Consulting Center of the Health Center of Kazeroun, 2014-2018. *J Hum Environ Health Promot.* 2019; 5(4): 183-7.