



Original Article

Seroepidemiological study of toxoplasmosis in hemodialysis patients of Ahvaz, Southwest of Iran



Open Access

Forough Kazemi^{1,2}, Somayeh Fallahizadeh^{*1,2}, Mohammad Hossein Feiz-Haddad^{2,3}

ARTICLE INFO

Article History:

Received 24 December 2017

Revised 4 January 2018

Accepted 5 January 2018

Published online 13 January 2018

Keywords:

Seroepidemiology;

Toxoplasma gondii;

Hemodialysis;

Ahvaz;

Iran

¹Student Research Committee, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran ²Department of Parasitology, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran ³Health Research Institute, Infectious and Tropical Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

Correspondence:

Somayeh Fallahizadeh, Department of Parasitology, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.
Email: somayeh.laboratory@gmail.com.

ABSTRACT

Introduction: *Toxoplasma gondii* (*T. gondii*) is an obligate intracellular protozoan that infects most warm-blooded vertebrates. Because of the immunodeficiency in hemodialysis patients, these individuals are at higher risk for reactivating chronic toxoplasmosis; therefore, a diagnosis of early and definitive in these individuals is highly desirable. Thus, the aim of this study is to investigate seroepidemiology of toxoplasmosis in hemodialysis patients of Ahvaz, southwest of Iran during 2015-2016.

Methods: This is a cross-sectional study. A total of 160 participants, 80 hemodialysis patients (test group) and 80 healthy persons (control group) were randomly selected. IgG and IgM against *T. gondii* was measured by CLIA (Chemiluminescence) and ELISA (Enzyme-linked immunosorbent assay) tests. To analyze the data from Chi-square test, and SPSS statistical software of version 21 was used.

Result: In hemodialysis patients, 35 (43.8%) and 25 (31.3%) were positive for the antibody of IgG by ELISA and CLIA, respectively. In healthy individuals, a positive result of 22 (27.5%) and 13 (16.3%) were obtained by ELISA and CLIA, respectively. In hemodialysis patients, 5 (6.3%) and 4 (5%) was positive by ELISA and by CLIA, respectively. In healthy individuals, a positive of 4 (5%) were obtained by ELISA and CLIA. The antibodies of IgG in hemodialysis patients was significantly higher than healthy individuals by ELISA and CLIA ($P<.05$).

Conclusion: The prevalence of *Toxoplasma* in hemodialysis patients was higher than healthy subjects. Our findings show a relatively high prevalence of the parasite among hemodialysis patients and healthy individuals in Ahvaz. So, the health authorities of Ahvaz city must pay more attention to control of the infection, especially in hemodialysis patients.

Introduction

Toxoplasmosis is a protozoan parasitic infection that has a worldwide distribution. *Toxoplasma gondii* (*T. gondii*) causes toxoplasmosis, that is, an obligate intracellular protozoan. *T. gondii* infects most of the warm-blooded vertebrates, for example, humans, other mammals and birds (1). Seroepidemiological studies show that *T. gondii*

antibodies are in serum of one-third of the world adult population (2). Prevalence rates are different in terms of the location and due to differences of culture and health information. Studies show that the toxoplasmosis prevalence rate is 16-40% in North America and Britain, as well as in Central and South America and Europe, it is estimated to be 80-50% (3, 4). Toxoplasmosis occurs mostly in tropical and subtropical areas. *T. gondii* requires 2 hosts to complete its life cycle including the family of cats as the definitive host and vertebrates such as birds and

mammals as an intermediate host. The main routes of transmission include the consumption of water contaminated with oocytes, raw or undercooked meat containing tissue cysts, as well as congenital transmission (5). Individuals of at high risk for toxoplasmosis are immunocompromised persons. Studies have also shown that the infection is related to nervous system disorders such as epilepsy, bipolar disorder and schizophrenia (6). Symptoms in individuals with healthy immune system are mild and transient that include mild fever and swollen lymph glands. In individuals with immune system dysfunction and hemodialysis, latent form of the parasite can be reactivated causing different clinical complications from swelling of lymph, damage to the central nervous system, epilepsy, cerebral calcifications, pneumonia and myocarditis (7). Because of the immunodeficiency in hemodialysis patients, these individuals may be at higher risk for reactivating chronic toxoplasmosis. Therefore, an early and definite diagnosis of and definitive in these individuals is highly advisable (8). Since hemodialysis patients are at higher risk for infection of *T. gondii*; thus, the aim of this study is to investigate seroepidemiological aspect of toxoplasmosis in hemodialysis patients of Ahvaz, southwest of Iran in 2015-2016.

Methods

Study area

Ahvaz is a city in the center of Khuzestan Province in Southwest of Iran. The city is 375 kilometers square and its population has been reported as 1,425,891 until 2006. The city has a desert like climate with temperatures above 50° C that is one of the hot cities in the world. The average annual rainfall is about 230 mm. According to the recent report by the World Health Organization (WHO), Ahvaz has the most polluted climate in the world.

Study population

This is a cross-sectional study. A total of 160 participants, 80 hemodialysis patients (test group) and 80 healthy persons (control group) referred to hospitals of Jundishapure University of Medical Sciences (southwest of Iran) were selected randomly. Informed consent was obtained from all participants included in the study (9).

Sampling collection and Serological diagnosis

Five ml blood samples taken from hemodialysis patients as well as healthy subjects were centrifuged. Then, 2 ml of isolated serum was placed in micro-

tube and was maintained until the time of testing at 20 °C. To test the samples, they were taken out of the freezer and were melted. Anti-Toxoplasma IgG and IgM antibodies were measured by CLIA (Chemiluminescence) and ELISA (Enzyme-linked immunosorbent assay) tests (10). For Measurement of IgG and IgM by chemiluminescence Architect Kit was used. In ELISA method, IgM and IgG were measured by Ab Capture ELISA and Indirect ELISA methods, respectively.

Ethical consideration

Before performing the research, it was explained to the participants. An informed consent was obtained from all participants included in the study.

Statistical analysis

To analyze the data from Chi-square tests and SPSS, statistical software of version 21 was used. The P-value less than 0.05 was considered significant.

Results

Tables 1 and 2 show IgG antibody of *T. gondii* by ELISA and CLIA in hemodialysis patients and healthy subjects, respectively: For the antibody of IgG, in hemodialysis patients were positive, that is, 35 (43.8%) and 25 (31.3%) by ELISA and CLIA respectively. In healthy individuals a positive result of 22 (27.5%) and 13 (16.3%) by ELISA and CLIA, was obtained respectively. The IgG antibody in hemodialysis patients was significantly higher than healthy subjects in both methods ELISA and CLIA ($P < 0.05$).

Table 1: Evaluation of *T. gondii* IgG antibody in hemodialysis patients and healthy individuals by ELISA

		Hemodialysi	Control	Total	
		s			
IgG	Positive	Count	35	22	57
		%	43.8%	27.5%	35.6%
	Negative	Count	45	58	103
		%	56.3%	72.5%	64.4%
Total		Count	80	80	160

Also, tables 3 and 4 show IgM antibody of *T. gondii* by ELISA and CLIA in hemodialysis patients and healthy subjects, respectively. Antibody of IgM, in hemodialysis patients showed positive data of 5 (6.3%) by ELISA and 4 (5%) by CLIA and in healthy individuals a positive of 4 (5%) was obtained by

Table 2: Evaluation of *T. gondii* IgG antibody in hemodialysis patients and healthy individuals by CLIA

		Hemodialysis		Control	Total
		Count	25	13	38
IgG	Positive	%	31.3%	16.3%	23.8%
		Count	55	67	122
	Negative	%	68.8%	83.8%	76.3%
		Total	Count	80	80

methods of ELISA and CLIA, respectively. There was no significant difference in the IgM antibody between hemodialysis patients and healthy subjects by both methods of ELISA and CLIA ($P > 0.05$) individuals by ELISA and CLIA ($P < 0.05$). The results show that the prevalence of *T. gondii* in hemodialysis

Discussion

T. gondii is an obligate intracellular protozoan from apicomplexa branches and order of coccidia (11). Pregnant women and individuals with immune system disorders are of high priority regarding *T. gondii* (12). Because of the immunodeficiency in hemodialysis patients, these individuals are at higher risk of reactivating toxoplasmosis infection. Therefore, an early definite diagnosis in these individuals is highly advisable (8). Since hemodialysis patients are at higher risk for infection of *T. gondii*; thus, the aim of this study was to investigate seroepidemiological of toxoplasmosis in hemodialysis patients of Ahvaz, Southwest of Iran in 2015-2016. On the other hand, toxoplasmosis occurs mostly in tropical and subtropical regions (5) and due to high heat in the city, the prevalence study of this parasite seems essential in hemodialysis patients in Ahvaz, southwest of Iran. Our findings show that the tests for the antibody of IgG, in hemodialysis patients were positive, that is, 35 (43.8%) and 25 (31.3%) by ELISA and CLIA respectively. In healthy individuals a positive result of 22 (27.5%) and 13 (16.3%) by ELISA and CLIA, was obtained respectively. Also antibody of IgM, in hemodialysis patients showed positive data of 5 (6.3%) by ELISA and 4 (5%) by CLIA and in healthy individuals a positive of 4 (5%) was obtained by methods of ELISA and CLIA, respectively. The antibody of IgG in hemodialysis patients was significantly higher than healthy patients is higher than healthy individuals; these results are consistent with results of previous studies. Similarly, in 2013, Saki et al. in Ahvaz measured IgG and IgM antibodies in hemodialysis patients and healthy individuals by ELISA and PCR tests and showed that the prevalence of this parasite in healthy subjects was 26%, but in hemodialysis patients it was found to be 29.3%.

Table 3: Evaluation of *T. gondii* IgM antibody in hemodialysis patients and healthy individuals by ELISA

		Hemodialysis	Control	Total	
IgM	Positive	Count	5	4	9
		%	6.3%	5.0%	5.6%
	Negative	Count	75	76	151
		%	93.8%	95.0%	94.4%
Total		Count	80	80	160

The results indicate a significant difference between two groups and the prevalence of this parasite was more in hemodialysis patients (8). Also, Rezavand et al. in 2016 studied the prevalence of *T. gondii* in Iran by ELISA and PCR tests and showed that of 90 patients under hemodialysis, 54 patients (60.0%) for anti-*T. gondii* IgG antibody and 3 patients (3.3%) for IgM were positive. Of 90 of healthy subjects, 34 cases (37.8%) were positive for IgG antibody and all healthy subjects were negative for IgM (13). In 2014-2015, Rasti et al. in Kashan and Qom (two cities in center of Iran) showed that the prevalence of *T. gondii* was significant difference between the two groups of control ($n = 120$) and hemodialysis ($n = 135$) and this parasite was greater in hemodialysis patients. So that was a positive for 63% hemodialysis patients and 33.3% of healthy individuals (14). Similarly, in 2013, Ebrahimzadeh et al. in Iran showed that for IgG antibody, 21 of 37 samples of hemodialysis patients was positive but only, 11 of 37 samples of healthy individuals was positive (15). Maraghi et al. in 2011 in Abadan and Khoramshahr (two cities in southwest of Iran) showed that this parasite was greater in hemodialysis patients. Since, for IgG antibody, 61 of 150 samples of hemodialysis patients were positive but only 39 of 150 samples of healthy individuals was reported positive. Also, for IgM antibody 13 cases of hemodialysis patients was positive but positive cases were not observed in the control group (16). Also in our findings, there was no significant difference in the IgM antibody between hemodialysis patients and healthy subjects

Table 4: Evaluation of *T. gondii* IgM antibody in hemodialysis patients and healthy individuals by CLIA

		Hemodialysis		Control	Total
IgM	Positive	Count	4	4	8
		%	5.0%	5.0%	5.0%
	Negative	Count	76	76	152
		%	95.0%	95.0%	95.0%
	Total	Count	80	80	160

by both methods of ELISA and CLIA ($P>0.05$) this can be due to the availability of low sample size. These results have also been obtained in other countries. For example, in 2005, Ocak et al. in Turkey showed that there were positive 195 of 255 hemodialysis patients for *T. gondii*, but in control group, from 50 healthy samples, 24 cases were positive. The prevalence of this parasite was greater in hemodialysis patients significantly. Also, the results showed that by increasing treatment duration of hemodialysis patients, the prevalence of this parasite was enhanced (17). The results of all these studies indicate that the prevalence of *T. gondii* in patients undergoing hemodialysis is higher than healthy subjects. The patients due to weakened immunity may activate latent form of toxoplasmosis, which is followed by a lot of complications caused by infection with *T. gondii*. Thus, the results of this research can be a help in planning to control and prevent the parasite in hemodialysis patients. One of the strengths of this study is using two powerful diagnostic tests for IgM and IgG antibodies but the limitation of this study is the low sample size. So, it is suggested to the succeeding researchers to go through the tests using samples with high volumes.

Conclusion

The prevalence of *Toxoplasma* in hemodialysis patients has been higher than healthy subjects. Because of immunodeficiency in hemodialysis patients, latent form of the parasite can be reactivated causing different clinical complications such as swelling of lymph nodes, damage to the central nervous system (e.g., epilepsy or cerebral calcification), pneumonia and myocarditis. Our findings show a relatively high prevalence of the parasite among hemodialysis patients and healthy individuals in Ahvaz. So, the health authorities of Ahvaz city must pay more attention to control of the infection, especially in hemodialysis patients.

Ethical disclosure

Before performing the research, it was explained to the participants. An informed consent was obtained from all participants included in the study.

Acknowledgements

We would like to acknowledge Ahvaz Jundishapur University of Medical Sciences and the study participants for their assistances.

Author Contributions

All the authors have accepted responsibility for the entire content of this submitted manuscript and approved submission.

Conflict of interest

The authors declare that they have no conflict of interest.

Funding/Support

None declared.

References

- Gharavi M, Rahnama N, Jahani M. Seroepidemiological survey of Toxoplasma infections of mentally retarded children. Iran J Public Health. 2005;34(1):19-22.
- Gharavi M, Jalali S, Khademvatan S, Heydari S. Detection of IgM and IgG anti-Toxoplasma antibodies in renal transplant recipients using ELFA, ELISA and ISAGA methods: comparison of pre-and post-transplantation status. Ann Trop Med Parasito. 2011;105(5):367-71. DOI: 10.1179/1364859411Y.0000000022.
- Jeannel D, Niel G, Costagliola D, Danis M, Traore BM, Gentililini M. Epidemiology of toxoplasmosis among pregnant women in the Paris area. Int J Epidemiol. 1988;17(3):595-602. PMID:3264821.
- Jones JL, Ogunmodede F, Scheftel J, Kirkland E, Lopez A, Schulkin J, et al. Toxoplasmosis-related knowledge and practices among pregnant women in the United States. Infect Dis Obstet Gynecol. 2003;11(3):139-45. PMID:15022874.
- Dubey JP. The history of Toxoplasma gondii—the first 100 years. J Eukaryot Microbiol. 2008;55(6):467-75. DOI:10.1111/j.1550-7408.2008.00345.x

6. Ahmadpour E, Daryani A, Sharif M, Sarvi S, Aarabi M, Mizani A, et al. Toxoplasmosis in immunocompromised patients in Iran: a systematic review and meta-analysis. *J Infect Dev Ctries*. 2014;8(12):1503-10.
7. Soltani S, Khademvatan S, Saki J, Shahbazian H. Detection of Toxoplasmosis in Renal Transplant Recipients by ELISA and PCR Methods in Ahvaz, South-West of Iran. *Jundishapur J Microbiol*. 2013;6(9). PMID:25500647.
8. Saki J, Khademvatan S, Soltani S, Shahbazian H. Detection of toxoplasmosis in patients with end-stage renal disease by enzyme-linked immunosorbent assay and polymerase chain reaction methods. *Parasitol Res*. 2013;112(1):163-8. DOI: 10.1007/s00436-012-3120-6.
9. Nasiri V, Esmailnia K, Karim G, Nasir M, Akhavan O. Intestinal parasitic infections among inhabitants of Karaj City, Tehran province, Iran in 2006-2008. *Korean J Parasitol*. 2009;47(3):265. DOI: 0.3347/kjp.2009.47.3.265.
10. Firouz ZE, Kaboosi H, Nasiri AF, Tabatabaie SS, Golhasani-Keshtan F, Zaboli F. A comparative serological study of toxoplasmosis in pregnant women by CLIA and ELISA methods in Chalus City Iran. *Iran Red Crescent Med J*. 2014;16(4). DOI: 10.5812/ircmj.15115.
11. Mousavi SSB, Faramarzi M. Do we Need to Screen Uremic Patients for Toxoplasmosis before Kidney Transplantation? *Shiraz E-Med J*. 2013;14(4).
12. Nissapatorn V, Leong TH, Lee R, Ibrahim J, Yen TS. Seroepidemiology of toxoplasmosis in renal patients. *Southeast Asian J Trop Med Public Health*. 2011;42(2):237. PMID:21710842.
13. Rezavand B, Poornaki AM, Mokhtari KR, Mohammad A, Andalibian A, Abdi J. Identification and determination of the prevalence of *Toxoplasma gondii* in patients with chronic renal failure by ELISA and PCR. *Asian Pac J Trop Dis*. 2016;6(5):347-9. DOI: 10.1016/S2222-1808(15)61044-1
14. Rasti S, Hassanzadeh M, Soliemani A, Hooshyar H, Mousavi SGA, Nikoueinejad H, et al. Serological and molecular survey of toxoplasmosis in renal transplant recipients and hemodialysis patients in Kashan and Qom regions, central Iran. *Ren Fail*. 2016;1-4. DOI: 10.3109/0886022X.2016.1172940.
15. Zadeh AE, Bamedi T, Etemadi S, Shahrakipour M, Saryazdipour K. Toxoplasmosis as a complication of transfusion in hemodialysis patients. *Iran J Ped Hematol Oncol*. 2014;4(1):22. PMID:24734160.
16. Maraghi S, Yadyad MJ, Sheikhi M, Shamakhteh F, Latifi SM. Study the anti-Toxoplasma antibodies (IgG and IgM) in hemodialysis patients of Abadan and Khoramshahr cities Southwest Iran in 2011 using ELISA. *Jundishapur J Microbiol*. 2013;6(7). DOI:10.5812/jjm.7113.
17. Ocak S, Duran N, Eskiocak AF, Aytac H. Anti-Toxoplasma gondii antibodies in hemodialysis patients receiving long-term hemodialysis therapy in Turkey. *Saudi Med J*. 2005;26(9):1378-82. PMID:16155651.