

Review Paper

A Scoping Review of Questionnaires on Stigma of Mental Illness in Persian



Farahnaz Rezvanifar¹, Seyed Vahid Shariat^{2,3}, Homayoun Amini^{4,5}, Maryam Rasoulia¹, *Mohammadreza Shalbafan¹

1. Mental Health Research Center, Iran University of Medical Sciences, Tehran, Iran.
2. Mental Health Research Center, Psychosocial Health Research Institute (PHRI), Iran University of Medical Sciences, Tehran, Iran.
3. School of Behavioral Sciences and Mental Health (Tehran Institute of Psychiatry), Iran University of Medical Sciences, Tehran, Iran.
4. Department of Psychiatry, Roozbeh Psychiatry Hospital, Tehran University of Medical Sciences, Tehran, Iran.
5. Psychosomatic Medicine Research Center, Tehran University of Medical Sciences, Tehran, Iran.



Citation Rezvanifar F, Shariat S, Amini H, Rasoulia M, Shalbafan M. [A Scoping Review of Questionnaires on Stigma of Mental Illness in Persian (Persian)]. Iranian Journal of Psychiatry and Clinical Psychology. 2020; 26(2):240-259. <http://dx.doi.org/10.32598/ijpcp.26.2.2619.1>

doi <http://dx.doi.org/10.32598/ijpcp.26.2.2619.1>



Received: 09 Sep 2019

Accepted: 12 Feb 2020

Available Online: 01 Jul 2020

Key words:

Social stigma, Mental disorders, Questionnaires and surveys

ABSTRACT

Objectives In order to plan according to the needs of psychiatric patients, there is a need to study different aspects of social stigma in target groups. Due to the limited access to questionnaires used for assessment of social stigma in Persian, this study aims to review all of the tools available for assessment of social stigma, which have been designed in Iran or have been translated to Persian.

Methods This is a conceptual review study. After extensive review of resources, articles related to social stigma in the health system and conducted on health care staff, students, caregivers or patients' families, psychiatric patients, mothers with children with autism, mothers with children with Down syndrome, and the general population were included in the study. We extracted the information of Persian questionnaires used in the reviewed articles. The questionnaires and additional information were collected by contacting the authors of the articles via E-mail.

Results There were 21 questionnaires for assessing social stigma in Persian language. Of these, 19 questionnaires could be accessed and analyzed. Most of the questionnaires were the translated version of other tools, and only 4 questionnaires had been developed and validated in Persian.

Conclusion Most of the questionnaires in Persian are used to measure social stigma in health care staff and students followed by caregivers or patients' families. There is limited number of questionnaires for the assessment of social stigma in general population and patients. This is in contrast to the other languages that have mainly focused on assessment of stigma in general population and patients. To date, no questionnaire in Persian exists for the assessment of social stigma in children and adolescents.

Extended Abstract

1. Introduction

P

atients with mental illness generally face two major issues related to their disorder: they should deal with not only different signs and

symptoms of their disorders (including anxiety, delusion, hallucination, etc.) but also with the concurrent social stigma of mental illness. The families of the patients with mental illness usually feel embarrassed and guilty about the disorder of their relative and would also experience similar social isolation and stigma.

* Corresponding Author:

Mohammadreza Shalbafan, PhD.

Address: Mental Health Research Center, Iran University of Medical Sciences, Tehran Iran.

Tel: +98 (21) 66551665

E-mail: shalbafan.mr@iums.ac.ir

Despite the progress in the treatment of mental disorders in recent decades, the social stigma of mental illness still exists and affects the patients with mental illness and their families as well as mental health personnel. For appropriate planning for the needs of the patients with mental illness, it is necessary to study different aspects of stigma in various sub-populations.

To adequately address the need for precise assessment of the social stigma of mental illness, we need valid and reliable assessment tools. Due to the limited access to questionnaires to assess the social stigma of mental illness in Persian, we planned to review all of the assessment tools related to the social stigma of mental illness, which have been designed in Iran or translated into Persian.

2. Methods

We searched for both Persian and English sources. We searched the English databases, including PubMed, Google Scholar, Scopus, and PsycINFO, along with the Persian databases, including IranMedex, Scientific Information Database, Irandoc, and Magiran. We did not limit our searches for a specific period. For English databases, we combined three keywords of "Stigma" OR its related words, "Mental Health" OR its related words, and "Iran" OR the related words using the AND operator. For Persian databases we used only the Persian equivalent terms for "stigma" and no Boolean search was used.

Our search yielded 289 articles and 248 theses. Only 25 articles and 7 theses met our inclusion criteria of studying stigma, using a questionnaire in English and Persian. After removing shared questionnaires, we found 21 distinct stigma questionnaires. Then, we emailed the corresponding authors of all of the articles, and informing them about our study asked them to send us a copy of the questionnaire and surveys that they used in their study. We also asked their permission to publish the full text of their questionnaire as an appendix to our article. Finally, we gathered the whole data on 19 questionnaires.

3. Results

Twenty-one questionnaires and surveys were used to assess 'stigma' in the Persian language that included: Social Distance Item, Perceived Dangerousness of Mental Patients Item, The Attribution Questionnaire, The Level of Familiarity Scale, Attitudes toward Seeking Professional Psychological Help, Knowledge and Attitude about Patients with Mental Disorders and Epilepsy, Experience of Caregiving Inventory, Internalization of Stigma for Mothers of Children with Disability, Opening Minds Scale for Health Care

Providers, Perceived Stigma of Mental Illness in Nurses, Opinion about Mental Illness, Depression Stigma Scale, Stigma Scale, The Parents Internalized Stigma of Mental Illness Scale, Family Caregiver's Stigma toward Patients with Mental Illness, Opening Minds Scale for Health Care Providers, Stigma in Patients with Mental Disorders, Perceived Stigma Questionnaire, Cognitive Appraisal of Stigma-related Stress, Rejection Sensitivity Questionnaire and Perceived Stigma Scale-Revised. We could not find the Persian version of the Rejection Sensitivity Questionnaire and Perceived Stigma Scale.

4. Discussion

Most of the questionnaires and surveys of the social stigma of mental illness in Persian were translated version from other languages except four questionnaires that have been originally developed and validated in Persian. These four questionnaires include the Internalization of Stigma for Mothers of Children with Disability, Family Caregiver's Stigma toward Patients with Mental Illness, Perceived Stigma Questionnaire, and Perceived Stigma of Mental Illness in Nurses. Only two questionnaires assessed the stigma of mental illness within the general population (Knowledge and Attitude about Patients with Mental Disorders and Epilepsy, Opinion about Mental Illness). However, eight questionnaires are available to assess the stigma of mental illness in caregivers or family members of patients with mental disorders.

There are ten questionnaires for assessment of stigma of mental illness in health care providers, and medical and nursing students as future care providers. Few questionnaires and surveys are available in Persian for assessment of stigma in patients themselves. In the world literature, most of the questionnaires have been developed to assess stigma in the general population and then in patient. However, most of the questionnaires and surveys studied in Iran are related to care providers and students and then related to caregivers or family members of the patients. Limited questionnaires are available for assessment of stigma in general population and patients. Furthermore, we did not find any questionnaires for assessment of stigma in children and adolescents. Most of the questionnaires and surveys that are being used in Persian are translations from an English questionnaire. Interestingly, some of the questionnaires are used for a different target population than the original questionnaire without clear justifications.

Ethical Considerations

Compliance with ethical guidelines

For all reviewed questionnaires, necessary permissions were obtained from their authors. Because of the type of article (Review), there was no need for obtaining ethical approval.

Funding

This research did not receive any grant from funding agencies in the public, commercial, or non-profit sectors.

Authors contributions

Conceptulization and design: Farahnaz Rezvanifar, Seyed Vahid Shariat, and Mohammadreza Shalbafan; Data collection: Farahnaz Rezvanifar, Mohammadreza Shalbafan, and Maryam Rasoulia; Initial draft preparation: Farahnaz Rezvanifar and Homayoun Amini; Editing & review: All authors.

Conflicts of interest

The authors declared no conflict of interest.