

Computational Analysis of High Risk Missense Variant in Human UTY Gene: A Candidate Gene of AZFa Sub-region

Mili Nailwal, Jenabhai Bhathibhai Chauhan *

- P.G. Department of Genetics, Ashok & Rita Patel Institute of Integrated Study & Research in Biotechnology and Allied Sciences (ARIBAS), Gujarat, India

Abstract

Background: The human Ubiquitously transcribed tetratricopeptide repeat gene, Y-linked (UTY) gene encodes histone demethylase involved in protein-protein interactions. UTY protein evidence at protein level predicted intracellular and secreted protein. UTY is also involved in spermatogenesis process.

Methods: The high-risk non-synonymous single nucleotide polymorphism in the coding region of the UTY gene was screened by SNP database and identified missense variants were subjected to computational analysis to understand the effect on protein function, stability and structure by SIFT, PolyPhen 2, PANTHER, PROVEAN, I-Mutant 2, iPTREE-STAB, ConSurf, ModPred, SPARKS-X, QMEAN, PROCHECK, project HOPE and STRING.

Results: A total of 151 nsSNPs variants were retrieved in UTY gene out of which one missense variant (E18D) was predicted to be damaging or deleterious using SIFT, PolyPhen 2, PANTHER and PROVEAN. Additionally, E18D variant showed less stability, high conservation and having role in post translation modification using i-Mutant 2 and iPTREE-STAB, ConSurf and ModPred, respectively. The predicted 3D model of UTY using SPARKS-X with z-score of 15.16 was generated and validated via QMEAN (Z-score of 0.472) and PROCHECK which plots Ramachandran plot (85.3% residues in most favored regions, 12.3% in additionally allowed regions, 2.0% in generously allowed regions and 4.0% were in disallowed regions) and it indicates a good quality model. STRING showed that UTY interacts with ten different proteins.

Conclusion: This study revealed that SNP data available on database was deduced to find out the most damaging nsSNPs *i.e.* rs3212293 (E18D). Therefore, it provides useful information about functional SNPs for future prospects concerning infertility in men.

Keywords: AZFa, Computational analysis, Male infertility, Missense, nsSNP, UTY gene.

To cite this article: Nailwal M, Chauhan JB. Computational Analysis of High Risk Missense Variant in Human UTY Gene: A Candidate Gene of AZFa Sub-region. J Reprod Infertil. 2017;18(3):298-306.

* Corresponding Author:
Jenabhai B. Chauhan,
Department of Genetics,
Ashok & Rita Patel
Institute of Integrated
Study & Research in
Biotechnology and Allied
Sciences (ARIBAS), New
Vallabh Vidyanagar-
388121, Dist-Anand,
Gujarat, India
E-mail:
jenabhaichauhan@aribas.
edu.in,
jbc109@gmail.com,
chauhanjena@gmail.com

Received: Feb. 2, 2017

Accepted: Apr. 16, 2017

Introduction

Infertility is a major public health problem in which approximately 15% of couples are not able to conceive an offspring (1) and 30% to 50% of infertile cases are related to only male factors (2). Genetic factor is considered as one of the most important etiology of infertility in men (3). Azoospermia factor (AZF) region on the Y chromosome deletion leads to severe spermatogenic

failure like phenotypic or sperm abnormality (4). AZF region in the long arm of Y chromosome (Yq11) comprises three main sub-regions namely AZFa, AZFb and AZFc (5) where essential genes are located for spermatogenesis (6). Among all the three sub-regions, AZFa is the smallest with molecular length of approximately 800 Kb (7). AZFa sub-region contains three main candidate

genes *i.e.* Ubiquitin-specific protease 9 (USP9Y), Dead box on the Y (DBY) and ubiquitously transcribed tetratricopeptide repeat gene, Y-linked (UTY).

It has been reported that human UTY gene is located to band 5C corresponding to AZFa sub-region and this band contains one or more genes that have function in spermatogenesis process and Y-specific growth gene (8). The corresponding UTY gene is located on chromosome Yq11.221 and is also known as Histone demethylase UTY, KDM6AL, KDM6C and UTY1. UTY gene has a total of 20 exons (9) and encodes for enzyme which is rich in tetratricopeptide repeats that may be included in protein-protein interaction (8, 10). The UTY gene is conserved between human and mouse (11). UTY gene is ubiquitously expressed and present in single copy (12).

Single nucleotide polymorphism (SNP) is a variation of single nucleotide which can be found anywhere throughout the genome (13). Mostly SNPs are neutral but some SNPs could predispose human to disease or affect drug response (14). Non-synonymous SNPs are found in coding region which can influence resulting protein structure and/or function with either neutral or deleterious effects (15, 16).

It is believed that autosomal and sex chromosomal abnormality accounts for 10-15% of cases in men with infertility (17). Furthermore, recently it has been suggested that genetic causes of infertility in men comprise Y chromosome microdeletion, genetic mutation and single nucleotide polymorphism as well as chromosomal aberration (18, 19).

Therefore, detection of molecular changes in the Yq11 region of infertile males is required. Deleterious SNPs for the UTY gene have not been converted to data through computational analysis. Thus, to inspect the potential association among the genetic mutation and phenotypic variation, different algorithms are used to compute the high risk missense mutations in coding regions which may have impact on protein structure and/or function of UTY. Considering the role played by UTY gene in male infertility, the study aimed to narrow down the candidate non-synonymous SNPs (nsSNPs) for the UTY through computational analysis which may influence the protein structure and/or function that may serve an important role in male infertility.

Methods

Genomic analysis of UTY was conducted using sorting intolerant from tolerant (SIFT), PolyPhen 2, protein variation effect analyzer (PROVEAN) and protein analysis through evolutionary relationships (PANTHER) to retrieve deleterious mutation followed by stability analysis using I-Mutant 2 and iPTREE-STAB. Additionally, conservation analysis prediction (ConSurf), 3D structure prediction, validation of model, structural effect and protein-protein interaction were done for UTY. Details of each tool are described below and highlighted in table 1.

Dataset: UTY SNPs (rsIDs) related data and protein sequence were extracted from the National Center for Biotechnology Information (NCBI- <http://www.ncbi.nlm.nih.gov>) and related information

Table 1. Computational approaches available as online tools

Server	Input	URL	Reference
SIFT	rs id's	http://sift.jcvi.org/	20
PolyPhen 2.0	Protein sequence in FASTA format, Position and Substitution	http://genetics.bwh.harvard.edu/pph2	21
PANTHER	Protein sequence, substitution and single organism	http://pantherdb.org/tools/csnpscoreForm.jsp	22
PROVEAN	Protein sequence and amino acid variations	http://provean.jcvi.org/index.php	23
I-Mutant 2.0	Protein Sequence, position and new residue	http://folding.biofold.org/cgi-bin/i-mutant2.0.cgi	24
iPTREE-STAB	Residue of deleted, introduced and neighbors mutation type	http://210.60.98.19/IPTREEr/iptree.htm	25
Consurf	Protein sequence	http://consurf.tau.ac.il/	26
ModPred	Protein sequence	http://montana.informatics.indiana.edu/ModPred/index.html	27
SPARKS-X	Protein Sequence	http://sparks-lab.org/yueyang/server/SPARKS-X/	28
HOPE Project	Protein Sequence, select a Residue to Mutate and select Mutation	http://www.cmbi.ru.nl/hope/home	29
QMEAN	PDB file model	https://swissmodel.expasy.org/qmean/cgi/index.cgi	30
PROCHECK	PDB file model	http://www.ebi.ac.uk/thornton-srv/software/PROCHECK/	31
STRING	Protein name and organism	http://string-db.org	32

regarding UTY gene and protein was collected from Online Mendelian Inheritance in Man (OMIM-<https://www.omim.org/entry/400009>).

Prediction of functional connection of missense mutation: Four tools were used namely SIFT, PolyPhen 2, PANTHER and PROVEAN to predict the functional context of missense mutation.

SIFT (Sorting intolerant from tolerant) tool predicts the single amino acid change impact on protein function that would be either tolerated or damaging. The SIFT score will be below or equal to 0.05 when amino acid substitution is predicted damaging and if the score is above 0.05, amino acid substitution is predicted to be tolerating (20). PolyPhen 2 (Polymorphism phenotyping version 2) predicted whether the amino acid substitution is probably damaging, possibly damaging or benign with a score ranging from 0.0 (benign) to 1.0 (damaging).

PANTHER cSNP (Protein analysis through evolutionary relationship- coding SNP) predicts functional effect of amino acid substitution on protein by calculating subPSEC (Substitution position-specific evolutionary conservation) score on the basis of alignment of evolutionary related proteins (33).

PROVEAN (Protein variation effect analyzer) predicts the effect of any type of protein sequence variation consisting of amino acid substitution and in-frame insertion and deletion change on proteins biological function. It analyzes the nsSNPs as deleterious or neutral when the score is below or above the threshold, respectively (23).

Prediction of stability change upon missense mutation: Generally, mutation leads to structural stability changes which affect the function of protein. Therefore, the stability check was carried out using I-Mutant 2 and iPTREE-STAB. I-Mutant 2 server prediction is based on support vector machine (SVM) for protein stability change of the folded protein in relation with single point mutation. It predicts direction ($\Delta\Delta G$ sign) as well as $\Delta\Delta G$ associated values of the protein stability changes upon single site mutations. iPTREE-STAB server is based on decision tree where it predicts the stability change ($\Delta\Delta G$ values) due to single site mutation. Thus, it predicts whether the outcome of the amino acid substitution is stabilizing or destabilizing (34).

Phylogenetic conservational analysis of UTY: Conservation prediction of UTY amino acids was analyzed using ConSurf tool. It uses high-throughput

characterization of functional regions of the protein. The degree of conservation of amino acid is calculated based on conservational score in the scale of 1-9 where 1-3 scores are variable, 4-6 scores indicate average conservation and 7-9 scores indicate high conservation (26).

Prediction of post translational modification sites for UTY: In UTY protein, post translational modification (PTM) sites were sequence based, predicted using ModPred. It accommodated 34 ensembles of logistic regression models that were trained independently on a united set of 126,036 non-redundant experimentally verified sites for 23 distinct polymorphism, retrieved from public databases and an ad-hoc literature investigation (27).

Prediction of 3D structure of UTY: SPARKS-X is a single method fold recognition technique which gives 3D structure. This tool was improved by changing the alignment scoring function as well as adding the SPINE-X techniques which upgraded prediction of secondary structure, backbone torsion angle and solvent accessible surface area (28).

Model validation for UTY: The predicted model of UTY from SPARKS-X was validated using qualitative model energy analysis (QMEAN) and PROCHECK. QMEAN z-score was another tool used for quality assessment where it analyzed the degree of nativeness of the predicted 3D structure of protein. The QMEAN score imitated the predicted global model reliability ranging between 0-1 (35). PROCHECK provides a detailed analysis on the stereochemistry quality of the 3D protein structure. Ramachandran plot with Phi/Psi was provided by the PROCHECK to validate the backbone structure of protein.

Prediction of structural effect of point mutation on UTY: Project HOPE was used to know the structural effect where it was used for molecular dynamics simulation to analyze the single point mutation. First BLAST was performed against PDB and built a homology model and retrieved tertiary structure information through WHATIF followed by the access to UniProt database features. Furthermore, the protein features were predicted using Distributed Annotation System (DAS) server (29, 36).

Prediction of protein-protein interactions: Search Tool for the Retrieval of Interacting Genes/Proteins (STRING) database provides a critical assessment and integration of protein-protein interactions whether it is direct (physical) or indirect (functional) association (37).

Table 2. nsSNP analysis by SIFT

Sr. No.	SNP	Amino acid change	Amino acid	Using homologues in the protein alignment	
				Prediction	Score
1	rs9341273	E34G	E	TOLERATED	1
			G	TOLERATED	0.19
2	rs3212293	E18D	E	TOLERATED	1
			D	DAMAGING	0
3	rs9341281	R1015Q	R	TOLERATED	1
			Q	DAMAGING	0
4	rs75596360	T261P	T	TOLERATED	1
			P	DAMAGING	0
5	rs112993031	S824F	S	TOLERATED	0.33
			F	DAMAGING	0

Table 3. nsSNP analysis by PolyPhen 2

dbSNP ID	Amino acid substitution	HumDiv		HumVar	
		Prediction	Score	Prediction	Score
rs61730117	E34G	Benign	0.002	Benign	0.002
rs3212293	E18D	Probably Damaging	0.979	Probably Damaging	0.983
rs9341281	R1015Q	Benign	0.006	Benign	0.000
rs75596360	T261P	Benign	0.026	Benign	0.027
rs112993031	S824F	Benign	0.190	Benign	0.042

Results

SNP datasets: The polymorphism data for UTY gene investigated in present work was retrieved from NCBI dbSNP¹ database and contained 151 missense, 4 non-sense, 2 splice site at 3' end, 73 UTR at 3' end, 1 splice site at 5' end, 5 UTR at 5' end, 84 coding synonymous, 6 frame shift, 2302 introns and 4 stop gained SNPs. Only non-synonymous coding SNPs of UTY were selected for this investigation.

Prediction of functional nsSNPs in UTY: The UTY single nucleotide variants were administrated to computational analysis through variety of tools.

According to SIFT, a total of 5 out of 151 nsSNPs of UTY gene from dbSNP analysis were predicted to be tolerated or deleterious (Table 2). SIFT classified 4 nsSNPs damaging (E18D, R1015Q, T261P and S824F) where tolerance index score was identified to be 0 (<0.05) and 1 nsSNP (E34G) tolerated with tolerance index score of 0.19. Only 1 missense variant was predicted to be probably

damaging (E18D) and the rest 4 missense variants were classified to be benign through PolyPhen 2 output (Table 3).

To increase the accuracy of computational techniques to compute utmost deleterious SNPs, different computational methods were used. Therefore, PANTHER and PROVEAN were used for further analysis. Out of 5 variants, one (E18D) was predicted to be damaging with SIFT and PolyPhen 2. PANTHER was performed to verify validation of results obtained from two tools and E18D was portrayed damaging with SIFT, PolyPhen 2 and now with PANTHER too (Table 4). E18D was selected for further confirmation through PROVEAN where the output came to be deleterious (Table 4).

Protein stability of UTY variant upon point mutation was found using I-Mutant 2.0 and iPTREE-STAB. E18D was subjected for stability prediction and showed decreased stability by both the tools (Table 5).

Table 4. nsSNP analysis by PANTHER and PROVEAN

Variant	PROVEAN		PANTHER
	PROVEAN Score	Prediction (cutoff= -2.5)	
E18D	-2.536	Deleterious	Possibly damaging

1- DataBase of Single Nucleotide Polymorphism

Table 5. Stability analysis of E18D variant

dbSNP ID	Amino acid substitution	I-Mutant 2.0			iPTREE-STAB	
		Stability	RI	DDG	The discriminated direction of thermal stability change	The predicted value of thermal stability change (Kcal/mol)
rs61730117	E18D	Decrease	6	-0.57	Decrease	-0.1221

Table 6. ModPred analysis of E18D variant

Residue	Modification	Score	Confidence
E18D	Proteolytic cleavage	0.51	Low
	Carboxylation	0.77	Medium
	ADP-ribosylation	0.63	Low

Table 7. QMEAN score of E18D variant

Parameters	Score
C_beta interaction energy	-88.52 (Z-score: -1.71)
All-atom pairwise energy	-14526.09 (Z-score: -1.93)
Solvation energy	-11.40 (Z-score: -2.67)
Torsion angle energy	-104.71 (Z-score: -3.11)
Secondary structure agreement	77.1% (Z-score: -0.33)
Solvent accessibility agreement	67.8% (Z-score: -2.43)
Total QMEAN-score	0.472 (Z-score: -3.22)

Conservation profiling of UTY: Amino acids are highly conserved when located in biologically active sites. If there is substitution of these amino acids, it leads to complete loss of biological activities (38). For this purpose, ConSurf server was used to predict the degree of evolutionary conservation in the protein UTY at each amino acid position. Although there is complete analysis of UTY protein, only those amino acids have been focused which were selected as high risk nsSNP. The conservational analysis revealed that E18D is highly conserved with a score of 8. Also, E18D is a conserved amino acid, so its functional role is very critical.

Post translational modification sites on UTY: To analyze the effect of nsSNPs on post translational modification process of human UTY protein, Mod Pred tool was used. ModPred predicted sites for proteolytic cleavage, carboxylation and ADP-ribosylation at E18 (Table 6).

3D modelling and biophysical validation of UTY: SPARKS-X modeled the 3D structure of UTY where 10 full length models were generated. The quality of full length models were predicted by z-score (Z-score >6). Using SPARKS-X, native model as well as mutated model with the most deleterious variant (E18D) was generated (Figure 1).

QMEAN tool was used where z-score gives prediction about the absolute quality of the model (Table 7). The quality was estimated by comparing other similar reference structures (39). The total QMEAN score was 0.472 along with its z-score -3.22 which comes under calculated model reliability value ranging between 0-1. According to SPARKS-X, z-score should be greater than 6 and the predicted model z-score was 15.16. This indicated that the predicted model was of good type.




Figure 1. A: Wild-type model showing glutamic acid at position 18 (GLU18) of UTY protein constructed using SPARKS-X and visualized by PyMol. B: Mutant model showing aspartic acid at position 18 (ASP18) of UTY protein constructed using SPARKS-X and visualized by PyMol

SPARKS-X predicted model was validated using another tool known as PROCHECK where Rama-chandran plot was used. Out of all the residues of UTY protein, 1024 (85.3%) were in most favorable regions, 148 (12.3%) were in additionally allowed regions, 24 (2.0%) were in generously allowed regions and 5 (0.4%) were in disallowed regions (Figure 2). Therefore, UTY protein structure can be considered as a relevant model.

Project HOPE was used to know the structural changes but no such information was retrieved. Therefore, HOPE used UniProt database and Reprot software was used to predict the mutational analysis. E18D substitution results into a change in glutamic acid residue to an aspartic acid residue at position 18 (Figure 3). The mutated residue (Aspartic acid-D) is smaller than the wild type residue (Glutamic acid-E) which may lead to loss of interaction.

Phen 2, PANTHER, PROVEAN, I-Mutant 2 and iPTREE-STAB for the screening and identification of functional mutations in UTY gene. By analyzing all the missense mutations through these tools, one nsSNP with position E18D was predicted to be high risk nsSNP.

The ConSurf tool output indicated that the nsSNP at position E18 was predicted in highly conserved region and may have promising role in UTY protein function. In addition, the E18 residue showed PTM sites for proteolytic cleavage, carboxylation and ADP-ribosylation. Therefore, amino acid polymorphism at E18 position is likely to have impact on post translational modification process of human UTY protein.

For UTY protein, 3 dimensional models were generated using SPARKS-X to visualize deviation between the wild type and mutant type protein models. According to HOPE in E18D, the wild type protein framework was disturbed due to shift of glutamic acid to aspartic acid upon missense mutation which may cause loss of interaction.

Validation of SPARKS-X generated model using QMEAN and PROCHECK servers indicated that model of UTY is good for further experiment as well as for better understanding of biological activity of UTY protein. Project HOPE was used to understand structural changes upon amino acid polymorphism. The wild and mutant amino acids differ sometimes due to specific properties which can disturb the structural and/or functional features of the native protein (47). Similarly, in missense mutation in the UTY protein at amino acid position 18 where Glu is converted to Asp, the mutant residue is smaller and loss of interaction can occur because of the smallness. Analysis of protein-protein interaction is one of the best ways to find out the organization of proteomes related to functional network (48). The use of functional network view for a particular genome is carried out to refine the statistical potential for human molecular genetics (49). STRING was used for protein-protein interaction analysis; reported interaction of UTY protein with different proteins may have an impact on many other pathways involved in the disease.

Conclusion

The present study symbolized the first detailed analysis where sequence and structure based algorithms are used to identify functional nsSNPs in UTY gene. Out of 151 nsSNPs, 1 (0.66%) was predicted to be deleterious by SIFT and PolyPhen 2.

Further analyses support that E18D variant is predicted to cause change in stability and functional interaction performance of UTY protein. This computational analysis of UTY will assist to understand and design future experimental research.

Acknowledgement

The authors express appreciation to Charutar Vidya Mandal (CVM) and SICART, Vallabh Vidyanagar, Gujarat for providing research work platform. We acknowledge the assistance of the Director, Ashok and Rita Patel Institute of Integrated Study and Research in Biotechnology and Allied Sciences (ARIBAS), New Vallabh Vidyanagar for providing all the facilities and constant encouragements to carry out this work. This work was supported by INSPIRE division of the Department of Science and Technology in the form of research fellowship to Ms. Mili Nailwal.

Conflict of Interest

The authors report no conflict of interest.

References

1. Naasse Y, Charoute H, El Houate B, Elbekkay C, Razoki L, Malki A, et al. Chromosomal abnormalities and Y chromosome microdeletions in infertile men from Morocco. *BMC Urol.* 2015;15:95.
2. Gurunath S, Pandian Z, Anderson RA, Bhattacharya S. Defining infertility--a systematic review of prevalence studies. *Hum Reprod Update.* 2011;17(5):575-88.
3. Kleiman SE, Almog R, Yogev L, Hauser R, Lehavi O, Paz G, et al. Screening for partial AZFa microdeletions in the Y chromosome of infertile men: is it of clinical relevance? *Fertil Steril.* 2012;98(1):43-7.
4. Nailwal M, Chauhan JB. Genetic investigation at azoospermia factor region for male infertility. *Indo Am J Pharm Sci.* 2016;3(12):1498-1510.
5. Masoudi R, Mazaheri-Asadi L. Y chromosome microdeletions in infertile men from south west of Iran. *International Conference on Chemical, Environmental and Biological Sciences*; 2015 March 18-19; Dubai: International Institute of Chemical, Biological and Environmental Engineering. 2015. p. 110-12.
6. Ambulkar P, Chuadhary A, Waghmare J, Tarnekar A, Pal A. Prevalence of Y Chromosome Microdeletions in Idiopathic Azoospermia Cases in Central Indian Men. *J Clin Diagn Res.* 2015;9(9):GC01-4.
7. Sun C, Skaletsky H, Rozen S, Gromoll J, Nieschlag E, Oates R, Page DC. Deletion of azoospermia fac-

- tor a (AZFa) region of human Y chromosome caused by recombination between HERV15 proviruses. *Hum Mol Genet.* 2000;9(15):2291-6.
8. Greenfield A, Carrel L, Pennisi D, Philippe C, Quaderi N, Siggers P, et al. The UTX gene escapes X inactivation in mice and humans. *Hum Mol Genet.* 1998;7(4):737-42.
 9. Shen P, Wang F, Underhill PA, Franco C, Yang W H, Roxas A, et al. Population genetic implications from sequence variation in four Y chromosome genes. *Proc Natl Acad Sci USA.* 2000;97(13):7354-9.
 10. Abilash VG, Ramgir Sh. Genetic and environmental factors involved in human male infertility: a review. *Asian J Pharm Clin Res.* 2015;8(6):34-43.
 11. Foresta C, Moro E, Ferlin A. Y chromosome microdeletions and alterations of spermatogenesis. *Endocr Rev.* 2001;22(2):226-39.
 12. Vineeth VS, Malini SS. A journey on Y chromosomal genes and male infertility. *Int J Hum Genet.* 2011;11(4):203-15.
 13. Drazen JM, Yandava CN, Dubé L, Szczerback N, Hippensteel R, Pillari A, et al. Pharmacogenetic association between ALOX5 promoter genotype and the response to anti-asthma treatment. *Nat Genet.* 1999;22(2):168-70.
 14. Akhouni F, Parvaneh N, Modjtaba EB. In silico analysis of deleterious single nucleotide polymorphisms in human BUB1 mitotic checkpoint serine/threonine kinase B gene. *Meta Gene.* 2016;9:142-50.
 15. Rasal KD, Shah TM, Vaidya M, Jakhesara SJ, Joshi CG. Analysis of consequences of non-synonymous SNP in feed conversion ratio associated TGF- β receptor type 3 gene in chicken. *Meta Gene.* 2015;4:107-17.
 16. Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc.* 2009;4(7):1073-81.
 17. Sen S, Pasi AR, Dada R, Shamsi MB, Modi D. Y chromosome microdeletions in infertile men: prevalence, phenotypes and screening markers for the Indian population. *J Assist Reprod Genet.* 2013;30(3):413-22.
 18. Liu XG, Hu HY, Guo YH, Sun YP. Correlation between Y chromosome microdeletion and male infertility. *Genet Mol Res.* 2016;15(2).
 19. Benoff SH, Millan C, Hurley IR, Napolitano B, Marmar JL. Bilateral increased apoptosis and bilateral accumulation of cadmium in infertile men with left varicocele. *Hum Reprod.* 2004;19(3):616-27.
 20. Ng PC, Henikoff S. SIFT: Predicting amino acid changes that affect protein function. *Nucleic Acids Res.* 2003;31(13):3812-4.
 21. Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, et al. A method and server for predicting damaging missense mutations. *Nat Methods.* 2010;7(4):248-9.
 22. Thomas PD, Kejariwal A, Guo N, Mi H, Campbell MJ, Muruganujan A, et al. Applications for protein sequence-function evolution data: mRNA/protein expression analysis and coding SNP scoring tools. *Nucleic Acids Res.* 2006;34(Web Server issue):W645-50.
 23. Choi Y, Sims GE, Murphy S, Miller JR, Chan AP. Predicting the functional effect of amino acid substitutions and indels. *PLoS One.* 2012;7(10):e46688.
 24. Capriotti E, Fariselli P, Casadio R. I-Mutant2.0: predicting stability changes upon mutation from the protein sequence or structure. *Nucleic Acids Res.* 2005;33(Web Server issue):W306-10.
 25. Huang LT, Saraboji K, Ho SY, Hwang SF, Ponuswamy MN, Gromiha MM. Prediction of protein mutant stability using classification and regression tool. *Biophys Chem.* 2007;125(2-3):462-70.
 26. Ashkenazy H, Erez E, Martz E, Pupko T, Ben-Tal N. ConSurf 2010: calculating evolutionary conservation in sequence and structure of proteins and nucleic acids. *Nucleic Acids Res.* 2010;38(Web Server issue):W529-33.
 27. Pejaver V, Hsu WL, Xin F, Dunker AK, Uversky VN, Radivojac P. The structural and functional signatures of proteins that undergo multiple events of post-translational modification. *Protein Sci.* 2014;23(8):1077-93.
 28. Yang Y, Faraggi E, Zhao H, Zhou Y. Improving protein fold recognition and template-based modeling by employing probabilistic-based matching between predicted one-dimensional structural properties of query and corresponding native properties of templates. *Bioinformatics.* 2011;27(15):2076-82.
 29. Venselaar H, Te Beek TA, Kuipers RK, Hekkelman ML, Vriend G. Protein structure analysis of mutations causing inheritable diseases. An e-Science approach with life scientist friendly interfaces. *BMC Bioinformatics.* 2010;11:548.
 30. Benkert P, Biasini M, Schwede T. Toward the estimation of the absolute quality of individual protein structure models. *Bioinformatics.* 2011;27(3):343-50.
 31. Laskowski RA, MacArthur MW, Moss DS, Thorn-

- ton JM. PROCHECK: a program to check the stereochemical quality of protein structures. *J App Cryst.* 1993;26:283-91.
32. von Mering C, Jensen LJ, Snel B, Hooper SD, Krupp M, Foglierini M, et al. STRING: known and predicted protein-protein associations, integrated and transferred across organisms. *Nucleic Acids Res.* 2005;33(Database issue):D433-7.
 33. Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, et al. PANTHER: a library of protein families and subfamilies indexed by function. *Genome Res.* 2003;13(9):2129-41.
 34. Teng S, Srivastava AK, Wang L. Sequence feature-based prediction of protein stability changes upon amino acid substitutions. *BMC Genomics.* 2010;11 Suppl 2:S5.
 35. Benkert P, Tosatto SC, Schomburg D. QMEAN: A comprehensive scoring function for model quality assessment. *Proteins.* 2008;71(1):261-77.
 36. Datta A, Mazumder MH, Chowdhury AS, Hasan MA. Functional and structural consequences of damaging single nucleotide polymorphisms in human prostate cancer predisposition gene RNASEL. *Biomed Res Int.* 2015;2015:271458.
 37. Jensen LJ, Kuhn M, Stark M, Chaffron S, Creevey C, Muller J, et al. STRING 8--a global view on proteins and their functional interactions in 630 organisms. *Nucleic Acids Res.* 2009;37(Database issue):D412-6.
 38. Miller MP, Kumar S. Understanding human disease mutations through the use of interspecific genetic variation. *Hum Mol Genet.* 2001;10(21):2319-28.
 39. Srivastava M, Gupta SK, Abhilash PC, Singh N. Structure prediction and binding sites analysis of curcumin protein of *Jatropha curcas* using computational approaches. *J Mol Model.* 2012;18(7):2971-9.
 40. Chen X, Sullivan PF. Single nucleotide polymorphism genotyping: biochemistry, protocol, cost and throughput. *Pharmacogenomics J.* 2003;3(2):77-96.
 41. Nailwal M, Chauhan JB. Analysis of consequences of non-synonymous SNPs of USP9Y gene in human using bioinformatics tools. *Meta Gene.* 2017; 12:13-7.
 42. Nailwal M, Chauhan JB. In silico analysis of non-synonymous single nucleotide polymorphisms in human DAZL gene associated with male infertility. *Syst Biol Reprod Med.* 2017;1-11.
 43. de Carvalho MD, De Mesquita JF. Structural modeling and in silico analysis of human superoxide dismutase 2. *PLoS One.* 2013;8(6):e65558.
 44. George Priya Doss C, Rajith B. A new insight into structural and functional impact of single-nucleotide polymorphisms in PTEN gene. *Cell Biochem Biophys.* 2013;66(2):249-63.
 45. Thusberg J, Vihinen M. Pathogenic or not? And if so, then how? Studying the effects of missense mutations using bioinformatics methods. *Hum Mutat.* 2009;30(5):703-14.
 46. Hicks S, Wheeler DA, Plon SE, Kimmel M. Prediction of missense mutation functionality depends on both the algorithm and sequence alignment employed. *Hum Mutat.* 2011;32(6):661-8.
 47. Hussain MR, Shaik NA, Al-Aama JY, Asfour HZ, Khan FS, Masoodi TA, et al. In silico analysis of Single Nucleotide Polymorphisms (SNPs) in human BRAF gene. *Gene.* 2012;508(2):188-96.
 48. Mohamoud HS, Hussain MR, El-Harouni AA, Shaik NA, Qasmi ZU, Merican AF, et al. First comprehensive in silico analysis of the functional and structural consequences of SNPs in human GalNAc-T1 gene. *Comput Math Methods Med.* 2014;2014:904052.
 49. Szklarczyk D, Franceschini A, Kuhn M, Simonovic M, Roth A, Minguetz P, et al. The STRING database in 2011: functional interaction networks of proteins, globally integrated and scored. *Nucleic Acids Res.* 2011;39(Database issue):D561-8.

Downloaded from <http://www.jri.ir>