Effects of *Ferulago angulata* Extract Against Oxidative Stress Induced by 6-hydroxydopamine in Rats

Sharifi F (M.Sc. Student)¹, Rafieirad M (Ph.D.)²*, Sazegar H (Ph.D.)³

1-Department of Physiology, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

2- Department of Biology, Izeh Branch, Islamic Azad University, Izeh, Iran

3-Department of Physiology, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

* Corresponding author: Department of Biology, Izeh Branch, Islamic Azad University, Izeh, Iran

Tel: +98-916-3116542, Fax: +98-61-43643374 Email: Rafieirad.m@gmail.com

Received: 7 Oct. 2014

Accepted: 18 Feb. 2015

Abstract

Background: Parkinson's disease is one of the prevalent degenerative diseases of the neural system. Oxidative stress, which has been recognized as the most important factor of Parkinson's disease, plays a main role in the death of neurons in this disease. Antioxidants have a proved role in the prevention of oxidative stress.

Objective: Considering the presence of evidence regarding antioxidant property of *Ferulago angulata*, effect of oral administration of hydroalcoholic extract of this plant was investigated on parameters of the oxidative damage caused by Parkinson's disease in adult rats.

Methods: In this research, 40 Wistar rats with the average weight of 200 to 250 g were randomly assigned to 5 groups, each with 8 rats. Also, to develop Parkinson's disease animal model, 2 μ l of 6-hydroxydopamine drug was injected into medial forebrain bundle on the left side of the animals and the treatment groups received *Ferulago angulata* extract with doses of 400, 200, and 100 mg/kg for 2 weeks as gavage. Then, the brain of the animals was isolated and parameters of Malondialdehyde and Thiol were measured.

Results: Results showed that Parkinson's disease induced significantly increased brain peroxidation lipid, which is one of the factors of oxidative stress, and reduced Thiol level (p<0.001). Further, treatment of Parkinson's disease with *Ferulago angulata* extract significantly decreased peroxidation lipid in striatum and hippocampus and increased Thiol level (p<0.001).

Conclusion: oral administration of hydroalcoholic extract of *Ferulago angulata* probably reduces oxidative damage and also reduces Malondialdehyde and increases Thiol in the treated groups due to its strong antioxidant property.

Keywords: Ferulago angulata, Malondialdehyde, Parkinson's disease, Thiol groups



Introduction

Parkinson's disease (PD) which is one of the prevalent neurodegenerative disorders is the most prevalent neuropathological disorder induced degeneration that is by of dopaminergic neurons of the dense part of substantia nigra and their terminals in striatum. Destruction of these neurons reduces dopamine neurotransmitter in this area [1, 2]. Reduction of dopamine leads to disorders such as bradykinesia, vibration, catalepsy, and posture instability [3]. Oxidative stress forms free radicals which are created in basal ganglia conditions, lead under pathologic to progressive damage and degeneration of nuclear neurons of substantia nigra, and induce Parkinson's disease [4, 5]. Other factors such as increase of lipid peroxidation, DNA destruction, and accumulation of iron play some role in the degeneration of dopaminergic neurons and Parkinson's disease [6]. Endogenous sources of oxidative stress include free radicals resulting from dopamine metabolism. Active radicals of oxygen are continually produced by Monoaminooxidase B enzyme or dopamine autoxidation in the dopaminergic neurons of medial brain due to metabolism of dopamine [7]. Oxidative stress is induced in brain when production of free radicals is higher than the ability of antioxidant system to delete additional reactive oxygen species (ROS)³. Natural chemical compounds with strong antioxidant activity are found in high concentrations in some plants [8]. Role of many plants has recently attracted the attention of human societies owing to their antioxidant property in treatment of neurogenerative diseases. Studies have shown that plant phenols such as flavonoids and phenolic acids can act as effective antioxidants for molecular level [9]. Secondary metabolites derived from plants such as phenol and total

flavonoid have a strong potential to clean free radicals and the plants which are enriched with antioxidant compounds can protect cells against oxidative damage [10, 11]. Ferulago angulata (FA) plant belongs to Apiaceae family and is a natural and native plant of mountainous regions of west of Iran [12]. Its extract which contains phenolic compounds and has antioxidant and anti-diabetic effects is considered one of important herbs in Iran [13-15]. According to studies by Soudfian et al. (2011), analysis of aerial parts of different species of ferulago by ethanol has led to identification of its several components, the most important of which are cis-ocimene, α pinene, β -germacrene, σ -terpinene, trans- β ocimene, germacrene D, limonene, bornyl camphene, acetate. myrcene, allo-neoocimene, *B*-pinene, bicyclogermacrene and sophrosyne, respectively [13]. Phenolic compounds, due to having antioxidant activity, are capable of inhibiting free radicals; thus, they could be effective in preventing many diseases like cancer. cardiovascular and nervous diseases [16]. This plant is applied as an inhibitor of acetylcholine [17]. It has also antimicrobial, antioxidant [18], and cytotoxic [19] properties., it is used as a flavor and strong tranquilizer of digestive system, and also for its antihelmentic property [20, 21]. Studying oxidative stress is done using different indices, one of which is the end peroxidation product of lipid or Malondialdehyde (MDA). This aldehyde is produced in a biomarker organism for the measurement of oxidative stress level. One of the other indices of studying oxidative stress is Thiol index. The most important role of Thiol in in vivo studies is its performance as an intracellular and extracellular reductive buffer which indicates that effect of Thiol bands as a defensive antioxidant mechanism has been



reduced. Reduction of Thiol level indicates reduction of antioxidant capacity of the brain [22]. Peroxidation of lipids changes chemical characteristics and physical structure of the cell membrane and reduces function. It also has a role in the pathogenesis of diseases such neurogenerative as diseases including Parkinson's disease [23]. Different molecules like aldehyde are created by peroxidation of lipids, one of which is MDA that is one of the indices of oxidative stress caused bv Parkinson's disease [24]. Thus in the current study, neuroprotective (neuronal protection) effects of Ferulago angulata extract (FAE) were investigated in animal model of Parkinson's disease through studying the effect oxidative induced on stress by 6hydroxydopamine in rats.

Materials and Methods

Animals and experimental groups

In this research, Wistar rats with the weight range of 200-250 g from the animal house of Ahvaz University of Medical Sciences were used for reproduction and maintenance. The animals were kept in a bright-dark place for 12 h at 21±2°C with free access to compressed food of Animal Feed Company, Shahreza, Isfahan, and piped, treated water of Izeh city, and were maintained in groups (each with four rats) inside standard cages in Animal center of Islamic Azad University, Izeh Branch. Protocol of this research was prepared based on the international regulations regarding laboratory animals and enacted in ethics committee of the University. The animals were randomly divided into the following groups:

1-Control group without inducing Parkinson's disease and treatment

2-Parkinson's group wihout treatment

3-Parkinson's group treated with 100 mg/kg of FAE with gastric gavage (i.g)



Journal of Medicinal Plants, Volume 14, No. 53, Winter 2015 4-Parkinson's group with 200 mg/kg of FAE with gastric gavage (i.g)

5-Parkinson's group treated with 400 mg/kg of FAE with gastric gavage (i.g) [25].

Parkinson's experimental model

The animals were anaesthetized by the intraperitoneal injection of 90 mg/kg hydrochloride ketamin + 10 mg/kg xylazine and then heads of the rats were fixed in stereotaxy for the brain surgery. Afterward, µI of 6-hydroxydopamine (6-OHDA) 2 neurotoxic drug purchased from American Sigma Company was injected into medial forebrain bundle (MFB) on the left side of the brain of the animals and the treatment groups received Ferulago angulata with different doses for 2 weeks as gavage after three-day recovery. Then, the brain of the animals was isolated and parameters of MDA and Thiol were measured [26].

Preparation Ferulago angulata extract

Ferulago angulata plant was collected in early spring from the surrounding of Izeh city, Iran, and then its leaves were separated and dried in open air under shade for 2 weeks. After drying, the leaves were finely powdered (with diameter of less than 0.4 mm) with an electric grinder and *Ferulago angulata* powder was rinsed in ethanol 72% for 3 days at room temperature. *Ferulago angulata* powder and ethanol mix were stirred occasionally and then alcohol and powder mix were finely filtered to obtain its extract. The obtained extract was distilled in vacuum to evaporate its alcohol completely. Finally, FAE was obtained (yield 29%) as a brown powder after full evaporation [27].

Malondialdehyde (MDA) Assays

In this test, the groups with 8 rats were used. Tissues of striatum and hippocampus

were homogenized separately with the specified amount of KCL 1.5 %. 0.5 ml of the homogenized solution was removed and 2.5 ml of the TCA solution 3% was added and kept in the water bath at 37°C for 10 min. Then, it was centrifuged at 3000 rpm for 10 min. 0.5 ml of the supernatant solution was removed after the centrifuge and 3 ml of phosphoric acid solution 0.1 and 1 ml of Thiobarbituric acid (TBA) 0.67 were added to each of them and put in boiling water for 45 min. The tubes were cooled in the ice container and 4 ml of butanol was added to each one. After vortex mixture, it was centrifuged at 2000 rpm for 20 min and finally absorption with the wavelength of 532 nm was read. After putting the numbers obtained from spectrophotometer and absorption а in standard curve line equation, concentration of MDA was evaluated [28].

Standard curve

Standard curve had to be first drawn and it was necessary to prepare MDA standard solution and measure wavelengths with the spectrophotometer (Figure 1). 0.5 ml of the standard solution with concentrations of 0.5,1, 2, 4, 6, 8, and 10 was removed and then 3 ml of the phosphoric acid 1% solution was added and other stages were performed like the above stages. Finally, absorption was read with the wavelength of 532 nm [28].

Total sulphydryl (-SH) groups assay

Total SH groups were measured using DTNB as the reagent. This reagent reacts with the -SH groups to produce a vellow coloured complex which has peak absorbance at 412 nm (Ellman, 1959). Briefly, 1 ml Tris-EDTA buffer (pH = 8.6) was added to 50μ L homogenate in 2 ml cuvettes and sample absorbance was read at 412 nm against Tris-EDTA buffer alone (A₁). Then, 20 μ L DTNB reagent (10 mM in methanol) was added to the mixture and after 15 min (stored in room temperature) the sample absorbance was read again (A_2) . The absorbance of DTNB reagent was also read as a blank (B). Total thiol concentration (mM) was calculated from the following equation: Total thiol concentration $(mM) = (A_2 - A_1 - B) \times 1.07 / 0.05 \times 13.6.$ [29].



Figure 1- The Standard Curve of MDA was constructed Over the Concentration Range of 0-10 µM



Statistical Analysis

Data of this research were presented as mean \pm SEM and then analyzed using suitable statistical methods in Excel and SPSS software by one way ANOVA and LSD test methods. Difference of the results between different groups was significant with minimum p<0.05.

Results

In this research, all the animals properly tolerated stereotaxic surgery and no mortality was observed during the study. The results of this study showed that the amount of MDA as lipid peroxidation index significantly increased in Parkinson's group compared with the control group in striatum and hippocampus tissues after MFB lesion due to the injection of 6-OHDA (p<0.001). At the next, comparison of MDA measurement between the Parkinson's group without treatment and treatment Parkinson's group which orally received the extract of Ferulago angulata with doses of 200 and 400 mg/kg for 14 days specified that MDA was significantly reduced in the group receiving FAE (p<0.05), while no significant reduction was found in the group receiving

100 mg/kg of the extract of *Ferulago angulata* in this tissue compared with the Parkinson's group. Comparison of MDA measurement between the Parkinson's group and Parkinson's groups which received extract of *Ferulago angulata*, 200 and 400 mg/kg led to finding a significant reduction in the striatum tissue (p<0.01), while it was not effective for the amount of MDA in this tissue in the group receiving FAE (100 mg/kg) (Figures 2 and 4).

In another section of this research, it was found that Thiol level was significantly decreased in Parkinson's group compared with the control in hippocampus tissue after MFB lesion in rats, and decreased significantly in striatum (p<0.001). Comparison between the Parkinson's group and treatment Parkinson's groups which orally received extract of Ferulago angulata, 100, 200 and 400 mg/kg for 14 days demonstrated that Thiol level significantly increased in hippocampus tissue in the groups receiving Ferulago angulata compared with Parkinson's group (p<0.05) and significant increase was also found in Thiol level in striatum tissue (p<0.001) (Figures 3 and 5).



Figure 2- Effect of *Ferulago angulata* on MDA levels in hippocampus tissue between Control group, PD and PD groups orally receiving 100, 200 and 400 mg/kg FAE for 14 days. Values are expressed as mean ± SEM (n = 8). *** p < 0.001 vs. control rats, #p < 0.05 vs. PD-treated rats





Figure 3- Effect of *Ferulago angulata* on thiol levels in hippocampus tissue between Control group, PD and PD groups orally receiving 100, 200 and 400 mg/kg FAE for 14 days. Values are expressed as mean ± SEM (n = 8). ** p < 0.01 vs. control rats, #p < 0.05 vs. PD-treated rats



Groups

Figure 4- Effect of *Ferulago angulata* on MDA levels in striatum tissue between Control group, PD and PD groups orally receiving 100, 200 and 400 mg/kg FAE for 14 days. Values are expressed as mean ± SEM (n = 8).



Figure 5- Effect of *Ferulago angulata* on thiol levels in striatum tissue between Control group, PD and PD groups orally receiving 100, 200 and 400 mg/kg FAE for 14 days. Values are expressed as mean ± SEM (n = 8). *** p < 0.001 vs. control rats, ###p < 0.001 vs. PD-treated rats



Discussion

In this research, effect of unilateral injection of 6-OHDA neurotoxic drug into MFB and its resulting oxidative stress after lesion and treatment with hydroalcoholic extract of Ferulago angulata were studied. The findings in this research showed that Parkinson's animal model in rats by the injection of 6-OHDA into MFB through reducing antioxidant substances in brain caused death of dopaminergic neurons inside the substantia nigra and striatum and oral administration of different amounts of FAE for two weeks was effective for the rats with Parkinson's disease, which led to reduction of MDA and increase of Thiol group, and prevented the disease progress. In this study, MFB lesion was used, because developing Parkinson's by the injection of 6-OHDA neurotoxic drug into MFB is more similar to the pathology of this disease in humans than other models and its symptoms are closer to human Parkinson's disease [30]. Based on the conducted studies. Parkinson's disease is accompanied by neuron oxidative degenerate ion and substantia nigra is specifically susceptible to oxidative damage due to (dopamine oxidative content and neuromelanin) and relatively low antioxidant content [2]. The evident characteristic of this disease is pathologically slow and gradual degeneration of dopaminergic neurons of substantia nigra [5,31]. Studies have that unilateral represented damage of dopaminergic system reduced dopamine level and increases regulation of dopaminergic postsynaptic receptors located in striatum neurons of the damaged side through the injection of 6-OHDA [32]. Oxidative stress weakens antioxidant systems of the brain by producing free radicals and other reactive oxygen species (ROS) and the brain is



Journal of Medicinal Plants, Volume 14, No. 53, Winter 2015 damaged by its destructive effects [33]. According to the conducted studies, free radicals and other ROS are important factors degenerative disorders for including Parkinson's disease, Alzheimer's disease, and some hepatic diseases and cardiovascular disorders [34]. ROS production and inducing oxidative stress are the known factors for the functional disorder of Parkinson's disease which oxidize lipids and proteins in the central nerves system [33]. In Parkinson's disease, hippocampus striatum and are more susceptible to damage caused by oxidative stress and free radicals than other parts of the brain [33]. In this disease, concentration of fatty acids has been reduced in substantia nigra, while MDA which is oxidative stress index is increased and also Thiol are decreased. Reduction of Thiol level indicates reduced antioxidant capacity of the brain [35]. One of the factors in the level of proteins which plays an effective role in its antioxidant properties is Thiol group and reduction of this group can be a good indicator for oxidative injuries [36]. Glutathione is one of the Thiol groups which plays an essential role in the protection of cells against degeneration caused by peroxidase hydrogen and oxygen species and these active oxygen species result from the disorder in dopamine metabolism. This disorder in dopamine metabolism reduces glutathione in nerve ending and degenerative diseases [37]. Many studies have shown that 6-OHDA neurotoxic drug creates its toxicity by creating free radicals [38]. This toxic substance is rapidly oxidized under physiological conditions and is converted into hydrogen peroxide which is one of the most destructive free radicals for live cells [39]. Many natural and artificial antioxidants are accessible which can be used for the treatment of Parkinson's disease [40]. Plants have

recently attracted the attention of human communities as natural antioxidants. Recent studies have demonstrated that Ferulago angulata contains phenol compounds [12]. It has been also recognized as one of the native and medicinal plants for the treatment of diseases [41]. This plant is also used for treating digestive diseases and chronic ulcer due to snakebite and headache [41]. In addition, antioxidant and antidiabetic effects of Ferulago angulata have been reported [42]. Aryan Poor et al. (2006) showed that antioxidants had a protective effect on Parkinson's disease [43]. Results of Salar et al. (2010) showed that hydraulic extract of plants improved Parkinson's disease in male rats due to antioxidant property [44]. Balooch Nejad et (2011) reported that intraperitoneal al. injection of antioxidants had a protective effect on 6-OHDA in Parkinson's experimental model [45]. Rafiei Rad et al. (2014) reported that different doses of FA was able to reduce MDA as a lipid peroxidation in ischemia [25]. Studies showed that FAE led to the reduction of MDA [46]. In other studies conducted by Mansouri et al. (2013), it was found that level of Thiol groups was significantly reduced after treatment with different doses of Gallic acid

by MFB lesion due to administration of 6-OHDA in rats [29]. The present results in this research were in line with those of the above studies and, considering findings of others and results of this research, it can be suggested that Ferulago angulata extract of has an antioxidant property and its consumption significantly increases antioxidant capacity. According to the conducted studies. antioxidant activity of *Ferulago angulata* may be due to the presence of phenols, particularly polyphenol compounds which are used as a natural antioxidant in food and pharmaceutical industries [12].

Conclusion

In this study, for the first time, it was shown that treatment with different doses of FAE could reduce MDA and increase level of Thiol in Parkinson's animal model.

Acknowledgment

The authors would thank the Research Deputy of Islamic Azad University, Izeh Branch, who was accompanied to carry out this research.

References

1. Sauer H and Oertel WH. Progressive degeneration of nigrostriatal dopamine neurons following intrastriatal terminal lesions with 6-hydroxydopamine: a combined retrograde tracingand immunocytochemical study in the rat. *Neuroscience* 1994; 59 (2): 401 - 15.

2. Gang Hu, Siamak B, Pekka J and Riitta A. Coffee and Tea consumption and the risk of

Parkinson's disease. *Move Disord* 2007; 22 (15): 224 - 8.

3. Sugawara T and Chan P. Reactive oxygen radicals and pathogenesis of neuronal death after cerebral ischemia. *Ant Red Signall* 2003; 5 (5): 597 - 607.

4. Chan PH. Reactive oxygen radicals in signaling and damage in theischemic brain. *J. Cereb Blood Flow Metab.* 2001; 21 (1): 2 - 14.



Sharifi et al.

5. Dauer W and Przedborski S. Parkinson's disease: mechanisms and models neuron. *Department Neurol* 2003; 39 (6): 889 - 909.

6. Schwarting RK, Huston JP. Behavioral and neurochemical dynamics of neurotoxic meso-striatal dopamine lesions. *Neurotoxicol.* 1997; 18: 689 - 708.

7. Valenzuela A and Garrido A. Biochemical bases of the pharmacological action of the flavonoid silymarin and of its structural isomer silibinin. *Biol. Res.* 1994; 27 (2): 105 - 12.

8. Pereira R, Fachinetto R and de Souza Prestes A. Antioxidant Effects of Different Extracts from Melissa officinalis, Matricaria recutita and Cymbopogon citratus. *Neurochemical Research* 2009; 34 (5): 973 -83.

9. Craig WJ. Health-promoting properties of common herbs. *Am. J. Clin. Nutr.* 1999; 70 (3): 491 - 9.

10. Kumaran A and Karunakaran RJ. Antioxidant and free radical scavenging activity of an aqueous extract of Coleus aromaticus. *Food Chem.* 2006; 97 (1): 109 -14.

11. Mathew S and Abraham TE. In vitro antioxidant activity and scavenging effects of Cinnamomum verum leaf extract assayed by different methodologies. *Food Chem. Toxicol.* 2006; 44 (2): 198 - 206.

12. Hosseini N, Akbari M, Ghafarzadegan R and Changizi Ashtiyani S.R.S. Total phenol, antioxidant and antibacterial activity of the essential oil and extracts of Ferulago angulata ssp. Angulata. *J. Med. Plants* 2012; 11 (43): 80 - 9.

13. Sodeifian G, Ansari K, Bamoniri A and Mirjalili BF. Study of chemichal composition

of the essential oil of *ferulago angulata* (Schelchet) boiss. from iran using super critical fluid extraction and nano scale injection. *Digest Journal of Nanomaterials and Biostructures* 2011; 6 (1): 161 - 8.

14. Zhou C, Huang Y and Przedborski S. Chemical Analysis and Antimicrobial Studies on Three Species of Ferulago from Greece. *Planta Med.* 2000; 66 (6): 560 - 3.

15. Taran M, Ghasempour HR, Shirinpour E. Antimicrobial activity of essential oils of Ferulago angulata subsp. *Jundishapur Journal of Microbiology* 2010; 3 (1): 10 - 14.

16. Hosseini N, Akbari M, Ghafarzadegan R, Changizi Ashtiyani S and Shahmohammadi R. Total Phenol, Antioxidant and Antibacterial Activity of the Essential Oil and Extracts of Ferulago angulata ssp. Angulate. *Journal of Medicinal Plants* 2012; 11 (43): 80 - 9.

17. Dall'Acqua S, Maggi F and Minesso P. Identification of nonalkaloid acetylcholinesterase inhibitors from Ferulago campestris (Besser) Grecescu (Apiaceae). *Fitoterapia* 2010; 81 (8): 1208 - 12.

18. Basile A, Sorbo S and Spadaro V. Antimicr al and antioxidant activities of coumarins from the roots of Ferulago obi campestris (Apiaceae). *J Molecules* 2009; 14: 939 - 52.

19. Rosselli S, Maggio AM, Faraone N and Spadaro V. Bastow Km roots of *Ferulago campestris* (Apiaceae) and ofsynthetic ester derivatives of aegelinolF. The cytotoxic properties of natural coumarins isolated fro. *Nat. Prod. Commun.* 2009; 4: 1701 - 6.

20. Baser KH, Demirci B, Demirci F, Hashimoto T and Asakawa Y. Ferulagone: A new monoterpene ester from Ferulago thirkeana essential oil. *Planta Med.* 2002; 68 (6): 564 - 7.



www.SID.ir

21. Ozturk B, Gur S and Coskun M. Relaxant effect of Ferulago syriaca root extract on human corpus cavernosum. *Eur. Urol. Suppl.* 2004; 3: 62.

22. Acikgoz O, Gonenc S and Kayatekin BM. Methamphetamine causes lipid peroxidation and an increase in superoxidedismutase activity in the rat striatum. *Brain Research* 1998; 813 (1): 200 - 2.

23. Moreira PI, Nunomura A and Nakamura M. Nucleic acid oxidation in Alzheimer disease. *Free Radical Biology Medicine* 2008; 44 (8): 1493 - 505.

24. Niki E. Lipid peroxidation: Physiological levels and dual biological effects. *Free Radical Biology & Medicine* 2009; 47 (5): 469 - 84.

25. Rafieirad M, Zangenehnezhad Z and Eydipour Z. Effects of oral Ferulago angulate hydroalcoholic extract on lipid peroxidation induced by ischemia hypoperfusion in the hipocamous rat. *International Journal of Biology, Pharmacy and Allied Sciences* 2014; 3 (2): 157 - 64.

26. Paxino G and Wasten C. The rat brain stereotaxic coordinates, 2 nd ed. Acad press Limit. San Diego. 2006, 25 - 8.

27. Sarkaki A, Rafieirad M, Farbood Y, Mansouri S and Moatamedi F. Cognitive deficiency induced by cerebral hypoperfusion/ischemia improves by exercise and grape seed extract. *Health Med. J.* 2012; 6 (4): 1097 - 105.

28. Ohkawa H and Ohishi N.K.Y. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* 1979; 95 (2): 351 - 8.

29. Mansouri MT, Farbood Y, Sameri MJ, Sarkaki A, Naghizadeh B and Rafeirad M. Neuroprotective effects of oral gallic acid against oxidative stress induced by 6hydroxydopamine in rats. *Food Chem.* 2013; 138 (2-3): 1028 - 33.

30. Deumens R, Blokland A and Prickaerts J. Modeling Parkinson's Disease in Rats: An Evaluation of 6-OHDA Lesions of the Nigrostriatal Pathway. *Experimental Neurol.* 2002; 175 (2): 303 - 17.

31. Joseph J and Eduardo T. Parkinson's disease and movement disorders, 4nd ed. Linppicott Williams & Wlikins. Philadelphia. 2002, pp: 116 - 663.

32. Schwarting RK and Huston JP. The unilateral 6-hydroxydopamine lesion model in behavioral brain research: analysis of functional deficits, recovery and treatments. *Prog. Neurobiol.* 1996; 50: 275 - 331.

33. Shukitt-Hale B, Carey A, Simon L and Mark DA. Effects of concord grape juice on cognitive and motor deficits in aging nutrition. *Nutrition* 2006; 22 (3): 295 - 302.

34. Hsiao G, Teng CM, Sheu JR and et al. Cinnamophilin as a novel antiperoxidative cytoprotectant and free radical scavenger. *Biochimica et Biophysica Acta* 2001; 1526: 77 - 88.

35. Dexter DT, Carter CJ and Wells FR. Basal lipid peroxidation in substantia nigra is increased in Parkinson's disease. *J. Neurochem.* 1989; 52: 381 - 9.

36. Hu ML. Measurement of protein thiol groups and glutathione in plasma. *Methods Enzymol.* 1994; 233: 380 - 5.



Sharifi et al.

37. Berlett S and Stadtman E. Oxidation in aging disease and oxidative stress. *Journal of Clinical Chem.* 1997; 272: 2031 - 6.

38. Tiffani Castiglioni E, Saneto RP, Proctor PH and Perez Polo JR. Participation of active oxygen species in 6-hydroxydopamine toxicity to a human neuroblastoma cell line. *Biochemical Pharmacol.* 1982; 31: 181 - 8.

39. Yeh CT, Ching LC and Yen GC. Inducing gene expression of cardiac antioxidant enzymes by dietary phenolic acids in rats. *Journal of Nutritional Biochem.* 2009; 20: 163 - 71.

40. Zhou C, Huang Y and Przedborski S. Oxidative stress in Parkinson's disease: a mechanism of pathogenic and therapeutic significance. *Ann. N. Y. Acad. Sci.* 2008; 1147: 93 - 104.

41. Zare Shahneh F , Valiyari S, Azadmehr A, Hajiaghaee R, Bandehagh A, Baradaran B. Cytotoxic activities of Ferulago angulata extract on human leukemia and lymphoma cells by induction of apoptosis. *Journal of Medicinal Plants Res.* 2013; 7 (11): 677 - 82. **42.** Demetzos C, Perdetzoglou D, Gazouli M

and Tan K. Chemical analysis and

antimicrobial studies on three species of Ferulago from Greece. *Planta Med.* 2000; 66: 560 - 3.

43. Arianpor R, Joghataiy M, Roghani M, Nobakht M and Askari H. Quercetin effects on behavioral and histological changes in experimental models of Parkinson's disease in the adult system nigrastriatal. *J. Iran. Anat. Sci.* 2006; 1: 225 - 39.

44. Salar F, Ziaiy A, Nasri S, Roghani M and Kamalinejad M. Neuronal protective effect of aqueous extract of barberry in rats model of Parkinson's disease. *J. Iran. Anat. Sci.* 2010; 4: 89 - 95.

45. Baluchnejadmojarad T, Roghani M and Mafakheri M. Protective effects of aqueous extract of silybum marianum in 6-hydroxydopamine-induced model of parkinsonism in male rat: a behavioral, biochemical and histological study. *J. Univer. Med.* 2011; 12 (4): 459 - 65.

46. Khanahmadi M and Janfeshan K. Study on antioxidation property of *Ferulago angulata* plant. *Asian J. Plant Sci.* 2006; 5: 521 - 6.

