Evaluation of Antibacterial Activity and Flavonoid Content of Two Capparis Species from Iran

Rahimifard N (Ph.D.)¹, Shojaii A (Ph.D.)^{2*}, Mahbobi M (M.D.)³, Hafezan Gh (M.D.)³, Bagheri F (M.Sc.)⁴, Asgarpanah J (Ph.D.)³

1-Food and Drug Laboratory Research center (FDLRC), Food and Drug Control Laboratories (FDCLs), Ministry of Health (MOH), Pharmaceutical sciences branch, Islamic Azad University, Tehran, Iran

2-Research Institute for Islamic and Complementary Medicine, Iran University of Medical Sciences, Tehran, Iran

3- Pharmacognosy Department, Pharmaceutical Sciences Branch, Islamic Azad University (IAUPS), Tehran, Iran

4- Microbiology Department, Pharmaceutical Sciences Branch, Islamic Azad University (IAUPS), Tehran, Iran

* Corresponding author: Research Institute for Islamic and Complementary Medicine, Iran University of Medical Sciences, Jomhuri Avenue, P.O. Box: 1145847111, Tehran, Iran Tel: +98-21-33950152 (ext 121)

Email: Shojaii.a@iums.ac.ir

Received: 17 March 2015

Accepted: 16 Sep. 2015

Abstract

Background: Due to multiple drug resistance against antimicrobial drugs for treatment of infectious disease, investigations to discovering new antibacterial compounds from natural sources have been designated.

Objective: According to antimicrobial effects of some Capparis species, this study was carried out to evaluate the antimicrobial activity of the extract and fractions of *Capparis cartilaginea* and *Capparis mucronifolia* against 6 bacterial strains.

Methods: Aerial parts of the Capparis species were extracted by maceration method using methanol and fractionated separately by liquid-liquid fractionation method. The antibacterial activity of the extract and fractions were studied against 6 bacterial strains using MIC microplate method. Total flavonoid content (TFC) of the extracts and fractions was determined using AlCl₃ reagent.

Results: The total extract and methanolic fractions of *C. mucronifolia* were the most effective fractions against the bacterial strains. Methanolic fraction of *C. cartilaginea* was the most effective fraction with MIC 10.42 μ g/ml against *Salmonella enterica*. The highest antibacterial activity of *C.mucronifolia* was against *Staphylococcus epidermidis* with MIC 7.8 μ g/ml. Methanolic fractions of Capparis species showed the highest TFC value in AlCl₃ colorimetric assay.

Conclusion: The results of this study indicate that the extracts and fractions of *Capparis* cartilaginea and *Capparis mucronifolia* have antimicrobial effect against 6 gram positive and gram negative strains. According to the high amount of flavonoids in methanolic fraction of *C. mucronifolia*, the antibacterial activity may be related to the flavonoid compounds of this plant.

Keywords: Capparis cartilaginea, Capparis mucronifolia, Antimicrobial activity, Flavonoid, MIC



Evaluation of ...

Introduction

Plants have been used for treatment of infectious disease for many years. Nowadays multiple drug resistance has developed due to the increasingly use of antimicrobial drugs for treatment of infectious disease [1]. Antibiotic resistance is posing an ever increasing therapeutic problem [2]. So, these problems lead to effort to discover new antibacterial compounds from natural source.

Medicinal plants have been used as effective treatments for numerous human diseases like infectious disease in traditional medicines for thousands of years. In recent decades, researchers have paid attention to identify safer herbs and biologically active compounds isolated from plant species [3]. Antibacterial activity of some plant extracts or their active compounds have been reported in different studies [4, 5].

Genus capparis (Capparaceae) consist of bout 150-200 species in tropical and subtropical countries throughout the world [6]. This Genus has four endemic species in Iran [7]. Capparis species have been used as antiinflammatory and analgesic (in gout and arthritis rheumatoid), diuretic, antibacterial and anthelmintic agent in Iranian traditional medicine [8]. Some component including Alkaloids, Terpens, Flavonoid and fatty acids were reported from different Capparis species [9]. These plants showed different biological effects including antioxidant, antibacterial, antihyperglycemic, hypolipidemic and analgesic effects in scientific studies [10-13].

Capparis cartilaginea and *Capparis mucronifolia* are two species which are grown in the southern parts of Iran. According to antibacterial effect of the other capparis species like *C. spinosa*, in this study antibacterial activity of the aerial parts of these plants were studied in some strains of gram



Journal of Medicinal Plants, Volume 14, No. 55, Summer 2015 positive and gram negative bacteria with MIC micro plate method.

Materials and Methods

Plant material

Fresh aerial parts of *C. cartilaginea* and *C. mucronifolia* were collected in May 2013 from Genow protected region in west north of Bandar Abbas and, Ahmadi region of Hadjiabad County respectively, Hormozgan Province, Iran. Specimens were identified by R. Asadpour and vouchers were deposited in the Herbarium of Pharmaceutical Sciences Branch, Islamic Azad University (IAU), Tehran under code number 1562 and 1564 respectively.

Extraction and isolation

2 kg of the aerial parts of plants were dried, grinded and extracted with methanol by maceration. The extraction was repeated for 3 times. Different fractions of the extracts were obtained by liquid-liquid fractionation using water, petroleum ether, chloroform and methanol. The extracts and fractions obtained were concentrated by rotary evaporator and finally dried and stored in a clean, dark container and cool place. The antibacterial activity was evaluated by determining the MIC microplate method.

Test microorganisms

The bacterial Gram-positive bacteria including *Staphylococcus aureus* (ATCC 6538), *Staphylococcus epidermidis* (ATCC 12228) and *Enterococcus faecalis* (ATCC 29212), and Gram-negative bacteria including *Escherichia coli* (ATCC 8739), *Pseudomonas aeruginosa* (ATCC 9027) and *Salmonella enterica* (ATCC 13311) were purchased from Mast diagnostic company as lyophilized ampoule.

Culture of microorganisms:

After recovery of the microorganism in TSB (Triptycase soy broth) for 24-48 h at 37 °C, the overnight microorganisms which be used for antimicrobial testing, obtained by subculture on plates containing TSA medium and incubation for 24 h at 37 °C.

Minimum inhibitory concentration (MIC) using microplate:

Micro-dilution assay was used to determine MIC of the extracts against 6 bacterial strains. MIC determined was by testing 8 concentrations of the extract against grampositive and Gram-negative bacteria, by microplate serial dilution method and using Mueller Hinton broth. The reconstituted extract was diluted to give concentrations of 1000, 500, 250, 125, 62.5, 31.25, 15.625, 7.812 µg/ml. The lowest concentration of the extract that could inhibit the bacterial growth was considered as MIC [14]. Chephalexin and gentamicin were used as positive controls.

Total flavonoid content:

The total flavonoid content (TFC) of С. cartilaginea С. mucronifolia and methanolic extracts and their fractions was determined using AlCl₃ reagent [15]. Briefly, 2.5 mL of each sample (and/or guercetin as the standard), previously dissolved in 90% ethanol, was mixed with 2.5 mL of a 2% AlCl₃ solution in 90% ethanol. After 40 min, the absorbance of the yellow color produced was measured at 425 nm. The TFC [as µg quercetin equivalents / mg of sample] for the sample was calculated on the basis of a linear calibration curve obtained using quercetin $(y=0.0169x+0.3526, r^2=0.995).$

Results

TFC of the Capparis extracts and fractions are presented in Table 1. Extract and fractions of C. mucronifolia showed higher amounts of TFC rather than *C. cartilaginea*.

The inhibitory effects of the methanolic extracts and fractions of C. cartilaginea and mucronifolia against different С. microorganisms are presented in table 2 and table 3 respectively. The extract and fractions of these plant showed different antibacterial activity against gram-positive and gram negative bacteria in microplate method. The total extract and methanolic fraction of C. mucronifolia were the most effective fractions of this plant against bacterial strains. The most effective fraction of C. cartilaginea was methanolic fraction with MIC 10.42 µg/ml against S. enterica. The most effectiveness of C. mucronifolia was against S. epidermidis with MIC 7.8 µg/ml for total extract and methanolic fraction.

Table 1- TFC of the Capparis extracts and fractions (µg quercetin/mg extract)							
Capparis species	Total extract	Petrolium	<i>ium Aqueous</i> Methanolic		Chloroformic		
		ether fraction	fraction	fraction	fraction		
C. mucronifolia	36.54±0.5	65.8±0.3	69.13±5.4	76.1±1.74	-		
C. cartilaginea	16.72±0.71	16.64±4.2	17.87±1.8	33.77±0.76	38.61±1.4		

Table 1- TFC of the Capparis extracts and fraction	s (μg quercetin/mg extract)

*Data are mean \pm SD, n=3



Microorganism	Total extract	Petrolium ether fraction	<i>Aqueous</i> fraction	<i>Methanolic</i> fraction	<i>Chloroformic</i> fraction	Chephalexin	Gentamicin
Pseudomonas aeruginosa	52.08	62.5	15.63	31.25	52.08		2.5
Esherichia coli	62.5	62.5	31.25	41.67	52.08		5
Salmonella enterica	31.25	15.63	31.25	10.42	13.02		23.4
Enterococcus faecalis	15.63	52.08	31.25	62.5	31.25	16.6	
Staphylococcus aureus	13.69	31.25	13.02	31.25	31.25	2.6	
Staphylococcus epidermidis	62.5	26.04	26.04	31.25	62.5	5.2	

Table 2- MIC	values of ext	act and fraction	ons of C. car	<i>tilaginea</i> (µg/ml)
--------------	---------------	------------------	---------------	--------------------------

• Data are mean of three replicate

					,	
Microorganism	Total extract	Petrolium ether fraction	Aqueous fraction	<i>Methanolic</i> fraction	Chephalexin	Gentamicin
Pseudomonas aeruginosa	>1000*	>1000	>1000	>1000		2.5
Esherichia coli	31.25	250	187.5	46.87		5
Salmonella enterica	46.87	250	125	62.5		23.4
Enterococcus faecalis	250	250	250	250	16.6	
Staphylococcus aureus	7.8	250	46.87	11.7	2.6	
Staphylococcus epidermidis	7.8	250	46.87	7.8	5.2	

Table 3- MIC values of extract and fractions of C. mucronifolia (µg/ml)

• * > 1000= extract or fraction was inactive,

• Data are mean of three replicate.

Discussion

Antibiotic resistance is posing an ever increasing therapeutic problem and these problems lead to effort to discover new antibacterial compounds from natural source [2].

C. cartilaginea and *C. mucronifolia* are two species of capparis genus which are endemic in the south of Iran. Some species of capparis



(like *C. spinosa*) showed significant antibacterial activity [12], so in this study antibacterial effect of the extract and fractions of aerial parts of *C. cartilaginea* and *C. mucronifolia* was determined by micro plate MIC method against 6 bacterial strains. The most effective fraction of *C. cartilaginea* was methanolic fraction against *S. enterica* and total extract and methanolic fraction of *C. mucrpnifolia* showed the best activity against *St. epidermidis*.

Previous antimicrobial study on C. cartilaginea showed MIC> 1000 μ g/ml for bacterial strains including Staphylococcus aureus, Bacillus subtilis and Microccocus flavus [16]. This studv showed the antibacterial effect of the extract and different fractions of C. cartilaginea against 6 bacterial strains using MIC method. Also, this study is a first report on the antibacterial activity of C. mucronifolia against gram positive and gram negative bacterial strains. The tested capparis species showed lower antibacterial activities in comparison to C. spinosa, against strains Phytochemical bacterial [17]. investigations of Capparis species showed the presence of flavonoids [18, 19]. In this study, TFC of the extracts and fractions of two Capparis species was determined. The findings showed higher TFC value in C. mucronifolia extract and fractions. The methanolic fractions

References

1. Davis J. Inactivation of the antibiotics and the dissemination of resistance genes. *Sci.* 1994; 264: 375 - 82.

2. Kim H, Park SW, Park JM, Moon KH and Lee CK. Screening and isolation of antibiotic resistance inhibitors from herb material Resistant Inhibition of 21 Korean plants. *Nat. Prod. Sci.* 1995; 1: 50 - 4.

3. Sen A and Batra A. Evaluation of Antimicrobial activity of different solvent extracts of medicinal plant: *Mellia Azedarach* L. *IJCPR*. 2012; 4 (2): 67 - 73.

4. Shahtalebi MA, Ghanadian M, Farzan A, Shiri N, Shokri D and Fatemi SA. Deodorant

of two Capparis species showed high amounts of total flavonoids in AlCl₃ colorimetric assay. Since the methanolic fraction of *C. mucronifolia* showed the best antibacterial activity, flavonoids may be responsible for antibacterial activity of this plant.

Conclusion

The results of this study indicate that the extracts and fractions of *C. cartilaginea* and *C. mucronifolia* have antimicrobial effect and total extract and methanolic fraction of *C. mucronifolia* showed the best antibacterial activity. Certainly further investigations are necessary to determine the active principles which responsible for antibacterial activity.

Acknowledgment

We thanks from Mrs Abdollahi and Mrs Abdoli (Research institute for Islamic and complementary medicine) for assistance in colorimetric assays.

effects of a sage extract stick: antibacterial activity and sensory evaluation of axillary deodorancy. J. Res. Med. Sci. 2013; 18: 833 - 9. 5. Tamokou JD, Kuiate JR, Tene M, Kenla Nwemeguela and Tane TJ Р. The antimicrobial activities of extract and compounds isolated from Brillantaisia lamium. Iran. J. Med. Sci. 2011; 36 (1): 24 -31.

6. Trombetta D, Occhiuto F, Perri D, Puglia C, Santagati N A and de Pasquale A. Antiallergic and antihistaminic effect of two extracts of *Capparis spinosa* L. flowering buds. *Phytother Res.* 2005; 19: 29 - 33.



Evaluation of ...

7. Mozafarian V. Dictionary of Iranian plants, Moaser Publication. 1375, p. 98

8. Aghili Khorasani H. Makhzan-Al-Advieh. Research institute for Islamic and complementary medicine Press 2008, Iran University of medical sciences. pp: 729 - 30.

9. Yang T, Wanga CH, Chou GX, Wua T, Cheng XM and Wang ZT. New alkaloids from *Capparis spinosa*: Structure and X-ray crystallographic analysis. *Food Chem.* 2010; 123: 705 - 10.

10. Eddouks M., Lemhadri A and Michel J B. Hypolipidemic activity of aqueous extract of *Capparis spinosa* L. in normal and diabetic rats. *J of Ethnopharmacol.* 2005; 98: 345 - 50.

11. Eddouks M, Lemhadri A and Michel JB. Caraway and caper: Potential antihyperglycaemic plants in diabetic rats. *J. Ethnopharmacol.* 2004; 94: 143 - 8.

12. Yang T, Liu YQ, Wang CH and Wang Z T. Advances on investigation of chemical constituents, pharmacological activities and clinical applications of *Capparis spinosa*. *Chine. J. Chin. Mat. Med.* 2008; 33: 2453 - 8.

13. Ghule BV, Murugananthan GB and Yeole PG. Analgesic and antipyretic effects of Capparis zeylanica leaves. *Fitoterapia* 2007; 78: 365 - 69.

14. Hakemi Vala M, Asgarpanah J, Hedayati MH, Shirali J and Bagheri Bejestani F. Antibacterial and cytotoxic activity of *Eremurus persicus* (Jaub and Spach) Boiss. *AJMR*. 2011; 5: 2349 - 52.

15. Nickavar B, Kamalinejad M, Haj-Yahya M and Shafaghi B. Comparison of the Free Radical Scavenging Activity of Six Iranian Achillea Species, *Pharm. Biol.* 2006; 44 (3): 208 - 212.

16. Mathana RA, Lindequist U, Gruenert R and Bednarski PJ. Studies of invitro anticancer, antimicrobial and antioxidant potentials of selected Yemeni medicinal plants from the island Soqotra. *BMC Complement*. *Altern. Med.* 2009; 9 (7): 1 - 11.

17. Mahboubi M and Maboubi A. Antimicrobial activity of *Capparis spinosa* as its usages in traditional medicine. *Herba Polonica* 2014; 60 (1): 39 - 48.

18. Ferheen S, Rasool MA, Imran M, Fareed G, Afza N and Malik A. Capspinosin, a new flavonoid from Capparis spinosa. *JCSP* 2013; 35: 984 - 86.

19. Sharaf M, El-Ansari MA, Saleh NAM. Quercetin triglycoside from *Capparis spinosa*. *Fitoterapia* 2000; 71: 46 - 9.

