

***Ex vivo*-Expansion of Cord Blood Cells and Its Clinical Application**

Kamran Alimoghaddam, Ph.D.[‡], Mandana Mohyedin, Ph.D., Sina Vatandoust, M.D.,
Zahra Goliaei, M.D., Poya Alijanipour, M.D., Farima Forouzia, M.D.

Hematology Department, Oncology and BMT Research Center,
Tehran University of Medical Sciences

[‡] Corresponding Address: P.O.Box: 14114, Hematology-Oncology and BMT Research Center,
Tehran University of Medical Sciences, Tehran, Iran
Email: alimg@ams.ac.ir

Received: 2/Jul/2006

Cord blood is a rich source of hematopoietic stem cells and could be potentially used for transplantation instead of conventional sources of stem cells (bone marrow or peripheral blood).

Cord blood cells are successfully used in pediatric and adult patients, but their major limitation is the low number of hematopoietic stem cells for patients of large body size.

There are several possible solutions for this problem, including use of third party donor, use of multi-unit cord blood, and finally *ex-vivo* expansion of cord blood hematopoietic stem cells to accelerate engraftment of transplanted cells. In this article we will discuss *ex-vivo* expansion from bench and clinical points of view.

Keywords: Cord Blood, Expansion, Transplantation

Introduction

Cord blood is a potential source for hematopoietic transplantation. The most important limitation of its use is the low number of hematopoietic stem cells (HSCs) in cord blood samples that might prevent a successful engraftment or cause delayed engraftment and higher morbidity and mortality during pancytopenic period after the transplantation. One of the possible theoretical methods to circumvent this unacceptable mortality rate is *ex vivo* expansion of hematopoietic cells to increase the number of stem cells and then use of this *ex vivo* expanded HSCs as a source of transplantation.

Biology of human umbilical cord blood hematopoietic stem cells

Cord blood stem cells are identified by their immunophenotypic and functional characteristics.

Immunophenotypic characterization of cord blood HSCs:

Certain immunophenotypic markers help define primitive cell populations, such as: CD34, CD38, thy-1, c-kit, HLA-DR, Rhodamine 123, etc.

CD34 antigen, an integral membrane glycoprotein, is a defining hallmark of HSCs. It has been suggested that this molecule

functions as a regulator of hematopoietic environment (1). In cord blood, the CD34+ cell content is about 1% of nucleated cells, which is similar to the bone marrow content of 1-3% (2). However, the frequency of CD34+ cells is much higher, up to 11%, at earlier stages of gestation, although this decreases with age of gestation (3).

One of the most frequently used markers is the CD38 antigen, which is absent on the more primitive CD34+ cells (4). The subset of CD34+CD38- cells in cord blood is fourfold higher than in adult bone marrow (5).

Another cell surface marker commonly used in immunophenotyping is HLA-DR. Primitive CD34+ cells in the bone marrow are DR-, while CD34+ cells from cord blood with similar functional properties express HLA-DR (6).

The function of Thy-1 on HSCs is unknown. It might be involved in HSCs development by mediating signals inhibiting the proliferation of primitive cells. Baum and colleagues demonstrated that only CD34+ cells expressing the Thy-1 antigen reconstituted human hematopoiesis in SCID mice (7).

The c-kit proto-oncogene encodes a transmembrane receptor with tyrosine kinase activity and is intimately involved in hematopoiesis (8). The ligand for this receptor is steel factor or SCF. The c-kit antigen is

expressed by 60% of CD34+ cord blood cells (9).

Expression of lymphoid and myeloid associated antigens on umbilical cord blood CD34+ cells has also been documented. Saeland *et al.* found that in contrast to adult bone marrow in which 25% of the CD34+ cells express CD10 and 18% express CD19, these markers are rarely expressed in CD34+ cell population derived from cord blood (10).

The vast majority of umbilical cord blood CD34+ cells (90%) co-express Flt3 (CD135). It is the receptor for early acting cytokine Flt3 ligand (11).

Several cell adhesion molecules are present on umbilical cord blood CD34+ cells, as is observed in bone marrow (BM) CD34+ cells. CD44 and LAM-1 adhesion receptors involved in the homing of hematopoietic cells have been found to be strongly expressed on umbilical cord blood CD34+ cells (10).

Functional characterization of cord blood HSCs.

Different methods are used to assay function of HSCs, including: colony forming cell (CFC) assay, long term culture – initiating cell (LTC-IC) assay, and SCID repopulating cell (SRC) assay.

Published data indicate that there are about 13 to 2400 GM-CFC, 8000 BFU-E, and between 1 and 10000 CFU-GEMM per ml of cord blood. (12). CFC proportion is higher in cord blood compared to BM, and is even higher in cord blood samples from pre-term infants (13).

Pattergell *et al.* compared the results of LTC-IC in BM, cord blood, and mobilized peripheral blood. They showed that mononuclear cells (MNCs) of peripheral blood produce more CFCs than cord blood and BM (14). However, others found significantly more progenitor cells in LTC-IC assay and longer cell production in cord blood MNCs compared with BM ones (15).

Vormoor *et al.* has reported stable human hematopoiesis in SCID mice transplanted with human cord blood cells. It has been suggested that the human cells capable of repopulation in such animals represent more primitive properties than LTC-ICs (16). Studies show that the frequency of SCID repopulating cells (SRCs) in cord blood is three folds higher than BM and six folds higher than peripheral blood (17).

Immunological characterization of cord blood HSCs.

There is less graft versus host disease (GVHD) with cord blood compared to BM transplantation, and we can transplant cord bloods with more HLA mismatches (18).

Therefore, cord blood may generate a lower immunological response compared to adult cell transplantation. T lymphocytes and NK cells are the most important cells in this response. Cord blood T cells have significantly less ability than adult T cell to produce IL-2 and express functional IL-2 receptor complexes. Moreover, the potential of cord blood cells to produce helper T cells derived cytokines (INF γ and IL-2) is lower, presumably due to the presence of immature, naive T cells in cord blood (19).

NK cells have been implicated in mediating the graft versus leukemia (GVL) effect. While cord blood generally manifests low NK activity compared to adult BM and blood, cord blood NK activity is readily augmented *in vitro* by cytokines such as IL-2 and IL-12, suggesting that cord blood NK cells should be as effective as adult BM or blood NK cells to mediate a GVL effect (20).

Cell cycle status of cord blood HSCs. Practically, all CD34+CD38- cells in cord blood are in G0/G1 status (21). These cells would present a useful target for retroviral transfection, as they respond to cytokine stimulation and rapidly enter S phase of the cycle. The proportion of primitive CD34+ in G0/G1 decrease from 98% to 55% after 48 hours of exposure to certain cytokines or unknown factors in cord blood plasma (22).

Important factors in cord blood expansion

1. Stromal cells:

Stroma-dependent culture systems have been developed to study long term hematopoiesis *in vitro*. Stromal cells are a source of growth factors and adhesion molecules to support stem cells (23). Cord blood mononuclear cells can not form long term colonies in culture without supportive environment (15). Some murine cell lines have been found to be useful for stroma dependent cultures of HSCs.

Kawada and colleagues (24) used a murine cell line for expansion of cord blood HSCs. They showed that direct adhesion of stromal cells to human progenitors significantly increased the number of CD34+CD38- cells. When cell lines were physically separated from human progenitor cells, they failed to get good results. It is suggested that stromal cells can support proliferation of HSCs by direct cell-cell interaction. It seems that stromal support improve expansion of cord blood hematopoietic stem cells (25).

2. Effect of serum on cord blood expansion:

Unknown factors present in cord blood plasma, probably capable of crossing the placenta, affect both by themselves and in the presence

of growth factors, hematopoietic progenitor cell proliferation and differentiation at all stages of hematopoietic development. Moreover, co-culturing the cord blood cells with mesenchymal cells may increase expansion and also CXCR-4 expression on *ex vivo* expanded cord blood cells (26).

In the absence of growth factors, CFU-GM expansion didn't occur in cord blood cell cultures supplemented with fetal calf serum or peripheral blood plasma (27). Broxmeyer *et al.* noted that cord blood plasma, but not fetal calf serum or peripheral blood plasma, increased both the size of secondary, replated CFU-GEMM and the number of times that CFU-GEMM could be replated to form colonies in culture containing SCF and erythropoietin (EPO) (28). As combining IL-1, IL-3, IL-6, IL-11, G-CSF and GM-CSF couldn't reproduce this effect, it is possible that the cord blood plasma effect may be attributable to a novel growth factor with synergy for SCF and EPO (28). IL-6, Flt3 ligand, and thrombopoietin (TPO) are the possible candidates (29).

Although experiments showed that expansion could be improved using serum, but to transfer expansion to the clinic, good manufacturing practice (GMP) standards are required. 235-fold expansion of cord blood CD34+ cells was obtained with a cocktail containing flt3 ligand, TPO, IL-6, and IL-11 at 5 weeks with serum-free medium [30]. Flt-3 ligand, SCF, and TPO are considered as early acting and indispensable cytokines. By adding IL-3 for these cytokines in serum free culture, the amplification of CD34+ cord blood cells was increased 20.9 fold as opposed to 9.3 fold without IL-3 after 7 days (31).

CD34+CD38- surface phenotype has been used to measure primitive cell numbers. One UK study demonstrated a lack of expression of CD38 on CD34+ cells in serum free cultures. This finding must be considered in *ex-vivo* cord blood expansion in serum free conditions, and CD34+CD38- phenotype should not be used to confirm the presence of primitive progenitor cells (32).

3. Purification:

Cells belonging to the stem cell compartment are rare, representing about 1 in 10,000 mononuclear cells in the adult bone marrow. Purifying the stem cell compartment becomes progressively easier as markers become available to select these cells positively (CD34+Thy-1+) or negatively (CD38-DR-). CD34+ selected cell fraction has been used most frequently as the starting cell population for the expansion of hematopoietic progenitor cells.

CD34+ selection is a time-consuming and expensive process, leading to approximately 50% loss of CD34+ cells (33). In the static culture, the increase in total number of cord blood mononuclear cells is less than CD34+ selected sample. However the CD34+ proportion increases significantly in the mononuclear cell culture and decreases in the CD34+ cell culture. This, in part, compensates the limited increase of cell number in the mononuclear cell culture. [34] In addition, cultures initiated with mononuclear cells have nearly the same performance as CD34+ cells in perfusion cultures (35).

4. Cytokines:

1) Stimulatory Cytokines

Hematopoietic cell proliferation and differentiation is regulated by stimulatory and inhibitory signals that are mediated by cytokines. Cytokines are secretory proteins produced by a variety of hematopoietic and non-hematopoietic cells. Each cytokine alone has modest effect on hematopoietic cell amplification, but they show additive or synergistic effect in combination with other growth factors (36, 37, 38, 39).

Moreover, the effect of cytokines in a culture system is not only a function of their concentration and synergistic interaction, but also other factors like culture conditions (40) and target cells. Cord blood progenitors and/or their progeny can produce cytokines including GM-CSF and IL-3, in culture, leading to autocrine or paracrine stimulation of cells (41).

Stem Cell Factor (SCF) is a potent hematopoietic growth factor [36] produced by bone marrow stromal cells and fibroblasts. It interacts with a variety of other growth factors to influence very early hematopoietic stem cells (36, 42). The c-kit acts as its receptor.

Flt3-Ligand (FL) provides a significant amplification of both committed and early progenitors (43). Flt3-L has been shown to act directly on quiescent cells causing them to enter the cycle (44). In addition, it has been demonstrated to induce the proliferation of CD34+CD38- bone marrow and cord blood cells that are non-responsive to other early acting cytokines (45, 46, 47).

Granulocyte-Colony Stimulating Factor (G-CSF) and **Granulocyte-Macrophage Colony Stimulating Factor (GM-CSF)** are produced by a variety of cell types including fibroblasts, endothelial cells, macrophages, and T lymphocytes. G-CSF can act on primitive as well as later hematopoietic progenitors, whereas the action of GM-CSF may be restricted to terminally differentiating cells (48).

Thrombopoietin (TPO) is an early acting cytokine (49). TPO stimulates and supports survival of primitive stem cells. It also regulates megakaryocyte proliferation and differentiation (49, 50, 51, 52). The C-mpl acts as its receptor.

Erythropoietin (EPO) secreted by renal tubular cells has limited proliferative effects; it modulates/stimulates survival and terminal maturation of erythroid progenitors. EPO also increases red blood cell production.

Interleukin-3 (IL-3) is a multilineage stimulator with direct megakaryocyte, mast cell/basophil, B cell, and eosinophil stimulatory activity. There is some evidence that IL-3 increases asymmetric division of the stem cells, leading to stem cell differentiation (53).

There is some evidence that **Interleukin-6 (IL-6)** dramatically stimulates expansion of human hematopoietic progenitor cells *in vitro* in the presence of SCF [54]. IL-6 is active on more immature hematopoietic progenitors (50). Its

Interleukin-11 (IL-11) is produced by bone marrow stromal cells and induces megakaryocyte colony forming and maturation. Although different combinations of cytokines have been used, standard expansion protocols have not yet been established due to complexity of cytokine interactions.

Different studies have shown that maximal expansion of hematopoietic cells is generally achieved using one or more cytokines acting on primitive cells in combination with cytokines acting on less-primitive cells (53). In addition, the cytokines acting through separate signaling pathways show more synergism (i.e., IL-6 in combination with SCF) (54). Some of these combinations are presented in Table 1.

II) Inhibitory Cytokines

Macrophage inflammatory protein-1 alpha (MIP-1 alpha) is secreted by monocyte/macrophage and T cells. It has been shown that MIP-1 α prevents *in vivo* murine stem cells from entering cell cycle (62).

Table-1: Effects of cytokines on *ex vivo* expansion of human stem/progenitor cells (53).

Reference	Cells	Culture	Cytokines	Duration	Expansion time		
					TNC	CFC	LTC-IC
Piacibello (55)	CD34+	IMDM/ 10%FCS	FL+TPO	<30 weeks		28 millions	270000
Dening-Kendall (56)	Cd34+	IMDM/ 20%FCS	SCF+ IL-3+ IL-6+ GM-CSF+ G-CSF	14 days	2500	CFU-GM: 49 BFU-E: 1	2.5
Kogler (57)	Cd34+	Serum- Free	FL+ SCF+ IL-3	7 days	138	CFU-GM: 264 CFU-GEMM: 94 BFU-E: 126	6.7
Piacibello (50)	CD34+	IMDM/ 10%FCS	SCF+ IL-3	8 weeks	12000	36	
			SCF+ FL	12-14weeks	<20000	<300	
Ohmizono (38)	CD34+	Serum- Free	SCF+ IL-3	14 days		BFU-E: 50 CFU-GM: 30 CFU-GEMM: 30	
Traycoff (58)	CD34+	Serum- Free	SCF+ IL-3	5-7 days			Reduced
Ruggieri (59)	CD34+	McCoys/ 10%Cord Serum	SCF+ IL-3 +GM-CSF	7 days	64	11	
Moore (60)	CD34+	Not stated	SCF+ IL-3 +IL-1 + EPO	14 days	2800	600	18
Durand (61)	CD34+	Serum- Free	SCF+ IL-3	21 days	1000	5000	
Migliaccio (42)	CD34+ SBA-	Serum- Free	SCF+ IL-3	3 weeks		20	
			SCF+ G-CSF			10	
			SCF+EPO			2-3	

IMDM: Iscov's modified Dulbecco's medium; FCS: fetal calf serum; FL: Flt3 ligand; TPO: thrombopoietin; SCF: stem cell factor; IL: interleukin; GM-CSF: granulocyte-macrophage colony-stimulating factor; G-CSF: granulocyte colony-stimulating factor; EPO: erythropoietin; TNC: total nucleated cells; CFC: colony forming cells; LTC-IC: long-term culture-initiating cells; CFU-GM: colony forming unit granulocyte-macrophage; BFU-E: burst-forming unit Erythroid; CFU-GEMM: colony forming unit granulocyte-erythroid-macrophage-megakaryocyte.

Some studies showed that MIP-1 α suppressed the growth of immature hematopoietic progenitors. Nevertheless, recent studies suggest that it has also a stimulatory effect on proliferation of more mature hematopoietic progenitors (63, 64, 65, 66). MIP-1 α may have a role in maintaining the long-term culture initiating cells (LTC-IC) (67). There is also some evidence that MIP-1 α inhibits bone marrow granulocyte-macrophage-colony forming cells (GM-CFC), while stimulating GM-CFC from cord blood CD34⁺ cells over the same dose range (68, 54, 93).

Transforming growth factor-beta (TGF-beta) is secreted by most cell types and affects most cell types. The major biological effect of TGF-beta on hematopoietic cell growth is the reversible inhibition of entry into the cell cycle (69). Many studies have shown the inhibitory activities of TGF-beta on hematopoiesis (70, 73), but recent evidence supports that TGF-beta can have both inhibitory and stimulatory effects, depending on the differentiation state of the target cell and other cytokines interacting with the cell (71, 74). There is also some evidence that the inhibitory effect of FCS and human serum on progenitor cell proliferation is caused by TGF-beta (71, 72).

Leukemia inhibitory factor (LIF) is a glycoprotein affecting a wide spectrum of cells (75). LIF prevents differentiation commitment of normal embryonic stem cells (75, 76). In human, it stimulates IL-3-dependent growth of primitive HSCs (77, 78). LIF induces *in vitro* proliferation of primitive HSCs and appears to be required for the survival of HSCs *in vivo* (79).

Since using a combination of just stimulatory cytokines leads to production of more mature cells rather than primitive stem cells, it seems that an ideal combination of cytokines should include both stimulatory and inhibitory cytokines.

Bioreactor systems

Recently, alternative culture techniques such as bioreactor systems have been developed to maintain growth factors and other required elements such as oxygen as well as waste products in constant and characteristic concentrations. This stirred culture system is particularly required when large-scale expansion culture is planned, especially for clinical purposes. By providing a closed system, this method protects the culture medium from infectious agents, which always threaten the conventional cultures especially during refeeding and other manipulations. Some studies have reported significant

improvement in cord blood expansion in bioreactor perfusion culture system, compared with static cultures (80, 81, 82).

Cord blood cryopreservation and expansion

The use of cord blood for transplantation would be much facilitated by banking of cord blood samples. Since cryopreservation remains the method of choice for long-term preservation of progenitor/stem cells, cord blood cryopreservation has become an important issue in banking and transplantation. It seems that the most suitable cryopreservation techniques used for cord blood samples are almost similar to those of bone marrow or peripheral blood progenitor/stem cell cryopreservation (83, 84).

Although several studies have shown that cryopreservation does not significantly reduce expansion potential (85), colonogenicity, and immunophenotypic properties of cord blood progenitor/stem cells, most of these studies have focused on quantitative measurements and they have not assessed the quality of recovered progenitor/stem cells after cryopreservation (86).

Rice *et al.* found that cryopreservation of *ex vivo* expanded cord blood cells doesn't deteriorate engraftment measures in NOD-SCID mouse model (87). This study raises the question whether expansion of cord blood cells should be done prior to or after cryopreservation to obtain the best clinical results.

Clinical studies of transplantation by ex-vivo expanded cord blood cells

Animal models demonstrate that engraftment of *ex vivo* expanded cord blood cells is possible, but it is delayed (88). Recent data demonstrated that *ex vivo* expanded cord blood cells may be useful in transplantation in adults. These expanded cells resulted in faster neutrophil engraftment (89, 90, 91, 92). Theoretically, transplantation of *ex vivo* expanded cord blood cells at the same time of non-expanded cord blood cells from the same donor could increase the number of immediately available progenitor cells responsible for short term engraftment.

Kogler *et al.* showed that expansion of 1/8 of sibling cord blood in the presence of G-CSF, TPO, and flt3-L and then simultaneous transplantation of a high risk leukemic patient using both expanded and non-expanded cells resulted in rapid and durable neutrophil engraftment (91). Pecora *et al.* evaluated the effect of supplementing unrelated umbilical cord blood with *ex vivo* expanded umbilical cord blood cells from the same donor in two

older adult patients with high risk CML and no alternative donor (92). They used a clinical grade perfusion system (Astrom Replicell) and observed that *ex vivo* expanded cells facilitate hematopoietic recovery.

References

1. Healy L, May G, Gale K, Grosveld F, Greaves M, Enver T. The stem cell antigen CD 34 functions as a regulator of hematopoietic cell adhesion. *Proc Natl Acad Sci USA*. 1995; 92: 12240-12244
2. Sutherland DR, Keating A, Nayar R, Anania S, Stewart AK. Sensitive detection and enumeration of CD34+ cells in peripheral and cord blood by flowcytometry. *Exp Haematol*. 1994; 22: 1003-1010
3. Thilaganathan B, Nicolaidis KH, Morgan G. Subpopulation of CD34 positive haemopoietic progenitors in fetal blood. *Bjh*. 1994; 87: 634-636
4. Sutherland H, Evaes CJ, Dragowska W, Lansorp PM. Characterisation and partial purification of human marrow cells capable of initiating of long-term hematopoiesis *in vitro*. *Blood*. 1989; 74: 1563-1569
5. Broxmeyer HE, Douglas GW, Hangoc G, Cooper S, Bard J, English D, Arrny M, Thomas L, Boyse EA. Human umbilical cord blood as a potential source of transplantable hematopoietic stem/progenitor cells. *Proc Natl Acad Sci USA*. 1989; 86: 3828-3832
6. Traycoff CM, Abboud MR, Laver J, Brandt JE, Hoffman R, Law P, Ishizawa L, Srour EF. Evaluation of the *in vitro* behavior of phenotypically defined populations of umbilical cord blood haematopoietic progenitor cells. *Exp Haematol* 1994; 22: 215-222
7. Baum CM, Weissman IL, Tsukamoto AS, Buckle AM, Peault B. Isolation of a candidate human hematopoietic stem cell population. *Proc Natl Acad Sci USA* 1992; 89: 2804-2808
8. Gunji Y, Nakamura M, Osawa H, Nagayoshi K, Nakauchi H, Miura Y, Yanagisawa M, Suda T. Human primitive hematopoietic progenitor cells are more enriched in Kit low cells than Kit high cells. *Blood* 1993; 82: 3282-3289
9. Reisbach G, Bartke I, Kempkes B, Kostka G, Ellwart J, Birner A, Thalmeier K, Mailhammer R, Bornkamm GW, Ullrich A. Characterization of hemopoietic cell populations from human cord blood expressing c-kit. *Exp Hematol* 1993; 21: 74-79
10. Saeland S, Duvert V, Caux C, Pandrau D, Favre C, Valle A, Durand I, Charbord P, de Vries J, Banchereau J. Distribution of surface membrane molecules on bone marrow and cord blood CD34+ hematopoietic cells. *Exp Hematol* 1992; 20:24-33
11. Rappold I, Ziegler BL, Kohler I, Marchetto S, Rosnet O, Birnbaum D, Simmons PJ, Zannettino AC, Hill B, Neu S, Knapp W, Alitalo R, Alitalo K, Ullrich A, Kanz L, Buhning HJ. Functional and phenotypic characterization of cord blood and bone marrow subset expressing flt3 (CD135) receptor tyrosin kinase. *Blood* 1997; 90: 111-125
12. Traycoff CM, Abboud MR, Laver J, Brandt JE, Hoffman R, Law P, Ishizawa L, Srour EF. Evaluation of the *in vitro* behavior of phenotypically defined populations of umbilical cord blood hematopoietic progenitor cells. *Exp Hematol*. 1994; 22(2): 215-22
13. Mayani H, Lansdorp PM. Thy-1 expression is linked to functional properties of primitive hematopoietic progenitor cells from human umbilical cord blood. *Blood* 1994; 83: 2410-4217
14. Pattengell R, Luft T, Henschler R, Hows JM, Dexter TM, Ryder D, Testa NG. Direct comparison by limiting dilution analysis of long term culture initiating cell in human bone marrow, umbilical cord blood and blood stem cell. *Blood* 1994;84:3653-3659.
15. Hows JM, Bradley BA, Marsh JC, Luft T, Coutinho L, Testa NG, Dexter TM. Growth of human umbilical cord blood in long term hematopoietic cultures. *Lancet* 1992; 340: 73-76
16. Vormoor J, Lapidot T, Pflumio F, Risdon G, Patterson B, Broxmeyer HE, Dick JE. Immature human cord blood progenitors engraft and proliferate to high levels in sever combined immunodeficient mice. *Blood* 1994; 83: 2489-2497
17. Wong JC, Doendens M, Dick JE. Primitive human hematopoietic cells are enriched in cord blood compared with adult bone marrow or mobilized peripheral blood as measured by the quantitative *in vivo* SCID- repopulating cell assay. *Blood* 1997; 89: 3919-3924
18. Cohen SB, Madrigal JA. Immunological and functional differences between cord and peripheral blood. *BMT* 1998; 21: 9-12
19. Konoplev S, Medeiros LJ, Bueso-Ramos CE, Jorgensen JL, Lin P. Immunophenotypic and functional characterization of human umbilical cord blood mononuclear cells. *Leukemia* 1999; 13: 87-89
20. Gaddy J. Clinical and basic science studies of human umbilical cord blood: implication for the graft versus leukemia effect following cord blood transplantation. in: Barret J, Jang Z eds. *Adoptive alloimmunotherapy- graft versus leukemia*. 1998
21. Hao QL, Thiemann FT, Petersen D, Smogorzewska EM, Crooks GM. Extended long-term culture reveals a highly quiescent and primitive human hematopoietic progenitor population. *Blood* 1996; 88: 3306-3313
22. Traycoff CM, Abboud MR, Laver J, Clapp DW, Srour EF. Rapid exit from G0/G1 phases of cell cycle in response to stem cell factor confers on umbilical cord blood CD34+ cells, an enhanced *ex vivo* expansion potential. *Exp Hematol* 1994; 22: 1264-1272
23. Wineman JP, Nishikawa S, Muller-Sieburg CE. Maintenance of high levels of pluripotent hematopoietic stem cells *in vitro*: effect of stromal cells and C-kit. *Blood* 1993; 81: 365-372
24. Kawada H, Ando K, Tsuji T, Shimakura Y, Nakamura Y, Chargui J, Hagihara M, Itagaki H, Shimizu T, Inokuchi S, Kato S, Hotta T. Rapid *ex vivo* expansion of human umbilical cord hematopoietic progenitors using a novel culture system. *Exp hematol*. 1993; 27: 904-915
25. Soleimani M, Mozdarani H, Pourfathollah AA, Mortazavi Y, Alimoghadam K, Mortazavi Y, Alimoghadam K, Nikogoftar M, Zonobi Z, Hajifathollahi A. A co-culture system for expansion nonenriched cord blood Stem/Progenitor cells. *Biotechnology* 2005; 4: 310-315
26. Soleimani M, Mozdarani H, Pourfathollah AA, Mortazavi Y, Alimoghadam K, Hajifathollahi A, Zonob: Increase of CXCR4 expression on expanded non-

- enriched cord blood CD34+ cells using MSCs. *Yakhteh*. 2005; 7; 74-79
27. Carow CE, Hangoc G, Broxmeyer HE. Human multipotential progenitor cells (CFU-GEMM) have extensive replicating capacity for second CFU-GEMM- an effect enhanced by cord blood plasma. *Blood* 1993; 81: 942-949
28. Ruggieri L, Heimfeld S, Broxmeyer HE. Cytokine-dependent *ex vivo* expansion of early subsets of CD34+ cord blood myeloid progenitors is enhanced by cord blood plasma but expansion of the more mature subsets of progenitors is favored. *Blood Cells* 1994; 20: 436-454
29. Broxmeyer HE, Lu L, Cooper S, Ruggieri L, Li ZH, Lyman SD. Flt3-ligand stimulates/costimulates the growth of myeloid stem/progenitor cells. *Exp Hematol* 1995; 23: 1121-1129
30. Lazzari L, Lucchi S, Rebullia P, Porretti L, Puglisi G, Lecchi L, Sirchia G. Long term expansion and maintenance of cord blood hematopoietic stem cells using thrombopoietin, flt3-ligand, IL-6 and IL-11 in a serum free culture system. *Bjh* 2001; 112(2): 397-404
31. Rossmann T, Schroder B, Bug G, Muller P, Klenner T, Knaus R, Hoelzer D, Ottmann OG. Interleukin 3 improves the *ex vivo* expansion of primitive human cord blood progenitor cells and maintains the engraftment potential of SCID repopulating cells. *Stem cell* 2001; 19(4): 313-320
32. Donaldson C, Denning-Kendall P, Bradley B, Hows J. The CD34+CD38- population is significantly increased in hematopoietic cell expansion cultures in serum free compared to serum replete conditions: dissociation of phenotype and function. *BMT* 2001; 27: 365-371
33. Dreger P, Viehmann K, Steinmann J, Eckstein V, Muller-Ruchholtz W, Loffler H, Schmitz N.G. CSF-mobilized peripheral blood progenitor cells for allogeneic transplantation: comparison of T cell depletion strategies using different CD34+ selection systems or CAMPATH-1. *Exp Hematol* 1995; 23: 147-154
34. Fietz T, Berdel WE, Rieder H, Reufi B, Hopp H, Thiel E, Knauf WU. Culturing human umbilical cord blood: a comparison of mononuclear vs CD34+ selected cells. *Bone marrow transplant* 1999; 23: 1109-1115
35. Sandstrom CE, Bender JG, Papoutsakis ET, Miller WM. Effects of CDE34+cell selection and perfusion on *ex vivo* expansion of peripheral blood mononuclear cells. *Blood* 1995; 86: 958-970
36. Broxmeyer HE, Hangoc G, Cooper S, Ribeiro RC, Graves V, Yoder M, Wagner J, Vadhan-Raj S, Benninger L, Rubinstein P. Growth characteristics and expansion of human umbilical cord blood and estimation of its potential for transplantation in adults. *Proc. Natl. Acad. Sci. USA* 1992; 89: 4109-4113
37. Migliaccio G, Migliaccio AR, Druzin ML, Giardina PJ, Zsebo KM, Adamson JW. Effects of recombinant human stem cell factor (SCF) on the growth of human progenitor cells *in vitro*. *J Cell Physiol* 1991; 148: 503-509
38. Ohmizono Y, Sakabe H, Kimura T, Tanimukai S, Matsumura T, Miyazaki H, Lyman SD, Sonoda Y. Thrombopoietin augments *ex vivo* expansion of human cord blood-derived hematopoietic progenitors in combination with stem cell factor and flt3 ligand. *Leukemia* 1997; 11: 524-530
39. Zauli G, Vitale M, Visani G, Marchisio M, Milani D, Capitani S. *In vitro* growth of human fetal CD34+ cells in the presence of various combinations of recombinant cytokines under serum-free culture conditions. *Br J Haematol* 1994; 86: 461-467
40. Laiuppa JA, Papoutsakis ET, Miller WM. Oxygen tension alters the effect of cytokines on the megakaryocyte, erythrocyte, and granulocyte lineages. *Exp Hematol* 1998; 26: 835-843
41. Schibler KR, Li Y, Ohls RK, Nye NC, Durham MC, White W, Liechty KW, Le T, Christensen RD. Possible mechanisms accounting for the growth factor independence of hematopoietic progenitors from umbilical-cord blood. *Blood* 1994; 84: 3679-384
42. Migliaccio G, Migliaccio AR, Druzin ML, Giardina PJ, Zsebo KM, Adamson JW. Long-term Generation of colony-forming cells in liquid culture of CD34+ cord blood cells in the presence of recombinant human stem cell factor. *Blood* 1992; 79 (10): 2620-2627
43. De Felice L, Di Pucchio T, Breccia M, Agostini F, Mascolo MG, Guglielmi C, Ricciardi MR, Screnci M, Tafuri A, Carmini D, Arcese W. Flt3-L enhances the early stem cell compartment after *ex vivo* amolification of cord blood CD34+ cells. *Bone Marrow Transplant*, 1998 ; 22(1): 66-67
44. Rice A, Flemming C, Case J, Stevenson J, Gaudry L, Vowels M. Comparative study of the *in vitro* behavior of cord blood subpopulations after short term cytokine exposure. *Bone Marrow Transplant* 1999; 23: 211-220
45. McKenna HJ, de Vries P, Brasel K, Lyman SD, Williams DE. Effects of flt3 ligand on the *ex vivo* expansion of human CD34+ hematopoietic progenitor cells . *Blood* 1995; 86: 3413-3420
46. Gabbianelli M, Pelosi E, Montesoro E, Valtieri M, Luchetti L, Samoggia P, Vitelli L, Barberi T, Testa U, Lyman S. Montesoro E. Multi-level effects of flt3 ligand on human hematopoiesis: expansion of putative stem cells and proliferation of granulomonocytic progenitors/monocytic precursors. *Blood* 1995; 86: 1661-1670
47. Shah AJ, Smogorzewska EM, Hannum C, Crooks GM. Flt3 ligand induces proliferation of quiescent human bone marrow CD34+CD38- cells and maintains progenitor cells *in vitro*. *Blood* 1996; 87: 3563-3570
48. Hogge DE, Cashman JD, Humphries RK, Eaves CJ. Differential and synergistic effects of human Granulocyte-Macrophage Colony Stimulating Factor and human Granulocyte Colony Stimulating Factor on hematopoiesis in human long term marrow cultures. *Blood* 1991; 77(3): 493-99
49. Liu J, Li K, Yuen PM, Fok TF, Yau FW, Yang M, Li CK. *Ex vivo* of enriched CD34+ cells from the neonatal blood in the presence of thrombopoietin, a comparison with cord blood and bone marrow. *Bone Marrow Transplant* 1999; 24: 247-52
50. Piacibello W, Sanavio F, Garetto L, Severino A, Bergandi D, Ferrario J, Fagioli F, Berger M, Aglietta M. Extensive amplification and self-renewal of human primitive hematopoietic stem cells from cord blood. *Blood* 1997; 89 (8): 2644-2653

51. Sitnicka E, Lin N, Priestley GV, Fox N, Broudy VC, Wolf NS, Kaushansky K. The effect of erythropoietin on the proliferation and differentiation of murine hematopoietic stem cells. *Blood* 1996; 87: 4998
52. Kobayashi M, Laver JH, Lyman SD, Kato T, Miyazaki H, Ogawa M. Thrombopoietin supports proliferation of human primitive hematopoietic cells in synergy with Steel factor and/or interleukin-3. *Blood* 1996; 88: 429
53. Cohen SBA, Gluckman E, Rubinstein P, Madrigal JA : Cord blood characteristics: role in stem cell transplantation, 1st ed. London: Martin Dunitz Ltd. , 2000: 169-203
54. Khalili M, Alimoghaddam K, Soleimani M, Ghodis P, Hayat P, Moezi L, Arjman A, Mohyedin M, Ghavamzadeh A. Evaluation of the best condition for *ex vivo* expansion of hematopoietic stem cells for the propose of cord blood transplantation. *Yakhteh medical journal*. 2006; 8; 39-44
55. Piacibello W, Sanavio F, Garetto L, Severino A, Dane A, Gammaitoni L, Aglietta M. Differential growth factor requirement of primitive cord blood hematopoietic stem cell for self renewal and amplification vs. proliferation and differentiation. *Leukemia* 1998; 12: 718-727
56. Denning-Kendall PA, Nicol A, Horsley H, Donaldson C, Bradley B, Hows JM. *Is in vitro* expansion of human cord blood cells clinically relevant? *Bone Marrow Transplant* 1998; 21: 1628-1632
57. Kogler G, Callejas J, Sorg RV, Fischer J, Migliaccio AR, Wernet P. The effect of different thawing methods, growth factor combinations and media on the *ex vivo* expansion of umbilical cord blood primitive and committed progenitors. *Bone Marrow Transplant* 1998; 21: 233-241
58. Traycoff CM, Kosak ST, Grigsby S, Srouf EF. Evaluation of *ex vivo* expansion potential of cord blood and bone marrow hematopoietic progenitor cells using cell tracking an limiting dilution analysis. *Blood* 1995; 85: 259-268
59. Ruggieri L, Heymfeld S, Broxmeyer HE. Cytokine-dependent *ex vivo* expansion of early subsets of CD34+ cord blood myeloid progenitors is enhanced by cord blood plasma, but expansion of the more mature subsets of progenitors is favored. *Blood Cells* 1994; 20: 436-454
60. Moore MAS, Hoskins I. *Ex vivo* expansion of cord blood-derived stem cells and progenitors. *Blood Cells* 1994; 20: 468-481
61. Durand B, Eddleman K, Migliaccio AR, Migliaccio G, Adamson JW. Long-term generation of colony forming cells (CFC) from CD34+ human umbilical cord blood cells . *Leuk Lymphoma* 1993; 11: 263
62. Clements JM, Craig S, Gearing AJ, Hunter MG, Heyworth CM, Dexter TM, Lord BI. Biological and structural properties of MIP-1 alpha expressed in yeast. *Cytokine* 1992; 4: 76-82
63. Broxmeyer HE, Sherry B, Lu L, Cooper S, Oh KO, Tekamp-Olson P, Kwon BS, Cerami A. Enhancing and suppressing effects of recombinant murine macrophage inflammatory proteins on colony formation *in vitro* by bone marrow myeloid progenitor cells. *Blood* 1990; 76: 1110-1116
64. Broxmeyer HE, Sherry B, Cooper S, Lu L, Maze R, Beckmann MP, Cerami A, Ralph P. Comparative analysis of the human macrophage inflammatory protein family of cytokines (chemokines) on proliferation of human myeloid progenitor cells. *J Immunol* 1993; 150: 3448-3458
65. Graham GJ, Wright EG, Hewick R, Wolpe SD, Wilkie NM, Donaldson D, Lorimore S, Pragnell IB. Identification and characterization of an inhibitor of hemopoietic stem cell proliferation. *Nature* 1990; 344: 442-444
66. Eaves CJ, Cashman JD, Wolpe SD, Eaves AC. Unresponsiveness of primitive chronic myeloid leukemia cells to macrophage inflammatory protein 1 alpha, an inhibitor of primitive normal hematopoietic cells. *Proc Natl Acad Sci USA* 1993; 90: 12015-12019
67. Verfaillie CM, Catanzarro PM, Li W. Macrophage inflammatory protein 1 α , interleukin 3 and diffusible marrow stromal factors maintain human hematopoietic stem cells for at least eight weeks *in vitro*. *J Exp Med* 1994; 179: 643-649
68. de Wynter EA, Durig J, Cross MA, Heyworth CM, Testa NG. Differential Response of CD34⁺ Cells Isolated from Cord Blood and Bone Marrow to MIP-1 α and the Expression of MIP-1 α Receptors on These Immature Cells . *Stem Cells* 1998; 16(5): 349-356
69. JR Keller, Jacobsen SE, Dubois CM, Hestdol K, Ruscetti Fw. Transforming growth factor-beta: a bidirectional regulator of hematopoietic cell growth. *Int J Cell Cloning* 1992; 10: 2-11
70. Ploemacher RE, van Soest PL, Boudewijn A. Autocrine transforming growth factor beta 1 blocks colony formation and progenitor cell generation by hemopoietic stem cells stimulated with steel factor. *Stem Cells* 1993; 11: 336-347
71. De Bruyn C, Delforge A, Bron D, Ley P, de Hemptinne D, Stryckmans P. Modulation of human cord blood progenitor cell growth by recombinant human interleukin 3 (IL-3), IL-6, granulocyte-macrophage colony stimulating factor (GM-CSF) and stem cell factor (SCF) in serum- supplemented and serum-free medium. *Stem Cells* 1994; 12: 616-625
72. Berthier R, Valiron O, Schweitzer A, Marguerie G. Serum-free medium allows the optimal growth of human megakaryocyte progenitors compared with human plasma supplemented cultures: role of TGF beta. *Stem Cells* 1993; 11: 120-129
73. McNiece IK, Bertoncello I, Keller JR, Ruscetti FW, Hartley CA, Zsebo KM. Transforming growth factor beta inhibits the action of stem cell factor on mouse and human hematopoietic progenitors. *Int J Cell Cloning* 1992; 10: 80-86
74. CS Rosenfeld. Transforming growth factor-beta 1 augments macrophage-colony stimulating factor activity on human marrow. *Stem Cells* 1994; 12: 527-532
75. Metcalf D. The leukemia inhibitory factor (LIF). *Int J Cell Cloning* 1991; 9: 95-108
76. Williams RL, Hilton DJ, Pease S, Willson TA, Stahi J, Gearing DP, Nicola NA, Metcalf D. Myeloid leukemia inhibitory factor (LIF) maintains the developmental potential of embryonic stem cells. *Nature* 1988; 336: 684-687

77. Leary AG, Wong GG, Clark SC, Smith AG, Ogawa M. Leukemia inhibitory factor differentiation-inhibiting activity/human interleukin for DA cells augments proliferation of human hematopoietic stem cells. *Blood* 1990; 75: 1960-1964
78. Verfaillie C, McGlave P. Leukemia inhibitory factor/human interleukin for DA cells: a growth factor that stimulates the *in vitro* development of multipotential human hematopoietic progenitors. *Blood* 1991; 77: 263-270
79. Escary JL, Perreau J, Dumenil D, Ezine S, Brulet P. Leukemia inhibitory factor is necessary for maintenance of haematopoietic stem cells and thymocyte stimulation. *Nature* 1993; 363: 361-364
80. Koller MR, Bender JG, Miller WM, Papoutsakis ET. Expansion of primitive human hematopoietic progenitors in a perfusion bioreactor system with IL-3, IL-6 and stem cell factor. *Biotechnology (NY)* 1993; 11(3): 358-363
81. Koller MR, Manchel I, Maher RJ, Goltry KL, Armstrong RD, Smith AK. Clinical scale human umbilical cord blood cell expansion in a novel automated perfusion culture system *BMT* 1998; 21: 653-663
82. Van Zant G, Rummel SA, Koller MR, Larson DB, Drubachevsky I, Palsson M, Emerson SG. Expansion in bioreactors of human progenitor population from cord blood and mobilized peripheral blood. *Blood cells* 1994; 20: 482-491
83. Donnell Thomas E, Blumer KG, Donnell Thomas E, Forman S. Hematopoietic cell transplantation. The 2nd edition. 1999; 45: 481-493
84. Donaldson C, Armitage WJ, Denning-Kendall PA, Nicol AJ, Bradley BA, Hows JM. Optimal cryopreservation of human umbilical cord blood. *BMT* 1996; 18: 725-731
85. Moezzi L, Pourfathollah AA, Alimoghaddam K, Soleimani M, Ardjmand AR. The Effect of Cryopreservation on Colonogenic Capacity and *In vitro* Expansion Potential of Umbilical Cord Blood Progenitor Cells. *Transplantation Proceedings* 2005; 37: 4500-4503
86. Wang SY, Hsu ML, Huang MZ, Hsu HC, Tzeng CH, Hung JH, Ho CK. The activity in *ex vivo* expansion of cord blood myeloid progenitor cells before and after cryopreservation. *Acta Haematologica* 2001; 105(1): 38-44
87. Rice AM, Wood JA, Milross CG, Collins CJ, Case J, Nordon RE, Vowels MR. Prior cryopreservation of *ex vivo* expanded cord blood cells is not detrimental to engraftment as measured in NOD- SCID mouse model. *J Hematotherapy and stem cell research* 2001; 10: 157-165
88. Guenechea G, Segovi JC, Albella B, Lamana M, Ramirez M, Regidor C, Fernandez MN, Bueren JA. Delayed engraftment of nonobese diabetic/severe combined immunodeficient mice transplanted with *ex vivo*-expanded human CD34+ cord blood cells. *Blood* 1999; 93: 1097
89. Stiff P, Pecora A, Parthasarathy M, Preti R, Chen B, Douville J, Malhotra D, Harrison J, Bayer R, Goltry K, Armstrong RD, Smith A. Umbilical cord blood transplants in adults using a combination of unexpanded and *ex vivo* expanded cells: preliminary clinical observations. *Blood* 1998; 92(10): 2668
90. Shpall EJ, Quinones R, Hami L, Jones R, Bearman S, Cagnoni P, Giller R, Nieto Y, Roman-Unfer S, McNiece I. Transplantation of cancer patients receiving high dose chemotherapy with *ex vivo* expanded cord blood cells. *BLOOD* 1998; 92(10): 646A-646A 2667 Part 1 Suppl. 1
91. Kogler G, Nummerger W, Fischer J, Niehues T, Somville T, Gobel V, Wernet P. Simultaneous cord blood transplantation of *ex vivo* expanded together with non-expanded cells for high risk leukemia. *BMT* 1999; 24: 397-403
92. Pecora AL, Stiff P, Jennis A, Goldberg S, Rosenbluth R, Price P, Goltry KL, Douville J, Armstrong RD, Smith AK, Preti RA. Prompt and durable engraftment in two older adult patients with high risk chronic myelogenous leukemia (CML) using *ex vivo* expanded and unmanipulated unrelated umbilical cord blood. *BMT* 2000; 25: 797-799
93. Alimoghaddam K, Khalili M, Soleimani M, Arjmand A, Ghodsi P, Hayat P, Moezi L, Mohyedin M, Ghavamadeh A. Evaluation the effects of MIP-1a on *ex vitro* expansion of hematopoietic progenitor cells in different culture media for the propose of cord blood transplantation. *Khoon (blood)* 2006