

The Effect of Fetal Olfactory Mucosa on Tissue Sparing and Locomotor Recovery after Spinal Cord Hemisection in Rats

Hamdollah Delaviz, M.Sc.^{1*}, Mohammad Taghi Joghataie, Ph.D.¹,
Mehdi Mehdizadeh, Ph.D.¹, Mehrdad Bakhtiyari, Ph.D.²,
Maliheh Nobakht, Ph.D.¹, Samideh Khoei, Ph.D.³, Amrollah Rozbehe, Ph.D.¹

1. Anatomy Department, Cellular and Molecular Research Center, School of Medicine, Iran University of Medical Sciences, Tehran, Iran
2. Anatomy Department, School on Medicine, Lorestan University of Medical Sciences, Khoramabad, Iran
3. Physic Medical Department, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

* Corresponding Address: P.O. Box: 14155/6183, Anatomy Department, Cellular and Molecular Research Center, School of Medicine, Iran University of Medical Sciences, Tehran, Iran
Email: nilohamdi@yahoo.com

Abstract

Received: 23/Jan/2008, Accepted: 20/Apr/2008

Objective: Olfactory ensheathing cells (OECs) has been shown to have a neuroprotective effect after transplanted in brain and spinal cord injury (SCI). This study was conducted to determine the possible beneficial results of transplantation of fetal olfactory mucosa (FOM) that was the source of OECs in the recovery of locomotor function and in spinal tissue sparing after spinal cord hemisection.

Materials and Methods: Forty-eight adult female Sprague-Dawley rats were spinally hemisected at the L1 level and were randomized into the three groups of 16 animals. The first group, immunosuppressed injured animals were received cyclosporine A (CsA) and FOM graft. The second group was received CsA and fetal respiratory mucosa (FRM) graft, and the control group; non-immunosuppressed rats were received saline and gel foam. Locomotor performance was assessed weekly for 8 weeks after lesion, using locomotive rating scale developed by Basso, Bresnahan and Beattie (BBB). After behavioral assessment, the spinal cord was examined by a histologist for spinal tissue sparing.

Results: From weeks 6-8, the functional recovery of the FOM rats significantly increased in comparison to the FRM, although a significant difference in tissue sparing was not apparent. From weeks, 2-8 the functional recovery of the FOM and FRM groups as well as tissue sparing of the FOM group increased significantly compared to the control group.

Conclusion: Thus, the FOM treatment may be effective to promote functional recovery and partially preserving tissue sparing.

Keywords: Ensheathing Glia, Fetal Olfactory Mucosa, Spinal Cord Hemisection, Transplantation

Yakhteh Medical Journal, Vol 10, No 3, Autumn 2008, Pages: 185-192

Introduction

Following spinal cord injury (SCI) in adult mammals, regeneration of the axons and functional recovery are not successful. Many factors act as a physical and chemical barrier for axon regeneration, such as inhibitory molecules, an inhospitable extraneuronal environment, tissue scaring, cell death, glial cells and some meningeal cells (1, 2). Repeating words In order to reduce these barriers and to stimulate the remaining intrinsic plasticity of neurons to regenerate axons, external intervention is needed (1). Cell and tissue types from various sources have been considered and evaluated for therapy in the SCI, including peripheral nerve grafts (3), Schwann cells (4), and embryonic spinal cord tissue (5).

In addition, transplantation of olfactory ensheathing cells (OECs) into the injured spinal cord is another existing strategy. OECs have the unique property of enfolding olfactory axons along the entire axonal path from the olfactory mucosa (PNS) to the outer layer of olfactory bulb (CNS) (6) and prevention of their exposures to inhibitory molecules (7). The OECs is more accessible via the olfactory mucosa with a simple biopsy through the external nares than through the olfactory bulb (OB) (8). OECs from the olfactory lamia propria (OLP) have an increased rate of mitotic activity rate and migration than OECs from the OB (9). Functional improvement has been reported after transplantation of lamina pro-

pria into the complete thoracic spinal (10), and transplantation of olfactory mucosa in clinical trials have shown promising results (11). This study illustrates the feasibility of therapies designed to target FOM, to preserve tissue, and to improve functional outcome following spinal cord hemisection injury.

Materials and Methods

Surgical procedure and transplantation

All experiments were performed according to the guidelines of the Iranian Council for Use and Care of Animals Guidelines and were approved by the Animal Research Ethical Committee of Iran Medical University. Forty-eight Spruge-Dawley rats (2-3 months old, female) weighing between 200- 250 g were used in this study. Anesthesia was induced via intraperitoneal injection of a combination of ketamine (80 mg/kg) and xylazine (10 mg/kg). Under an operating microscope, a dorsal laminectomy of the T12 vertebrae was performed and the superior articular joint of the left side was removed. The L1 spinal segment was identified. Hemisection on the left side was performed by using a pair of iridectomy scissors, effectively disrupting all major unilateral descending pathways (12). Median sagittal vein was used as an anatomical landmark to demarcate the median plane. The FOM and FRM were removed from E18 as described previously. (8) One piece (0.5 -1mm) of OM or RM was gently situated between the rostral and caudal sites of the lesion and another piece (1 mm) was placed over the lesion site and covered with gel foam. Sixteen rats were randomly chosen to receive FOM, another 16 rats to receive FRM, and a final 16 to receive gel foam as the control group. (Two rats from the FRM group died as a result of anesthetic complications during preparation for transplantation). The overlying muscles were sutured in layers, and the skin was closed with wound clips. The FOM and FRM groups received intraperitoneal injection of cyclosporine A (Sandimmune, Sandoz, Basel, Switzerland) at a dose of 10 mg for every kg of body weight to prevent rejection of the graft (13) and the control group received saline. Cyclosporine and saline injections started 2 days before the transplantation procedures and continued throughout the experiment, daily. Otherwise, all three groups following surgery were maintained under the same conditions with free access to food and water. Postoperative treatments included saline (1.0 cc s.c.) for rehydration and penicillin-G (0.35ml/kg i.m) as a prophylactic antibiotic. Their bladders were manually expressed twice a day for the first 3 days.

Immunostaining for identification of OECs in fetal olfactory mucosa

The FOM and FRM were removed from E18 and immediately transferred to separate 35 mm Petri dishes containing 4 % paraformaldehyde in 0.1 M PB, (pH=7.4, 4°C) for two hours. The tissues were then cryoprotected (in 30% sucrose) overnight and embedded in tissue

freezing medium. Transverse sections (20 µm – thick) of the tissues were made with a freezing microtome (leica cryostat, CM 3000), mounted onto gelatin-coated glass slides, and immunostained with anti- p75 (Mouse anti-nerve growth factor- receptor monoclonal antibody; cat number MAB365, Chemicon) 1:100, overnight. The following day, the sections were washed in PBS and incubated with a secondary anti-body 1:200 (molecular probes, Eugen, Oregon, USA, Alexa Fluor 488nm wave length) in PBS for one hour at room temperature. The slides were then washed with PBS, cover slipped and viewed with a fluorescence microscopy (Fig 1). Control sections without the primary antibody were prepared for each tissue as well.

Assessment of hindlimb motor function

Hindlimb motor function was assessed based on the Basso, Beattie, and Bresnahan (BBB) Locomotor Rating Scale (14) at one day post-surgery and weekly for a total of 8 weeks. For BBB assessment, the rats were allowed to move individually for 5 minute on a smooth, nonslip floor in an open field (200x100 cm). Hindlimb motor function was scored from 0 to 21 based on the performance of the ipsilateral hindlimb by an observer who was not a participant in the experiment. Expectedly spinal cord hemisection of the thoracolumbar region resulted in pronounced paralysis on the ipsilateral hindlimb (impaired hindlimb), with the partially recovery by 3 weeks (15). Hemisections were considered complete when they included the dorsal column, Lissaur's tract, lateral and ventral column, and gray matter.

Estimating the amount of tissue sparing

following behavioral assessment, five rats from each of the groups were deeply anesthetized with an overdose of ketamine (200 mg/kg body weight) and xylazine (20 mg/kg body weight), perfused through their hearts with 0.9% NaCl in distilled water (200 ml) followed by 4% paraformaldehyde in 0.1 M PB, pH 7.4 (4°C, 500 ml). A 5-mm-long spinal cord segment with epicenter of the transplanted tissue were removed, fixed in the same solution for 24 hours, cryoprotected in 30% sucrose overnight and cut into 50µm serial horizontal sections by a leica cryostat. Every fourth horizontal section from rostral to the caudal of the segment, mounted onto gelatin-coated glass slides, was stained with cresyl violet or H&E dehydrated and cover slipped. To determine the volume of spared spinal tissue, low-power magnification (x4) images of the spinal cord sections were taken with the aid of a digital camera (DP 11) attached to the microscope (Olympus AX70). By using an image analysis computer system (olysia) on every fourth horizontal section of a 5-mm- long spinal segment, the total area of the left spinal cord and the area of damaged spinal tissue (cavity) in left half of the spinal cord were determined. The border of the damaged area was defined as an obvious discontinuity in density of small cells and the absence of healthy-looking spinal neurons (16). As

previously described (16), the measurements of each section were summed per rat and multiplied by 4 to give the total area of the left half and total area of the damaged tissue of a 5-mm-long spinal segment. This value was expressed as a percentage of the total volume of the left half of a 5-mm-long segment from the same cord level from uninjured rats ($n=5$).

Statistical analysis

All data are expressed as mean \pm SD. One-way ANOVA was used for data analysis, followed by the Tukey test for post hoc analysis. A P -value < 0.05 was considered to be statistically significant.

Results

Olfactory ensheathing cells are present in the lamina propria of fetal olfactory mucosa

The lamina propria beneath the olfactory epithelium contains olfactory nerve bundles surrounded by OECs which identified immunohistochemically with an antibody to P75 NGFR (Fig 1C). The OM removed by its posterior position on the nasal septum and identified from RM by semicircular line (Fig 1A).

The FOM transplants improve locomotor recovery

One day after operation, the hemisected animals displayed loss of ipsilateral hindlimb function with no observable movement. Of the 46 rats that received left

hemisection, the six rats showed consistent weight supported plantar steps and consistent forelimb – hind limb (FL-HL) coordination (BBB= 15). For one week after operation, subsequent histological evaluation showed that in these cases the full hemisection of the L1 segment was incomplete and they were thus excluded from the study. Of the remaining 40 rats, the 4 rats were killed and horizontal and parasagittal section confirmed the full hemisection (Fig 2). The inflammatory reactions were observed (have seen delete) at the lesion epicenter one week after lesion (Fig 2A). Then it followed by infarction and cysts formation four weeks after the hemisection (Fig 2B).

The following BBB scores increased considerably for the FOM, FRM and control groups, respectively, on the 8th day post surgery (mean \pm SD): 10.41 ± 1.31 , 10.16 ± 1.85 and 8.75 ± 1.76 (Fig 3). From weeks 2-8, the FOM and FRM rats demonstrated a significant increase in the movements of their hind limbs compared to the control group ($p < 0.05$). The FOM and FRM rats recovered at similar rates during the 5th week post-injury but from weeks 6-8, the FOM treated rats slightly improved their walking behaviors in comparison to the FRM group. Furthermore, the FOM group showed consistent plantar stepping and consistent forelimb-hindlimb (FL-HL) coordination, whereas FRM demonstrated limited FL-HL coordination.




Fig 1: (A) Photograph of the nasal septum of a fetal (E18) rat indicating the locations of the OM and the RM separated from each other by a semicircular line (arrow). (B) H&E staining of the nasal septum: arrow indicates the transitional zone between OM and RM. (C) Anti-p75 immunoreactivity (light green) indicates olfactory ensheathing cells lying into the lamina propria of the OM (arrowhead): the olfactory epithelium and lamina propria of the RM are indicated by double arrow. (D) The lamina propria of the RM contains no p75 immunoreactivity. Scale bar: 50 μ m (B, C, D). OM, olfactory mucosa; RM, respiratory mucosa.




Fig 2: (A) Cresyl violet stained, 20 μm thick sections, sagittal sections through the injury site one week after injury. Inflammatory reaction (arrowhead) formed at the lesion epicenter. (B) H&E stained, 20 μm thick horizontal of L1 lumbar spinal cord 28 days after lesion. Stars indicate cystic cavities. Arrow indicates median fissure; D, distal; P, proximal; Scale bar: A, 100 μm ; B, 200 μm .

On the 8th week, the FRM animals achieved a plateau score of 15 ± 1.27 , whereas the FOM rats reached an average score of 16.5 ± 0.90 . One-way ANOVA results showed significant difference between FOM and FRM groups during the weeks 6 to 8 ($p < 0.05$).




Fig 3: The Means BBB scores of the different groups following hemisection. During the two weeks post-injury all groups recovered nearly at similar rates, but from weeks 2-8, the FOM and FRM rats demonstrated significant performance improvement over the control group. Values represent means \pm SD. $n=12$, * $p < 0.05$: compared to FRM group.

Histomorphology of the grafts and lesion site after eight weeks

Histological evaluation of the spinal cord from the epicenter of the lesion site revealed that the FOM grafts,

survived, grew and proliferated to fuse with the parenchyma of the host tissue. There were no obvious glial boundaries at the host-graft interface (Fig 4, A, D). In the FRM rats, there were a few cavities at the host-graft interface (Fig 4, B, E). In the both group, a clear boundary consisting of a different cytoarchitecture was observed at the host-graft interface (Fig 4, A, B). In the control rats, there were large cysts that replaced all of the area previously occupied by gray matter and white matter (Fig 4, C, F). In some of the rats from the control group, fibrous tissue had replaced the lesion site (Fig 4C).

Transplantation of olfactory mucosa improve partially spinal tissue sparing

To ensure consistency in the analyses of tissue sparing, all sections were seen through the same illumination level, and microscope and digital camera settings were maintained constant throughout the image capturing sessions. The volume of the left half of a 5-mm-long spinal cord segment of a normal uninjured rat at the L1 level was (mean \pm SD): $11.85 \pm 0.12 \text{ mm}^3$. In contrast, the spared tissue of the left half in the FOM, FRM, and control groups were $5.65 \pm 0.54 \text{ mm}^3$, $5.0 \pm 0.94 \text{ mm}^3$, and $4.25 \pm 0.11 \text{ mm}^3$, respectively (Fig 5). Although an ANOVA revealed no significant difference in the percentage of spared tissue between the FOM and FRM rats, there was a modest increase in tissue sparing as a result of FOM treatment in comparison to the control group ($p < 0.05$). Whereas there was no significant difference in the percentage of spared tissue between the FRM group and control rats, indicating that the FOM grafts enhanced tissue sparing better than FRM after hemisection.




Fig 4: Parasagittal and transverse sections (50 μm) at the lesion epicenter at the L1 level, after a hemisection injury. In the FOM rats, the grafts (A, D) fused with the host tissue and formed a seamless continuous spinal cord, whereas in some of the FRM animals, (B, E) there were a few cavities at the host-graft interface (arrowhead) at the 8th week after transplantation. At the lesion epicenter, the control group showed characteristic necrosis and cavitations of the spinal parenchyma, (C, D). Arrows indicate central canal or median fissure, arrowheads indicate cavity, T, transplant; F, fibroses tissue; P, proximal; D, distal; scale bar: A, 200 μm ; B, C, 300 μm ; D, 100 μm ; E, 200 μm ; F, 400 μm .




Fig 5: Graph representing the spared spinal tissue in the left half of the spinal cord at the L1 level at the lesion epicenter in different groups. FOM showed a significant difference in the percentage of spared tissue compared to the control groups. Values represent means \pm SD, n=5.*p<0.05 compare to the control group.

Discussion

This experiment shows that the fetal olfactory mucosa has neuroprotective effects that improve locomotor recovery, axonal regeneration and attenuation of lesion size compared to the cyclosporine and fetal respiratory mucosa. Our behavioral and histological data demonstrate that the animals used in this study had completely hemisectioned spinal cords (Fig 2). The ascending and descending tract were interrupted after transection of the spinal cord and spinal tissue undergo apoptosis and inflammatory reaction as observed at the first week of the post-injury (17). In this study, we confirm following L1 hemisection in rats, the ipsilateral hindlimb is initially paralyzed and partially recovers within 2 to 3 weeks (18). The functional recovery and promotion of axonal regeneration by OECs from the olfactory bulb have been studied in various adult rat spinal cord injury models (19-22). Unlike rats, the olfactory bulb in humans is comparatively small and relatively inaccessible (23). Present study shows the transplantation of

FOM into the spinal cord injury is safe and feasible. The FOM rats showed that grafts entirely filled the lesion site between the rostral and caudal host tissue and formed an interface without cysts or scars (Fig 4, A, D). The absence of scar formation may reflect the mild host immune reaction attributable to immunosuppression by CsA (24). The scar tissue was visible in nonimmunosuppressive rats. OECs transplants did not prevent glial scar formation (19). Importantly, our data indicate that the FOM treatment was important to decrease damaged area and cavity formation following a hemisectioned lesion. Morphometric analyses indicated that the FOM rats had a 48% of preservation tissue around the injury site compared to a similar segment in the uninjured rats; this value was significantly greater than the volume of the spared tissue in the control group (35%). It has been documented that small changes in tissue sparing can significantly impact locomotor function (25). Thus, a 6% increase in tissue sparing in the FOM rats in comparison to the FRM group, and a similar 12.5% increase in comparison to the control group may indeed contribute to improve functional recovery. The improvement of functional recovery in FRM compared to the control group may be the effect of CsA, rather than FRM. The fibroblast and endothelial cells in FRM are not reported to assist spinal cord regeneration (8). Although, Schwann cells and macrophages can assist axonal regeneration but they are present in low number in both olfactory and respiratory lamina propria (8, 21). Whereas cyclosporine A promotes axonal regeneration in rats submitted to transverse section of the spinal cord (26) and reduces delayed motor neuron death after spinal cord ischemia in rabbits (27). Cyclosporin-A promotes neuroprotection by diminishing both demyelination and neuronal cell death, resulting in a better motor outcome after spinal cord injury (24). We cannot rule out that the growth factor in connective tissue of the grafts may facilitate endogenous repair (28). Also, there are reports that the extra cellular matrix (ECM) are neuroprotective and assist spinal cord repair (29, 30).

The protection effect of trophic factors of transplanted olfactory mucosa at the lesion site (31) may benefit functional recovery. Intrathecal injections of bFGF alone could preserve more spared tissue and enhance the functional recovery after spinal cord contusion (32). Immunostaining for trophic factors at the lesion site of OM-transplanted rats in the transected facial nerve showed increased expression of NGF, BDNF, and FGF-2 (31). The increase of the BBB scores on the 42nd day after the operation in the FOM rats in the present study shows that the olfactory mucosa may provide a weaker but long-lasting secretion of neurotrophins and bFGF at the lesion site (31). In other experiments, OECs from LP transplantation into the cervical dorsolateral funiculus crush provides the greatest opportunity to promote re-

generation or reconnect local circuitry (9). Transplantation of olfactory lamina propria promotes locomotor recovery after delaying transplantation into the transected spinal cord (8) and promotes partial recovery in paraplegic rats (14). It may be the OECs work optimally in concert with other cells or connective tissue element (23). In this study, not only OECs and CsA but also neural progenitor cells in the olfactory epithelium are probably responsible for tissue sparing and promotion of recovery in the FOM rats. Adult olfactory epithelium contains multipotent progenitor cells that give rise to neurons and non-neural cells (33). Human adult olfactory neural progenitor cells rescue axotomized rodent rubrospinal neurons and promote functional recovery (34). Thus, FOM have the potential to provide an efficient and renewable source of cells for autologous transplantation studies for spinal cord injury.

Conclusion

This study shows that transplantation of fetal olfactory mucosa with its lamina propria and of olfactory neuroepithelium results in promotion of tissue sparing and in functional recovery in mammals with partial spinal cord injury. Whatever the eventual outcome of these investigations, further work on the cell biology of the FOM and its behavior following transplantation will advance our understanding to the development of ways of spinal cord repair.

Acknowledgement

This research study was financially supported by a grant from Iran University of Medical Sciences (Tehran – Iran, No; 364). Also, we wish to thank Dr Naghmeh Fouladi from King College for excellent technical support and Hamoun Delaviz from Toledo, Ohio University, for revising of this article.

References

1. Santoes-Benito FF, Ramon-Cueto A. Olfactory ensheathing glia transplantation: A Therapy to promote repair in the mammalian central nervous system. *Anat Rec (New Anat)* 2003; 271: 77-85
2. Boyd JG, Skihar V, Kawaja M, Doucette R. Olfactory ensheathing cells: historical perspective and therapeutic potential. *Anat Rec (New Anat)* 2003; 271B: 49-60
3. Fernandez E, Pallini R, Minciacchi D, Sbriccoli A. Peripheral nerve autografts to the rat spinal cord: study on the origin and course of regenerating fibers. *Acta Neurochir* 1986; 82: 57-63
4. Guest JD, Rao A, Olson L, Bunge MB, Bunge RP. The ability of human Schwann cell grafts to promote regeneration in the transected nude rat spinal cord. *Exp Neurol* 1997; 148: 502-522
5. Bregman BS. Spinal cord transplants permit the growth of serotonergic axons across the site of neo-

- natal spinal cord transection. *Dev Brain Res* 1986; 34: 265-279
6. Doucette R. PNS-CNS transitional zone of the first cranial nerve. *J Comp Neurol*; 1991; 312: 451-466
 7. Pasterkamp RJ, De Winter F, Holtmaat AJ, Verhaagen J. Evidence for a role of the chemorepellent Semaphorin III and its receptor Neuropilin-1 in the regeneration of primary olfactory axons. *J Neurosci* 1998; 18: 9962-9976
 8. Lu J, Feron F, Ho SM, Mackay-Sim A, Waite PM. Transplantation of nasal olfactory tissue promotes partial recovery in paraplegic adult rats. *Brain Res* 2001; 889: 344-357
 9. Richter MW, Fletcher PA, Liu J, Tetzlaff W, Roskams AJ. Lamina propria and olfactory bulb ensheathing cells exhibit differential integration and migration and promote differential axon sprouting in the lesioned spinal cord. *J Neurosci* 2005; 25(46):10700-10711
 10. Lu J, Feron F, Mackay-Sim A, Waite PM. Olfactory ensheathing cells promote locomotor recovery after delayed transplantation into transected spinal cord. *Brain* 2002; 125: 14-21
 11. Lima C, Pratas-Vital J, Escada P, Hasse-ferreira A, Capucho C, Peduzzi J. Olfactory mucosa autograft in human spinal cord injury: A pilot clinical study. *JSCM* 2006; 29: 191-203
 12. Houle J, Jin Y. Chronically injured supraspinal neurons exhibit only modest axonal dieback in response to a cervical hemisection lesion. *Exp Neurol* 2001; 169: 208-217
 13. Hentall ID, Noga BR, Sagen J. Spinal allografts of adrenal medulla block nociceptive facilitation in the dorsal horn. *J Neurophysiol* 2001; 85: 1788-1792
 14. Basso DM, Beattie MS, Bresnahan JC. A sensitive and reliable locomotor rating scale for open field testing in rats. *J Neurotrauma* 1995; 12: 1-21
 15. Qin DX, Zou XL, Luo W, Zhang W, Zhang HT, et al. Expression of some neurotrophins in the spinal motoneurons after cord hemisection in adult rats. *Neurosci Lett* 2006; 410: 222-227
 16. Takami T, Oudega M, Bates ML, Wood PM, Kleitman N, Bunge BB. Schwann cell but not olfactory ensheathing glia transplants improve hindlimb locomotor performance in the moderately contused adult rat thoracic spinal cord. *J Neurosci* 2002; 22(15): 6670-6681
 17. Hains BC, Black JA, Waxman SG. Primary cortical motor neurons undergo apoptosis following axotomizing spinal cord injury. *J Comp Neurol* 2003; 462: 328-341
 18. Hains BC, Johnson KM, McAdoo DJ, Eaton MJ, Hulsebosch CE. Engraftment of serotonergic precursors enhances locomotor function and attenuates chronic central pain behavior following spinal hemisection injury in the rat. *Exp Neurol* 2001; 171: 361-378
 19. Ramon-Cueto A, Cordero MI, Santos-Benito FF, Avila J. Functional recovery of paraplegic rats and motor axon regeneration in their spinal cords by olfactory ensheathing glia. *Neuron* 2000; 25: 425-435
 20. Verdu E, Garcia-Alias G, Fores J, Pez-vales R, Navarro X. Olfactory ensheathing cells transplanted in lesioned spinal cord prevent loss of spinal cord parenchyma and promote functional recovery. *GLIA* 2003; 42: 275-286
 21. Mackay-Sim A. Olfactory ensheathing cells and spinal cord repair. *Keio J Med* 2005; 54 (1): 8-14
 22. Sasaki M, Hains BC, Lankford KL, Waxman SG, Kocsis JD. Protection of corticospinal tract neurons after dorsal spinal cord transection and engraftment of olfactory ensheathing cells. *GLIA* 2006; 53: 352-359.
 23. Franklin RJ. Obtaining olfactory ensheathing cells from extra-cranial sources a step closer to clinical transplant-mediated repair of the CNS? *Brain* 2002; 125: 2-3
 24. Ibarra A, Correa D, Willms K, Merchant M.T, Guizar-Sahagun G, Grijalva I, Madrazo I. Effects of cyclosporin-A on immune response, tissue protection and motor function of rats subjected to spinal cord injury. *Brain Res* 2003; 979(1): 165-178
 25. Basso DM, Beattie MS, Bresnahan JC. Graded histological and locomotor outcomes after spinal cord contusion using the NYU weight-drop device versus transection. *Exp Neurol* 1996; 139: 244-256
 26. Teichner A, Morselli E, Buttarelli, FR, Caronti B, Pontieri FE, Venturini G, et al. Treatment with cyclosporine A promotes axonal regeneration in rats submitted to transverse section of the spinal cord. *J Hirnforsch* 1993; 34(3): 343-349
 27. Sato M, Horinouchi T, Sakurai M, Murakami N, Sato S, Kato M. Cyclosporin A reduces delayed motor neuron death after spinal cord ischemia in rabbits. *Ann Thorac Surg* 2003; 75: 1294-1299
 28. Mumm JS, Shou J, Calof AL. Colony forming progenitors from mouse olfactory epithelium: evidence for feedback regulation of neuron production. *Proc Natl Acad Sci USA* 1996; 11176-11172
 29. Carbonetto S, Gruver MM, Turner DC. Nerve fiber growth in culture on fibronectin, collagen and glycosaminoglycan substrates. *J Neurosci* 1983; 3: 2324-2335
 30. Dityatev A, Schachner M. Extracellular matrix molecules and synaptic plasticity. *Nat Rev Neurosci* 2003; 4:456-468
 31. Guntinas-Lichius O, Wewetzer K, Tomov TL, Az-zolin N, Kazemi S, Streppel M, et al. Transplantation of olfactory mucosa minimizes axonal branching and promotes the recovery of vibrissae motor performance after facial nerve repair in rats. *J Neurosci* 2002; 22(16): 7121-7131
 32. Rabchevsky AG, Fugaccia I, Turner AF, Blades DA, Mattson MP, Scheff SW. Basic fibroblast growth factor (bFGF) enhances tissue sparing and functional recovery following moderate spinal cord injury. *J Neurotrauma* 1999; 16: 817-829
 33. Huard JM, Youngentob SL, Goldstein BJ, Luskin MB, Schwob JE. Adult olfactory epithelium contains multipotent progenitor cells that give rise to neurons and non-neural cells. *J Comp Neurol* 1998; 400: 469-486

Delaviz et al.

34. Xiao M, Klueber K.M, Lu C, Guo Z, Marshalla CT, Wang H, et al. Human adult olfactory neural progenitors rescue axotomized rodent rubrospinal neurons

and promote functional recovery. *Exp Neurol* 2005; 194: 12-30

Archive of SID