

Investigation The Role of Gender on The *HLA-DRB1* and *-DQB1* Association with Type 1 Diabetes Mellitus in Iranian Patients

Arezou Sayad, Ph.D.^{1,2}, Mohammad Taghi Akbari, Ph.D.², Mohammad Pajouhi, M.D.⁴, Feridoon Mostafavi, M.D.⁵, Anooshirvan Kazemnejad, Ph.D.⁶, Mahdi Zamani, Ph.D.^{1,3*}

1. Department of Neurogenetics, Iranian Centre of Neurological Research, Tehran University of Medical Sciences, Tehran, Iran
2. Department of Medical Genetics, Faculty of Medical Science, Tarbiat Modares University, Tehran, Iran
3. Department of Medical Genetics, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran
4. Department of Medical Science, Endocrine Research Center, Faculty of Medical Science, Tehran University of Medical Sciences, Tehran, Iran
5. Department of Endocrinology, Children Medical Center, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran
6. Department of Biostatistics, Faculty of Medical Science, Tarbiat Modares University, Tehran, Iran

* Corresponding Address: P.O. Box: 1417613151, Department of Neurogenetics, Iranian Centre of Neurological Research, Tehran University of Medical Sciences, Tehran, Iran
Email: Mzamani@tums.ac.ir

Received: 20/Nov/2011, Accepted: 14/Oct/2012

Abstract

Objective: Type 1 diabetes mellitus (T1D) is an autoimmune and multifactorial disorder. Subsequent analysis on human leukocyte antigen (HLA) region shows that *HLA-DRB1* and *-DQB1* genes have the strongest association with T1D. In this study, for the first time, we investigated the influence of gender on the *HLA-DRB1* and *-DQB1* association with type 1 diabetes mellitus in Iranian patients in order to determine gender dependent HLA heterogeneity in Iranian T1D patients.

Materials and Methods: In this case control study, the *HLA-DRB1* and *-DQB1* typing were performed on 105 Iranian T1D patients and 100 healthy controls. The data were evaluated by using Fisher exact test.

Results: Our results indicate that *DRB1*04:01*, *DQB1*03:02* alleles and *DRB1*04:01-DQB1*03:02* haplotype were significantly more frequent in male T1D patients than females. Also, *DRB1*03:01*, *DRB1*15:01*, *DQB1*06:01* alleles, *DQB1*03:01/05:01* genotype, *DRB1*03:01-DQB1*02:01* and *DRB1*15:01-DQB1*06:01* haplotypes were significantly higher in female T1D group than males. Furthermore, our results showed that *DRB1*04:01* and *DQB1*03:02* alleles were significantly more frequent in male T1D patients 1-5 years old at onset than females with similar condition. The *DRB1*03:01* allele and *DRB1*03:01-DQB1*02:01* haplotype were significantly higher in female T1D patients 6-10 years old at onset than males with similar condition. The *DRB1*15:01* allele and *DRB1*15:01-DQB1*06:01* haplotype were significantly more frequent in female T1D patients 16-20 years old at onset than males with similar condition.

Conclusion: Our findings suggest that gender has a significant influence on the distribution of *HLA-DR* and *-DQ* alleles, genotypes and haplotypes. Also, distribution of the *HLA-DRB1* and *-DQB1* alleles, genotypes and haplotypes vary based on the gender of T1D patients in different age at onset.

Keywords: Type 1 Diabetes, *HLA-DRB1*, *HLA-DQB1*, Gender, Age at Onset

Cell Journal(yakhteh), Vol 15, No 2, Summer 2013, Pages: 108-115

Citation: Sayad A, Akbari MT, Pajouhi M, Mostafavi F, Kazemnejad A, Zamani M. Investigation the role of gender on the *HLA-DRB1* and *-DQB1* association with type 1 diabetes mellitus in Iranian patients. Cell J. 2013; 15(2): 108-115.

Introduction

Type 1 diabetes (T1D) is a chronic autoimmune disease which is caused by the absence of insulin synthesis. It is characterized by the immunologically T lymphocyte mediated destruction of the insulin producing cells (beta cells) in the pancreatic islets of langerhans (1). The rate of the disease concordance in siblings (6%) and monozygotic twins (35-50%) shows that both genetic and environmental factors are contributed to the disease risk (2, 3). Therefore, the T1D is known as multifactorial disorder. The human leukocyte antigen (*HLA*) genes are located on the short arm of chromosome 6. The *HLA* genes are contributed in 50% of the genetic risk for T1D (4, 5). For the first time, the association of T1D and *HLA* class I alleles was studied in 1973 (6). Association of these alleles is secondary to the association of *HLA-DR* and *-DQ* alleles of the *HLA* class II which were investigated later (4). The *HLA-DRB1* and *-DQB1* alleles are identified as the most important alleles which are associated with T1D in many ethnic groups (7-9). Due to strong linkage disequilibrium between the *HLA-DRB1* and *-DQB1* alleles, the highest genetic risk is conferred by their haplotypes in compared with particular alleles (10-16). The studies about the incidence of T1D showed that, Finland, Sardinia and Scandinavia have the highest worldwide prevalence of T1D (>35/100,000/year). T1D is less frequent in Asian populations (0.4-1.1/100000/year) compared to Caucasians. The Japanese, Chinese and Koreans have a very low T1D incidence, approximately 1/100000/year (17). The high incidence of T1D in Scandinavians correlates with the high frequency of the *DRB1*04:01-DQA1*03:01-DQB1*03:02* haplotype. The high incidence of T1D in Sardinians is due to both the high frequency of another susceptible haplotype (*DRB1*03:01-DQA1*05:01-DQB1*02:01*), and the low frequency of protective haplotype (*DRB1*15:01-DQA1*01:02-DQB1*06:02*) (18). In Japanese population, the absence of the *DR3* alleles correlates with the low incidence of T1D. The share of *DR3/4* genotype is also less in Koreans than in Caucasians (19). There are differences in the *HLA-DRB1* and *-DQB1* gene profiles according to the gender and age at onset (20, 21). Some studies reported that *HLA-DRB1*03:01/09:01* was significantly more frequent in female T1D patient, whereas *HLA-DRB1*03:01/04:01* had

more frequency in male T1D patients (21, 22). Another study showed that *DQ6.2 (DQA1*01:02-DQB1*06:02)* was more negatively associated with T1D in females than males (23). Finding whether gender influence on susceptibility to or protection against T1D, can help to better follow-up the T1D patients according to their sex category. Also, knowing the gender dependent *HLA* heterogeneity among T1D patients indicates influence of sex dependent factor, as the hormones, on the functional effects of the *HLA* molecules. The individuals carrying alleles which are associated with younger age at onset in each female or male group should take care under preventive treatment.

In the present study, we aimed to examine the influence of the gender on susceptible and protective alleles, genotypes and haplotypes of the *HLA-DRB1* and *-DQB1* to find gender dependent *HLA* heterogeneity in Iranian T1D patients.

Materials and Methods

Subject and control groups

In this case control study, one hundred and five unrelated Iranian patients with T1D mellitus with a mean standard deviation age of 9.4 ± 4 years and an age range of 1-17 years (45% male and 55% female) were studied in the Department of Neurogenetics of the Iranian Center of Neurological Research. Patients with other type of diabetes, such as type 2 diabetes, maturity onset diabetes of youth and latent diabetes were not included in this study. Subjects were collected from the different hospitals located in Tehran and were diagnosed as having type1 diabetes mellitus by endocrinologist according to the WHO criteria. All patients as well as controls were of Iranian descent with mixed ethnic origin, and they were selected from different ethnic groups. One hundred healthy, ethnically and sex matched individuals with no personal or family history of autoimmune disorders (mean age of 36.2 ± 7.8 years and age range of 23-68 years) were selected as a control group in the same geographical area as patients (47% male and 53% female). All guardians were informed of the study and gave their written consent. The study was approved by the Ethnic Committee of the Tarbiat Modares University.

HLA-DRB1 and -DQB1 typing and data collection

DNA was extracted from peripheral venous blood

specimens using salting out method (24). The high resolution *HLA-DRB* typing was performed by Innolipa DRB kit (Innogenetics NV, Belgium). It was conducted according to the manufacturer's recommendations. HISTO TYPE SSP Low Resolution Kits (BAG Health care, Germany) were used for *HLA* Typing of *DQB1* as instructed by the manufacturer. Questionnaire information (such as sex, age at onset, ethnic, autoimmune or genetic history) of all patients and controls were collected. According to the results of *HLA-DRB* and *-DQB* typing, we determined susceptible and protective alleles, genotypes and haplotypes (25). Individuals were divided into two groups of male and female. Moreover, each male and female group was categorized into four groups based on age categories at onset (1-5, 6-10, 11-15 and 16-20). Distribution of susceptible and protective alleles, genotypes and haplotypes in each group was assessed.

Statistical methods

The data were evaluated by using Fisher's exact test. Odds ratio (OR) was calculated. The p values less than 5% were considered statistically significant.

All data were analyzed by SPSS 18.

Results

All patients and controls were divided into two groups of male and female. Furthermore, each male and female group was categorized into four groups based on age categories at onset (1-5, 6-10, 11-15 and 16-20). Distribution of susceptible and protective alleles, genotype and haplotype in each group was determined.

Distribution of susceptible and protective HLA-DRB1 and -DQB1 alleles according to the gender of T1D

*HLA-DRB1*04:01* allele had a 15.5 times higher risk of developing T1D in males than females. *HLA-DQB1*03:02* allele ($p=0.01$; OR: 6.273) was significantly higher in male T1D patients compared to females. Female carriers of *DQB1*06:01* allele had a 14 times higher risk of developing T1D than males. *DRB1*03:01* ($p=0.01$; OR: 0.244) and *HLA-DRB1*15:01* ($p=0.03$) alleles were significantly more frequent in female T1D patients than males (Table 1).

Table 1: Distribution of susceptible and protective *HLA-DRB1* and *-DQB1* alleles according to the gender of T1D

Allele	Gender	T1D	Control	P	OR (CL)
<i>HLA-DRB1*0401</i>	M	31	3	0.01	15.5 (1.813-132.543)
	F	2	3		
<i>HLA-DRB1*0301</i>	M	18	13	0.01	0.244 (0.079-0.751)
	F	34	6		
<i>HLA-DRB1*1301</i>	M	0	9	0.4	-
	F	1	7		
<i>HLA-DRB1*1501</i>	M	0	17	0.03	-
	F	4	11		
<i>HLA-DQB1*0201</i>	M	43	17	0.2	-
	F	31	21		
<i>HLA-DQB1*0302</i>	M	46	4	0.01	6.273 (1.507-26.106)
	F	11	6		
<i>HLA-DQB1*0301</i>	M	9	24	1	-
	F	15	40		
<i>HLA-DQB1*0601</i>	M	1	21	0.01	0.071 (0.007-0.682)
	F	6	9		

M; Male, F; Female, NS; Not significant and P; Value of Fisher's exact test.

Distribution of susceptible and protective HLA-DRB1 and -DQB1 genotypes were based on the gender of T1D

Frequency of susceptible and protective HLA-DRB1 and -DQB1 genotypes with respect to the gender of T1D patients and healthy controls are shown in table 2. The DQB1*03:01/05:01 (p=0.03) genotype was significantly higher in female T1D patients than males. Although HLA-DRB1*03:01/04:01 and HLA-DQB1*02:01/03:02 genotypes were more frequent in male patients compared to females, but their differences were not significant.

Distribution of susceptible and protective HLA-DRB1 and -DQB1 haplotypes according to the gender of T1D are as follows

HLA-DRB1*04:01-DQB1*03:02 (p: 0.03) haplotype was significantly more frequent

in male T1D patients than female. HLA-DRB1*03:01-DQB1*02:01 (p=0.02; OR: 0.241) and DRB1*15:01-DQB1*06:01 (p=0.01) haplotypes were significantly higher in female T1D patients than males. These results are summarized in table 3.

Distribution of susceptible and protective HLA-DRB1 and -DQB1 alleles according to the gender of T1D in four age categories at onset groups are as follows

The HLA-DRB1*04:01 (p=0.03; OR: 17) and HLA-DQB1*03:02 (p=0.001) alleles were significantly more frequent in male T1D patients 1-5 years old at onset than females with similar condition. HLA-DRB1*03:01 (p=0.02; OR: 0.194) and HLA-DRB1*15:01 (p=0.03) alleles were significantly higher in female T1D patients 6-10 and 16-20 years old at onset, respectively, than males with similar conditions (Table 4).

Table 2: Distribution of susceptible and protective HLA-DRB1 and -DQB1 genotypes according to the gender of T1D

Genotype	Gender	T1D	Control	P	OR (CL)
HLA-DRB1*0301/0401	M	12	0	0.1	-
	F	1	1		
HLA-DQB1*0201/0302	M	11	0	0.2	-
	F	3	1		
HLA-DQB1*0301/0501	M	0	12	0.03	-
	F	4	7		
HLA-DQB1*0301/0601	M	0	10	0.3	-
	F	1	4		

M; Male, F; Female, NS; Not significant and P; Value of Fisher's exact test.

Table 3: Distribution of susceptible and protective HLA-DRB1 and -DQB1 haplotypes according to the gender of T1D

Haplotype	Gender	T1D	Control	P	OR (CL)
DRB1*0401-DQB1*0302	M	31	0	0.03	-
	F	0	1		
DRB1*0301-DQB1*0201	M	18	11	0.02	0.241 (0.072-0.8)
	F	34	5		
DRB1*1501-DQB1*0601	M	0	15	0.01	-
	F	4	6		

M; Male, F; Female, NS; Not significant and P; Value of Fisher's exact test.

Distribution of susceptible and protective HLA-DRB1 and -DQB1 genotypes and haplotypes based on the gender of T1D in four age categories at onset groups are as follows

HLA-DQB1*02:01/03:02 genotype was more frequent in female patients compared to males,

but the difference was not significant. HLA-DRB1*03:01-DQB1*02:01 ($p=0.007$; OR: 0.121) and DRB1*15:01-DQB1*06:01 ($p=0.01$) haplotypes were significantly higher in female T1D patients 6-10 and 16-20 years old at onset groups, respectively, than males with similar conditions (Table 5).

Table 4: Distribution of susceptible and protective HLA-DRB1 and -DQB1 alleles according to the gender of T1D in four age categories at onset groups

Allele	Gender	Age categories at onset												Control
		1-5	P	OR (CL)	6-10	P	OR (CL)	11-15	P	OR (CL)	16-20	P	OR (CL)	
DRB1*0401	M (n:31)	17	0.035	17 (1.2-223.1)	6	0.2	-	8	0.06	-	0	NS	-	3
	F (n:2)	1			1			0			0			3
DRB1*0301	M (n:18)	7	NS	-	5	0.02	0.194 (0.04-0.77)	10	NS	-	0	NS	-	12
	F (n:34)	8			15			7			0			7
DRB1*1501	M (n:0)	0	NS	-	0	NS	-	0	NS	-	0	0.03	-	17
	F (n:4)	0			0			0			4			11
DQB1*0302	M (n:46)	17	0.001	-	12	0.08	-	12	0.2	-	5	0.1	-	4
	F (n:11)	0			3			17			1			6
DQB1*0601	M (n:1)	0	NS	-	0	NS	-	0	0.1	-	1	0.056	-	21
	F (n:6)	0			0			2			4			9

M; Male, F; Female, NS; Not significant and P; Value of Fisher's exact test.

Table 5: Distributions of susceptible and protective HLA-DRB1 and -DQB1 genotypes and haplotypes according to the gender of T1D in four age at onset groups

Genotype/haplotype	Gender	Age categories at onset												Control
		1-5	P	OR (CL)	6-10	P	OR (CL)	11-15	P	OR (CL)	16-20	P	OR (CL)	
Genotype DQB1*0301/0501	M (n:0)	0	NS	-	0	NS	-	0	0.4	-	0	0.07	-	12
	F (n:4)	0			0			4			3			7
Haplotype DRB1*0401-DQB1*0302	M (n:30)	13	0.07	-	8	0.1	-	10	0.9	-	0	NS	-	0
	F (n:1)	0			0			0			0			1
DRB1*0301-DQB1*0201	M (n:18)	4	0.1	-	4	0.007	0.121(0.02-0.55)	10	0.3	-	0	NS	-	11
	F (n:34)	8			15			4			0			5
DRB1*1501-DQB1*0601	M (n:0)	0	NS	-	0	NS	-	0	NS	-	0	0.01	-	15
	F (n:4)	0			0			0			4			6

M; Male, F; Female, NS; Not significant and P; Value of Fisher's exact test.

Discussion

In this study, we investigated the distribution of susceptible and protective *HLA-DRB1* and *-DQB1* alleles, genotypes and haplotypes based on the gender to find gender-dependent HLA heterogeneity in Iranian T1D patients.

Type 1 diabetes mellitus is a multifactorial and polygenic disease with concordance rate of 30-50% in twins, indicating the significance of genetic factors (26, 27). Among various genes in HLA region, *HLA-DRB* and *-DQB* (HLA class II) contribute the strongest risk to T1D (28, 29). However, differences in *HLA-DRB1* and *-DQB1* haplotypes have been associated with T1D in different population. Exact mechanism of susceptibility to T1D still remains unclear.

In current study, it was found out that *HLA-DRB1*04:01* and *DQB1*03:02* alleles and *DRB1*04:01-DQB1*03:02* haplotype were significantly more frequent in male patients than females. Chan et al. already showed that frequency of *DRB1*04* was significantly higher in male T1D patients than females (22). Tait et al. reported that *DR3-/DR4+* genotype was higher in male patients than female group, which is consistent with our results (30). Another study observed that *DRB1*03:01/04:01* genotype was mainly in male T1D group (22). We demonstrated that *DRB1*03:01/04:01* genotype was more frequent in male patients than females, but it was not significant. Chen et al. reported *DQB1*02:01* allele was higher in male T1D group than females (20), whereas *DQB1*02:01* allele was not significantly more frequent in Iranian male T1D patients. Also, there was observed that *HLA-DRB1*03:01*, *15:01* and *DQB1*06:01* alleles, *DRB1*03:01/05:01* genotype, *DRB1*03:01-DQB1*02:01* and *DRB1*15:01-DQB1*06:01* haplotypes were significantly higher in female T1D patients than male. Similar to our data, most studies demonstrated *DRB1*0301* allele was more frequent in female T1D patients than male (22, 30). In Korean population, it was observed that *DRB1*03:01/09:01* was mainly in female T1D group (22), but as *DRB1*03:01/09:01* genotype was rare among Iranian population, we did not obtain significant any association be-

tween *DRB1*03:02/09:01* genotype and gender in T1D patients. A study in Sweden by Graham et al. 1999 found that *DQB1*06:02* was more negatively associated with T1D in females than males, while we demonstrated that *DQB1*06:01* was more frequent in female T1D group than males (23).

We observed significant differences in age categories at onset in susceptible and protective alleles, genotypes and haplotypes of *DRB1* and *DQB1* stratified by the associated gender. We showed a gender difference in T1D patients which were diagnosed in category of 1-5 years old at onset, with an excess of males in the *DRB1*04:01* and *DQB1*03:02* groups. Also, frequencies of *DRB1*03:01* allele and *DRB1*03:01-DQB1*02:01* haplotype in female patients diagnosed in category of 6-10 years old were significantly more than male patients. Similar to our results, Tait et al. showed a gender difference in T1D patients diagnosed at age <12 years with an excess of males in *DR3-/DR4+* group and females in *DR3+/DR4-* group (30). Chan et al. demonstrated that male T1D patients were significantly higher in the frequency of *DRB1*03:01/04:01* than female with an inverse relation to the age at onset, while *DRB1*03:01/09:01* was observed more frequent in female T1D patients than males with an inverse relation to the age at onset (22). Finding whether gender influence on susceptibility to or protection against T1D, can help to better flow up the T1D patients according their sex category. Also, knowing the gender dependent HLA heterogeneity among T1D patients indicates the influence of sex dependent factor, as the hormones, on the functional effects of the HLA molecules. The individuals carrying alleles which are associated with younger age at onset in each female or male group should take care under preventive treatment.

Conclusion

We can conclude that there is gender dependent HLA genetic heterogeneity of type 1 diabetes in Iranian patients. Also, distribution of *HLA-DRB1* and *-DQB1* alleles, genotypes and haplotypes according to the gender are significantly different in

diagnosed age at onset.

Acknowledgements

This work was supported by a grant from the Tarbiat Modares University as a Ph.D. thesis research program. We gratefully acknowledge the T1D patients and the healthy volunteers participating in this study. The authors declare that they have no conflict of interest in this article.

References

- Atkinson MA, Maclaren NK. The pathogenesis of insulin-dependent diabetes mellitus. *N Engl J Med*. 1994; 331(21): 1428-1436.
- Barnett AH, Eff C, Leslie RD, Pyke DA. Diabetes in identical twins. *Diabetologia*. 1981; 20(2): 87-93.
- Deschamps I, Lestrade H, Busson M, Hors J. Evaluation of recurrence risk in siblings of diabetic children: importance of age and birth order in relation to HLA genotypes. *Diabetes Res*. 1984; 1(3): 125-130.
- Bartsocas CS, Gerasimidi-Vazeou A. Genetics of type 1 diabetes mellitus. *Pediatr Endocrinol Rev*. 2006; 3 Suppl 3: 508-513.
- Steenkiste A, Valdes AM, Feolo M, Hoffman D, Concannon P, Noble J, et al. 14th International HLA and Immunogenetics Workshop: report on the HLA component of type 1 diabetes. *Tissue Antigens*. 2007; 69 Suppl 1: 214-225.
- Singal DP, Blajchman MA. Histocompatibility (HL-A) antigens, lymphocytotoxic antibodies and tissue antibodies in patients with diabetes mellitus. *Diabetes*. 1973; 22(6): 429-432.
- Ikegami H, Fujisawa T, Kawabata Y, Noso S, Oghara T. Genetics of type 1 diabetes: similarities and differences between Asian and Caucasian populations. *Ann N Y Acad Sci*. 2006; 1079: 51-59.
- Ionescu-Tirgoviste C, Guja C, Herr M, Cucca E, Welsh K, Bunce M, et al. Low frequency of HLA DRB1*03-DQB1*02 and DQB1*0302 haplotypes in Romania is consistent with the country's low incidence of Type 1 diabetes. *Diabetologia*. 2001; 44 Suppl 3: B60-66.
- Djoulah S, Busson M, Sasazuki T, Maillere B, Yasunaga S, Kimura A, et al. A new predictive model for insulin dependent diabetes mellitus susceptibility based on combinations of molecular HLA-DRB1 and HLA-DQB1 pockets. *Tissue Antigens*. 1999; 54(4): 341-348.
- Undlien DE, Friede T, Rammensee HG, Joner G, Dahl-Jørgensen K, Søvik O, et al. HLA-encoded genetic predisposition in IDDM: DR4 subtypes may be associated with different degrees of protection. *Diabetes*. 1997; 46(1): 143-149.
- Sheely MJ, Scharf SJ, Rowe JR, Neme de Gimenez MH, Meske LM, Erlich HA, et al. A diabetes-susceptible HLA haplotype is best defined by a combination of HLA-DR and -DQ alleles. *J Clin Invest*. 1989; 83(3): 830-835.
- Erlich HA, Zeidler A, Chang J, Shaw S, Raffel LJ, Klitz W, et al. HLA class II alleles and susceptibility and resistance to insulin-dependent diabetes mellitus in Mexican-American families. *Nat Genet*. 1993; 3(4): 358-364.
- Cucca F, Muntoni F, Lampis R, Frau F, Argiolas L, Silveti M, et al. Combinations of specific DRB1, DQA1, DQB1 haplotypes are associated with insulin-dependent diabetes mellitus in Sardinia. *Hum Immunol*. 1993; 37(2): 85-94.
- Ronnigen KS, Keiding N, Green A. Correlations between the incidence of childhood-onset type 1 diabetes in Europe and HLA genotypes. *Diabetologia*. 2001; 44 Suppl 3: B51-59.
- Thorsby E, Ronningen KS. Particular HLA DQ molecules play a dominant role in determining susceptibility or resistance to type 1 (insulin-dependent) diabetes mellitus. *Diabetologia*. 1993; 36(5): 371-377.
- Petrone A, Bugawan TL, Mesturino CA, Nisticò L, Galgani A, Giorgi G, et al. The distribution of HLA class II susceptible/protective haplotypes could partially explain the low incidence of type 1 diabetes in Continental Italy (Lazio region). *Tissue Antigens*. 2001; 58(6): 385-394.
- Gorodezky C, Alaez C, Murguía A, Rodríguez A, Baladares S, Vazquez M, et al. HLA and autoimmune diseases: Type 1 diabetes (T1D) as an example. *Autoimmun Rev*. 2006; 5(3): 187-194.
- Nejentsev S, Koskinen S, Sjoroos M, Reijonen H, Schwartz EI, Kovalchuk L, et al. Distribution of insulin-dependent diabetes mellitus (IDDM) - related HLA alleles correlates with the difference in IDDM incidence in four populations of the Eastern Baltic region. *Tissue Antigens*. 1998; 52(5): 473-477.
- Park Y. Why is type 1 diabetes uncommon in Asia?. *Ann N Y Acad Sci*. 2006; 1079: 31-40.
- Chen BH, Chiang CH, Lin SR, Chao MG, Tsai ST. The influence of age at onset and gender on the HLA-DQA1, DQB1 association in Chinese children with insulin dependent diabetes mellitus. *Human Immunol*. 1999; 60(11): 1131-1137.
- Caillat-Zucman S, Garchon HJ, Timsit J, Assan R, Boitard C, Djilali-Saiah I, et al. Age-dependent HLA genetic heterogeneity of type 1 insulin-dependent diabetes mellitus. *J Clin Invest*. 1992; 90(6): 2242-2250.
- Chan SH, Thai AC, Lin YN, Liu KF, Wee GB. Influence of gender and age at onset on the HLA association in Chinese with insulin-dependent diabetes mellitus. *Hum Immunol*. 1995; 44(3): 175-180.
- Graham J, Kockum I, Sanjeevi CB, Landin-Olsson M, Nystrom L, Sundkvist G, et al. Negative association between type 1 diabetes and HLA DQB1*0602-DQA1*0102 is attenuated with age at onset. *Eur J Immunogenet*. 1999; 26(2-3): 117-127.
- Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res*. 1998; 16(3): 1215.
- Sayad A, Akbari MT, Pajouhi M, Mostafavi F, Zamani M. The influence of the HLA-DRB1, HLA-DQB1 and polymorphic positions of the HLA-DRβ1 and HLA-DQβ1 molecules on risk of Iranian type 1 diabetes mellitus patients. *Int J Immunogenet*. 2012; 39(5): 429-436.
- Sanjeevi CB. Immunology of diabetes society congress. *Immunology of diabetes: autoimmune mecha-*

- nisms and the prevention and cure of type 1 diabetes. 1st ed. USA: Annals of The NY Academy of Sciences; 2002; 958.
27. Dorman JS, Bunker CH. HLA-DQ locus of the human leukocyte antigen complex and type 1 diabetes mellitus. A HUGE review. *Epidemiol Rev.* 2000; 22(2): 218-227.
 28. Zamani M, Pociot F, Spaepen M, Raeymaekers P, Nerup J, Cassiman JJ. Linkage and association of the HLA gene complex with IDDM in 81 Danish families: strong linkage between DR1571+ and IDDM. *J Med Genet.* 1996; 33(11): 899-905.
 29. Zamani M, Cassiman JJ. Reevaluation of the importance of polymorphic HLA class II alleles and amino acids in the susceptibility of individuals of different populations to type I diabetes. *Am J Med Genet.* 1998; 76(2): 183-194.
 30. Tai BD, Harrison LC, Drummond BP, Stewart V, Varney MD, Honeyman MC. HLA antigens and age at diagnosis of insulin-dependent diabetes mellitus. *Hum Immunol.* 1995; 42(2): 116-122.
-

Archive of SID