

Suppression of Gelatinase Activity in Human Peripheral Blood Mononuclear Cells by Verapamil

Fatemeh Hajighasemi, Ph.D.*, Neda Kakadezfuli, M.Sc.

Department of Immunology, Faculty of Medicine, Shahed University, Tehran, Iran

*Corresponding Address: P.O.Box: 14155-7435, Department of Immunology, Faculty of Medicine, Shahed University, No. 31, Shahid Abdollahzadeh Street, Keshavarz Blvd., 1415635111, Tehran, Iran

Email: resoome@yahoo.com

Received: 02/Jan/2013, Accepted: 16/Mar/2013

Abstract

Objective: Gelatinases are a large group of proteolytic enzymes that belong to the matrix metalloproteinases (MMPs). MMPs are a broad family of peptidases, which proteolyse the extracellular matrix and have an important role in inflammation. Verapamil is a calcium channel blocker extensively used in the treatment of numerous cardiovascular diseases such as arrhythmia and hypertension. The anti-tumor and anti-inflammatory effects of verapamil have also been shown. In this study, the effect of verapamil on gelatinase activity in human peripheral blood mononuclear cells (PBMCs) has been assessed *in vitro*.

Materials and Methods: In this experimental study, PBMCs from healthy adult volunteers were isolated by ficoll-hypaque-gradient centrifugation. The cells were then cultured in complete RPMI-1640 medium and after that incubated with different concentrations of verapamil (0–200 μ M) in the presence or absence of phytohemagglutinin (PHA) (10 μ g/ml) for 48 hours. The gelatinase A (MMP-2)/gelatinase B (MMP-9) activity in cell-conditioned media was then evaluated by gelatin zymography. Statistical comparisons between groups were made by analysis of variance (ANOVA).

Results: Verapamil significantly decreased the MMP-2/MMP-9 activity in human PBMCs after 48 hours incubation time compared with untreated control cells. The association was dose-dependent.

Conclusion: In this study verapamil exhibited a dose-dependent inhibitory effect on gelatinase A and gelatinase B activity in human PBMCs. It seems that the anti-inflammatory properties of verapamil may be in part due to its inhibitory effects on gelatinase activity. Regarding the beneficial effects of MMPs- inhibitors in the treatment of some cardiovascular diseases, the positive effect of verapamil on such diseases may be in part due to its anti-MMP activity. Verapamil with its inhibitory effects on gelatinases activity may be a useful MMP-inhibitor. Given the beneficial effect of MMP-inhibitors in some cancerous, inflammatory and autoimmune disorders, it seems likely that verapamil could also be used to treat these diseases.

Keywords: Verapamil, Gelatinase, Mononuclear Cells

Cell Journal(Yakhteh), Vol 16, No 1, Spring 2014, Pages: 11-16

Citation: Hajighasemi F, Kakadezfuli N. Suppression of gelatinase activity in human peripheral blood mononuclear cells by verapamil. Cell J. 2014; 16(1): 11-16.

Introduction

Gelatinases are a family of proteolytic enzymes belonging to the matrix metalloproteinases (MMPs) (1). MMPs are a group of zinc-dependent enzymes, which proteolyse the extracellular matrix (2, 3). Gelatinases degrade collagen type IV and V and are divided into gelatinase A (MMP-2) and gelatinase B (MMP-9) with molecular weights of 72 and 92 KD, respectively (4, 5). Gelatinases play an important role in inflammation, autoimmunity, cancer progression and metastasis (6-9). Verapamil is a calcium (Ca^{2+}) channel blocker extensively used in the treatment of numerous cardiovascular diseases, such as arrhythmia and hypertension (10, 11). Anti-tumor and anti-inflammatory effects of verapamil have also been shown (12, 13). The inhibitory effect of verapamil on lipopolysaccharide (LPS)-induced pro-inflammatory cytokine production and NF- κ B activation has been shown *in vivo* (14, 15), and the therapeutic properties of a Ca^{2+} channel blocker in inflammatory bowel disease have been described (16). In addition, down regulation of proinflammatory factors such as superoxide and nitric oxide (NO) by verapamil through a Ca^{2+} channel-independent pathway (13) and the inhibitory effect of verapamil on MMP-9 activity in murine mammary tumor cells has been reported (12).

Mononuclear cells play an important role in inflammation (17, 18) through several mechanisms such as regulating the extracellular turnover. This occurs via the production of a number of mediators such as inflammatory cytokines and MMPs (19-21). Production of gelatinases by peripheral blood mononuclear cells (PBMCs) has also been shown (22). Given the anti-inflammatory effects of verapamil and the important role of mononuclear cells and MMPs in inflammation, in this study we assessed the effect of verapamil on gelatinase (MMP-2 and MMP-9) activity in human PBMCs.

Materials and Methods

This experimental study was approved by The Deputy Director of Research in the Faculty of Medicine at Shahed University.

Reagents

RPMI-1640 medium, penicillin, streptomycin, PHA (phytohemagglutinin) and trypan blue (TB) were obtained from Sigma (USA). MTT (3-[4,

5-dimethyl thiazol-2,5-diphenyltetrazolium bromide]) was purchased from Merck (Germany). Fetal calf serum (FCS) was from Gibco (USA). Verapamil was purchased from Sobhandarou Pvt. Co. Ltd (Tehran, Iran). Microtiter plates, flasks and tubes were from Nunc (Falcon, USA).

Preparation of verapamil

Verapamil was dissolved in distilled water and stored as a stock at -20°C until use. The stock was diluted in culture medium in order to prepare appropriate concentrations before use.

Peripheral blood mononuclear cells isolation

PBMCs from the venous blood of healthy adult volunteers were isolated by ficoll-hypaque-gradient centrifugation. Subsequently, the cells were washed three times in phosphate buffer saline (PBS). The cells were then resuspended in RPMI-1640 medium supplemented with 10% FCS and were incubated in 5% CO_2 at 37°C .

Cell culture and treatment

The method used for cell culture and treatment has been described in detail previously (23). Briefly, human PBMCs were cultured in RPMI-1640 medium supplemented with 10% FCS, penicillin (100 IU/ml) and streptomycin (100 $\mu\text{g}/\text{ml}$) at 37°C in 5% CO_2 . The cells were seeded at a density of 1×10^6 cells/ml and then treated with different concentrations of Verapamil (0-200 μM) in the presence of PHA (10 $\mu\text{g}/\text{ml}$) for 48 hours. Afterward the supernatants from the cell cultures were collected, centrifuged and stored at -20°C for subsequent tests. All experiments were done in triplicate.

Evaluation of MMP-2 and MMP-9 activity by gelatin zymography

MMP-2 and MMP-9 activity in cell-conditioned media were evaluated using the gelatin zymography technique according to the modified Kleiner and Stetler-Stevenson method (1994, 24) as previously described (25). Briefly, cell culture supernatants were subjected to SDS-PAGE on 10% polyacrylamide gel copolymerized with 2 mg/ml gelatin in the presence of 0.1% SDS under non-reducing conditions at a constant voltage of 80 V for three hours. After electrophoresis,

the gels were washed in 2.5% Triton X-100 for one hour to remove the SDS and then incubated in a buffer containing 0.1 M Tris-HCl, pH=7.4 and 10 mM CaCl_2 at 37°C overnight. Afterwards, the gels were stained with 0.5% Coomassie brilliant blue (Coomassie blue dissolved in 40% ethanol, 10% acetic acid) for 1 hour and then destained. Proteolytic enzyme activity was detected as clear bands against a blue background indicating lysis of gelatin. The supernatants from serum-free cultured HT1080 cells obtained from NCBI (National Cell Bank of Iran, Pasteur Institute of Iran, Tehran) were used as a molecular weight marker for MMP-2 and MMP-9 as described before (26).

The relative intensity of the gelatin lysis bands compared to the control was measured using UVI Pro gel documentation system (Vilber Lourmat, Marne-la-Vallee Cedex 1, France) and expressed as relative gelatinolytic activity.

Statistical analysis

MMP-2 and MMP-9 activity measurement in cell-conditioned media was performed in three independent experiments and the results were expressed as mean \pm SEM. Statistical comparisons between groups were made by analysis of variance (ANOVA). $P < 0.05$ was considered significant. Multiple comparisons were tested using the Tukey method (5%) for statisti-

cally significant differences. The software SPSS 11.5 and Excel 2003 were used for statistical analysis and graph making respectively.

Results

Effect of verapamil on gelatinase-A (MMP-2) and gelatinase-B (MMP-9) activity in human PBMCs in different concentrations are shown in figures 1 (A, B) and 2 (A, B).

Verapamil effect on gelatinase-A (MMP-2) activity in PHA-stimulated human PBMCs

Verapamil significantly decreased the gelatinase-A (MMP-2) activity in PHA-stimulated human PBMCs in a dose-dependent fashion after 48 hours incubation compared with untreated control cells (Fig 1A, B). The decrease in gelatinase-A activity was shown at 200 μM concentration of verapamil.

Verapamil effect on gelatinase-B (MMP-9) activity in PHA-stimulated human PBMCs

Verapamil significantly decreased the gelatinase-B (MMP-9) activity in PHA-stimulated human PBMCs in a dose-dependent fashion after 48 hours incubation time compared with untreated control cells (Fig 2A, B). The decrease in gelatinase-B activity was seen at a verapamil concentration of 200 μM .




Fig 1: Effect of verapamil on MMP-2 activity in PHA-stimulated human PBMCs. The human PBMCs (1×10^6 cells/ml), were treated with different concentrations of verapamil (0-200 μM) in the presence of PHA (10 $\mu\text{g}/\text{ml}$) for 48 hours. At the end of treatment, MMP-2 activity in conditioned medium was measured by gelatin zymography. A. Zymogram of MMP-2 activity in human PBMCs treated with verapamil. Lane 1 represents untreated human PBMCs. Lanes 2 to 4 represent verapamil at 20, 100 and 200 μM concentrations respectively. Lane 5 represents the control. B. MMP-2 activity in human PBMCs, was measured by scanning the zymograms and by densitometric analysis of the MMP-2 bands. Data are represented as the mean \pm SEM of the three independent experiments.

*, $P < 0.05$ was considered significant.




Fig 2: Effect of verapamil on MMP-9 activity in PHA-stimulated human PBMCs. Human PBMCs (1×10^6 cells/ml), were treated with different concentrations of verapamil (0-200 μ M) in the presence of PHA (10 μ g/ml) for 48 hours. At the end of treatment, MMP-9 activity in the conditioned medium was measured by gelatin zymography. **A.** Zymogram of MMP-9 activity in human PBMCs treated with verapamil. Lane 1 represents untreated human PBMCs. Lanes 2 to 4 represent verapamil at 20, 100 and 200 μ M concentrations respectively. Lane 5 represents the control. **B.** MMP-9 activity in human PBMCs, was measured by scanning the zymograms and by densitometric analysis of the MMP-9 bands. Data are mean \pm SEM of three independent experiments.

*; $P < 0.05$ was considered significant.

Discussion

In this study, a 200 μ M concentration of verapamil inhibited the gelatinase A (MMP-2) and gelatinase B (MMP-9) activity in human PBMCs. These results are consistent with the study by Farías, et al. (12) in which verapamil suppressed the MMP-9 activity in murine mammary tumor cells. In the Farías, et al. study a 50 μ M concentration of verapamil decreased the MMP-9 activity, while in our study the decrease in MMP-9 activity was seen at 200 μ M. The discrepancy between our results and those of Farías, et al. may in part be due to the type and origin of the cells used. The cells used by Farías et al. were mouse mammary tumor cells while we used human PBMCs which are normal cells. Thus it seems that verapamil exerts an inhibitory effect on MMP activity in tumor cells at a much lower concentration relative to normal cells. Further experiments to verify this hypothesis are warranted.

The anti-inflammatory effects of verapamil have been reported by several investigators (14, 15, 27, 28). For example in the study by Matsomuri, et al. (14) verapamil reduced the production of inflammatory cytokines (IFN- γ , TNF α and IL-1 β) in PBMCs and in the study by Das, et al. (27) verapamil suppressed the recall of peritoneal macrophages in response to tiogliclate. A verapamil induced decrease in the serum levels of TNF- α and IFN- γ in septic shock mice has been also reported (28). As

gelatinases play an important role in inflammation (6), the anti-inflammatory properties of verapamil may be in part due to its inhibitory effects on gelatinase activity. The inhibitory effects of verapamil on LPS-induced secretion of pro-inflammatory cytokines and NF-Kappa B activation has been shown *in vivo* (14, 15). Suppression of the LPS-induced secretion of pro-inflammatory factors such as superoxide and nitric oxide (NO) by verapamil through a calcium channel-independent pathway has been shown (13). Inflammatory factors such as cytokines and prostaglandin E (2) are important regulators of MMPs activity (29, 30). Thus inhibitory effect of verapamil on MMP activity shown in this study may in part be due to its suppressive effect on inflammatory mediators.

The potential implication of anti-MMPs in the treatment of ischemic heart failure has been reported (31) and the positive effect of verapamil on ischemic heart failure (32) may in part be due to its anti-MMP activity. It should be noted that in our study, the concentration of verapamil which inhibited gelatinase activity *in vitro*, was higher than that usually used in cardiovascular patients.

In our previous study, verapamil showed a significant cytotoxic effect against human PBMCs at ≥ 1000 μ M concentrations (33). Therefore a decrease in gelatinase activity at a verapamil concentration of 200 μ M is not due to its cytotoxic effect and other mechanism(s) may be involved in this

process.

Yue et al. (34) in a study of calcium channel blockers, showed that nifedipine increased and amlodipine decreased the MMP-2 activity in rat fibroblasts, while verapamil and diltiazem had no effect on MMP activity. The difference between our results and the Yue, et al. study may in part be due to differences between the lines and origin of the cells used. As already discussed, we used human PBMCs. However, the study by Yue, et al. used rat fibroblasts and another study performed by Yue et al. (35) in which verapamil had no effect on MMP-2 activity, the cells used were rat vascular smooth muscle cells.

According to our review of the literature this is the first report documenting the inhibitory effect of verapamil on gelatinase activity in human PBMCs. As gelatinases are important mediators of cancer and inflammation (6, 36), the anti-tumoral and anti-inflammatory effects of verapamil reported by other studies (12, 13) may in part be due to its inhibitory effects on gelatinase activity.

Given the important role of inflammation in cardiovascular diseases (37) the therapeutic effect of verapamil in cardiac disease (32) may in part be due to its anti-inflammatory properties mediated by inhibition of MMP(s)-activity.

Taken together our results suggest that verapamil, along with its long-term usage in cardiovascular diseases, may have potential implication for the development of MMP-inhibitors. Further studies of the effect of verapamil on gelatinase activity in other cell lines as well as inflammatory and autoimmune diseases are warranted.

Conclusion

In this study, verapamil showed inhibitory effects on gelatinase (MMP-2/MMP-9) activity in human PBMCs. Thus verapamil may be of potential use in the preparation of MMP-inhibitors. MMPs have important role in inflammation, so verapamil, along with its long-term usage in cardiovascular disease, may be a good candidate for the development of anti-inflammatory agents.

Acknowledgments

This study was financially supported by Shahed University. The authors declare no con-

flict of interests in relation to this article.

References

1. Fink K, Boratyński J. The role of metalloproteinases in modification of extracellular matrix in invasive tumor growth, metastasis and angiogenesis. *Postepy Hig Med Dosw (Online)*. 2012; 66: 609-628.
2. Xie XY, Yang C, Ren M, Hao SY, Zhu P, Yan L. Inhibition of matrix metalloproteinase 9 expression in rat dermal fibroblasts using small interfering RNA. *J Am Podiatr Med Assoc*. 2012; 102(4): 299-308.
3. Vandembroucke RE, Dejonckheere E, Van Lint P, De-meestere D, Van Wonterghem E, Vanlaere I, et al. Matrix metalloprotease 8-dependent extracellular matrix cleavage at the blood-CSF barrier contributes to lethality during systemic inflammatory diseases. *J Neurosci*. 2012; 32(29): 9805-9816.
4. Sillat T, Saat R, Pöllänen R, Hukkanen M, Takagi M, Kontinen YT. Basement membrane collagen type IV expression by human mesenchymal stem cells during adipogenic differentiation. *J Cell Mol Med*. 2012; 16(7): 1485-1495.
5. Legrand C, Gilles C, Zahm JM, Polette M, Buisson AC, Kaplan H, et al. Airway epithelial cell migration dynamics. MMP-9 role in cell-extracellular matrix remodeling. *J Cell Biol*. 1999; 146(2): 517-529.
6. Lagente V, Boichot E. Role of matrix metalloproteinases in the inflammatory process of respiratory diseases. *J Mol Cell Cardiol*. 2010; 48(3): 440-444.
7. Nakagawa P, Liu Y, Liao TD, Chen X, González GE, Bobbitt KR, et al. Treatment with N-acetyl-seryl-aspartyl-lysyl-proline prevents experimental autoimmune myocarditis in rats. *Am J Physiol Heart Circ Physiol*. 2012; 303(9): H1114-1127.
8. Durlík M, Gardian K. Metalloproteinase 2 and 9 activity in the development of pancreatic cancer. *Pol Przegl Chir*. 2012; 84(8): 377-382.
9. Wang N, Zhu M, Tsao SW, Man K, Zhang Z, Feng Y. Up-regulation of TIMP-1 by genipin inhibits MMP-2 activities and suppresses the metastatic potential of human hepatocellular carcinoma. *PLoS One*. 2012; 7(9): e46318.
10. Iskenderov BG, Sisina ON, Burmistrova LF. Selection of rational combinations of indapamide with various of calcium antagonists in patients with arterial hypertension. *Kardiologia*. 2011; 51(4): 22-27.
11. Hegazi MO, Aldabie G, Al-Mutairi S, El Sayed A. Junctional bradycardia with verapamil in renal failure-care required even with mild hyperkalaemia. *J Clin Pharm Ther*. 2012; 37(6): 726-728.
12. Farías EF, Aguirre Ghiso JA, Ladeda V, Bal de Kier Joffé E. Verapamil inhibits tumor protease production, local invasion and metastasis development in murine carcinoma cells. *Int J Cancer*. 1998; 78(6): 727-734.
13. Liu Y, Lo YC, Qian L, Crews FT, Wilson B, Chen HL, et al. Verapamil protects dopaminergic neuron damage through a novel anti-inflammatory mechanism by inhibition of microglial activation. *Neuropharmacology*. 2011; 60(2-3): 373-380.
14. Matsumori A, Nishio R, Nose Y. Calcium channel blockers differentially modulate cytokine production by peripheral blood mononuclear cells. *Circ J*. 2010; 74(3): 567-571.
15. Li G, Qi XP, Wu XY, Liu FK, Xu Z, Chen C, et al. Verapamil modulates LPS-induced cytokine production via inhibition of NF-kappa B activation in the liver. *Inflamm Res*. 2006; 55(3): 108-113.
16. Di Sabatino A, Rovedatti L, Kaur R, Spencer JP, Brown JT, Morisset VD, et al. Targeting gut T cell Ca2+ release-activated Ca2+ channels inhibits T cell cytokine production

- and T-box transcription factor T-bet in inflammatory bowel disease. *J Immunol.* 2009; 183(5): 3454-3462.
17. Shan J, Ma JM, Wang R, Liu QL, Fan Y. Proliferation and apoptosis of peripheral blood mononuclear cells in patients with oral lichen planus. *Inflammation.* 2012 ; 36(2): 419-425.
 18. Lin WH, Wu CR, Lee HZ, Kuo YH, Wen HS, Lin TY, et al. Induced apoptosis of Th2 lymphocytes and inhibition of airway hyperresponsiveness and inflammation by combined lactic acid bacteria treatment. *Int Immunopharmacol.* 2013; 15(4): 703-711.
 19. Li H, Li L, Min J, Yang H, Xu X, Yuan Y, et al. Levels of metalloproteinase (MMP-3, MMP-9), NF-kappaB ligand (RANKL), and nitric oxide (NO) in peripheral blood of osteoarthritis (OA) patients. *Clin Lab.* 2012; 58(7-8): 755-762.
 20. Garcia-Rodriguez S, Callejas-Rubio JL, Ortego-Centeno N, Zumaquero E, Ríos-Fernandez R, Arias-Santiago S, et al. MAPK1 gene expression on peripheral blood mononuclear cells and correlation with T-helper-transcription factors in systemic lupus erythematosus patients. *Mediators Inflamm.* 2012; 2012: 495934.
 21. Bayard F, Godon O, Nalpas B, Costentin C, Zhu R, Sossan P, et al. T-cell responses to hepatitis B splice-generated protein of hepatitis B virus and inflammatory cytokines/chemokines in chronic hepatitis B patients. ANRS study: HB EP 02 HBSP-FIBRO. *J Viral Hepat.* 2012; 19(12): 872-880.
 22. Hajjighasemi F, Hajjighasemi S. Zymographic analysis of gelatinases patterns in peripheral blood mononuclear cells. *J Med Sci.* 2011; 2(6): 904-909.
 23. Hajjighasemi F, Mirshafiey A. Propranolol effect on proliferation and vascular endothelial growth factor secretion in human immunocompetent cells. *J Clin Immunol Immunopathol Res.* 2010; 2(2): 22-27.
 24. Kleiner DE, Stetler-Stevenson WG. Quantitative zymography: detection of picogram quantities of gelatinases. *Anal Biochem.* 1994; 218(2): 325-329.
 25. Hajjighasemi F, Hajjighasemi S. Effect of propranolol on angiogenic factors in human hematopoietic cell lines in vitro. *Iran Biomed J.* 2009; 13(4): 223-228.
 26. Yodkeeree S, Garbisa S, Limtrakul P. Tetrahydrocurcumin inhibits HT1080 cell migration and invasion via down-regulation of MMPs and uPA. *Acta Pharmacol Sin.* 2008; 29(7): 853-860.
 27. Das R, Burke T, Van Wagoner DR, Plow EF. L-Type calcium channel blockers exert an anti-inflammatory effect by suppressing expression of plasminogen receptors on macrophages. *Circ Res.* 2009; 105(2): 167-175.
 28. Wyska E. Pretreatment with R(+)-verapamil significantly reduces mortality and cytokine expression in murine model of septic shock. *Int Immunopharmacol.* 2009; 9(4): 478-490.
 29. Lee SJ, Lee EJ, Kim SK, Jeong P, Cho YH, Yun SJ, et al. Identification of pro-inflammatory cytokines associated with muscle invasive bladder cancer; the roles of IL-5, IL-20, and IL-28A. *PLoS One.* 2012; 7(9): e40267.
 30. Kim YS, Shin SI, Kang KL, Chung JH, Herr Y, Bae WJ, et al. Nicotine and lipopolysaccharide stimulate the production of MMPs and prostaglandin E(2) by hypoxia-inducible factor-1 α up-regulation in human periodontal ligament cells. *J Periodontal Res.* 2012 ; 47(6): 719-728.
 31. Yokoyama U, Ishiwata R, Jin MH, Kato Y, Suzuki O, Jin H, et al. Inhibition of EP4 signaling attenuates aortic aneurysm formation. *PLoS One.* 2012; 7(5): e36724.
 32. Yui H, Imaizumi U, Beppu H, Ito M, Furuya M, Arisaka H, et al. Comparative effects of verapamil, nifedipine, and nitroglycerin on myocardial ischemia/reperfusion injury. *Anesthesiol Res Pract.* 2011; 2011: 521084.
 33. Hajjighasemi F, Kakadezfuli N. Sensitivity of monocytic cells to verapamil in vitro. *Res J Biol Sci.* 2012; 7(5): 209-214.
 34. Yue H, Uzui H, Shimizu H, Nakano A, Mitsuke Y, Ueda T, et al. Different effects of calcium channel blockers on matrix metalloproteinase-2 expression in cultured rat cardiac fibroblasts. *J Cardiovasc Pharmacol.* 2004; 44(2): 223-230.
 35. Yue H, Lee JD, Shimizu H, Uzui H, Mitsuke Y, Ueda T. Effects of magnesium on the production of extracellular matrix metalloproteinases in cultured rat vascular smooth muscle cells. *Atherosclerosis.* 2003; 166(2): 271-277.
 36. Roth M, Eickelberg O, Kohler E, Ernet P, Blockt L H. Ca²⁺ channel blockers modulate metabolism of collagens within the extracellular matrix. *Proc Natl Acad Sci USA.* 1996; 93(11): 5478-5482.
 37. Edsfeldt A, Nitulescu M, Grufman H, Grönberg C, Persson A, Nilsson M, et al. Soluble urokinase plasminogen activator receptor is associated with inflammation in the vulnerable human atherosclerotic plaque. *Stroke.* 2012; 43(12): 3305-3312.