The Effect of Extremely Low Frequency Pulsed Electromagnetic Field on *In Vitro* Fertilization Success Rate in NMRI Mice

Leili Hafizi, M.D.¹, Ameneh Sazgarnia, Ph.D.², Nezhat Mousavifar, M.D.¹, Mohammad Karimi, M.D.^{1*}, Saleh Ghorbani, M.D.¹, Mohammad Reza Kazemi, M.D.¹, Neda Emami Meibodi, B.Sc.³, Golkoo Hosseini, M.D.⁴, Hesam Mostafavi Toroghi, M.D.⁴

1. Department of Obstetrics and Gynecology, Imam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, Iran

 Department of Medical Physics, Mashhad University of Medical Sciences, Mashhad, Iran
Department of Biomedical Engineering, Engineering Faculty, Islamic Azad University, Mashhad, Iran
Biochemistry of Nutrition Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

*Corresponding Address: P.O.Box: 9137913316, Department of Obstetrics and Gynecology, Imam Reza Hospital, Mashhad University of Medical Sciences, Mashhad, Iran

E-mail: Karimim851@mums.ac.ir

Received: 23/Aug/2012, Accepted: 15/Jan/2013

Abstract -

Objective: The effects of exposure to electromagnetic fields (EMF) on reproduction systems have been widely debated. In this study, we aimed to investigate whether low frequency EMF could ameliorate the *in vitro* fertilization success rate in Naval medical research institute (NMRI) Mice.

Materials and Methods: In this randomized comparative animal study, ten NMRI mice were randomly divided into 2 equal groups (control and experimental). 10 IU of human chorionic gonadotropin (hCG) was injected intraperitoneally to both groups in order to stimulate ovulating, and ovums were then aspirated and kept in KSOM (modified version of sequential simplex optimization medium with a higher K⁺ concentration) culture medium. Metaphase II ovums were separated, and sperms obtained by "swim out" method were added to metaphase II ovums in the culture medium. The experimental group was exposed to 1.3 millitesla pulsed electromagnetic field at 4 kilohertz frequency for 5 hours. To assess the efficacy, we considered the identification of two-pronuclear zygote (2PN) under microscope as fertilizing criterion.

Results: Total number of collected ovums in the control and experimental groups was 191 and 173, respectively, from which 58 (30.05%) and 52 (30.36%) ovums were collected from metaphase II, respectively. *In vitro* fertilization (IVF) success rate was 77% in extremely low frequency- pulsed electromagnetic field (ELF-PEMF) for exposed group (experimental), whereas the rate was 68% for control group.

Conclusion: Despite increased percentile of IVF success rate in exposed group, there was no statistically significant difference between 2 groups, but this hypothesis has still been stated as a question. Further studies with larger sample sizes and different EMF designs are suggested.

Keywords: In Vitro Fertilization, Electromagnetic Fields, Mice Cell Journal (Yakhteh), Vol 15, No 4, Winter 2014, Pages: 310-315.

Citation: Hafizi L, Sazgarnia A, Mousavifar N, Karimi M, Ghorbani S, Kazemi MR, Emami Meibodi N, Hosseini G, Mostafavi Toroghi H. The effect of extremely low frequency pulsed electromagnetic field on in vitro fertilization success rate in NMRI mice. Cell J. 2014; 15(4): 310-315.

Introduction

We are surrounded by many various magnetic, electric and electromagnetic fields (EMF). Natural electromagnetic fields are one of the factors maintaining the life in our planet. Pulsed electromagnetic fields (PEMF) stimulate many sub-cellular responses in living systems, but the range of action differs considerably depending on the properties of fields' and types of cells and organisms. Plasma membrane, especially gap junctions and proteins connecting two adjuvant cells, are the most susceptible to such effects (1, 2).

Although electromagnetic field has been subjected to various researches in recent years, it has been considered as a completely diverse and controversial subject, yet. Most of these researches have focused on machine life and effects of many electric instruments around us that mostly use50 Hertz EMF, which is as the target frequency evaluated by these studies. Some studies have insisted on the toxicological aspect of electromagnetic fields on various body systems and organs(3-5) including apoptosis (1, 6, 7), congenital anomalies (2), alteration in ion homeostasis (9), free radicals generation (9-11) and DNA damage (12, 13), while some other researches have considered the therapeutic effects (14-17), especially on bone tissue(17, 18). For example, extremely low frequency (ELF) is routinely applied in non-union factures (19), diabetic ulcer (20) and osteoarthritis treatment (21, 22). In this category, the range of the frequency is almost wide, but mostly below 50 Hertz. Other positive effects of EMF include pain alleviation (acute and chronic) (23), inflammation reduction (24), nerve regeneration (25), improving blood flow (26), enhancement of delivery and effects of medications, angiogenesis (15, 27), decreasing blood glucose and serum cortisol concentration (28) and so on.

There have been extensive researches about the effects of EMF on male reproductive system, but investigations about female reproductive system are scarce. About 15% of couples worldwide fail to give birth to a child (29). Assisted reproductive technology (ART) includes all methods involving laboratory manipulation of gametes (sperm or oo-cytes) and/or embryos for the purposes of repro-

duction. Although many diverse methods have been invented and employed, *in vitro* fertilization (IVF) is the most popular and preferable method in ART. Nowadays, IVF success rate is about 35% with a very short range of changes in different centers (30).

Regarding the potential *in vitro* effects of extremely low frequency- pulsed electromagnetic field (ELF-PEMF) on reproduction, growth and development, we aimed to investigate the effects of ELF-PEMF on IVF success rate in NMRI mice.

Materials and Methods

The study was designed as a randomized comparative animal study.

Animals

Ten female and 2 male Naval Medical Research Institute (NMRI) mice (20 to 25 g) were obtained from the Razi Institute (Mashhad, Iran) and housed under standard laboratory conditions. The maintenance and care of the mice complied with National Institutes of Health (NIH) guidelines for the humane use of laboratory animals. They were kept at constant room temperature $(21 \pm 2^{\circ}C)$ under a normal 12-hour light/12-hour dark cycle with free access to food and water. The study protocol was approved by the Ethics Committee of Research Council of Mashhad University of Medical Sciences. To identify possible effect of PEMF exposure on IVF success rate, female NMRI mice were randomly divided into 2 groups. Five mice were randomly selected for control group and five for experimental group.

During ovulation time, vaginal opening gets wet and its color changes into pink with high folding, so we used this sign to control the synchronization of their ovulation time.

Induction of ovulation

To super ovulate the mice, we injected 10 IU of Pregnant Mare's Serum Gonadotropin (PMSG; Sigma-Aldrich, St. Louis, USA) intraperitoneally, and 48 hours later, we injected 10IU human chorionic gonadotropin (hCG; Organon, Holland) intraperitoneally, as well. Thirteen hours after hCG injection, the mice were made unconscious and their abdomens were surgically opened under sterile conditions. The ovaries and fallopian tubes were dissected and transported into culture media. All ovums were aspirated under sterile conditions, transported into KSOM (Merck, Germany) culture media, and incubated at 37° C temperature with 5% of CO₂.

Microscopic evaluation

In the next step, we put aspirated ovums in hyaluronidasesolution (Sigma-Aldrich, USA) for 1 minute to remove granulosa cell layer. Then, ovums, observed under stereo microscope (Olympus, Japan) for quality control, were counted consequently. Metaphase II ovums were identified by their polar bodies and separated from other ovums.

In vitro fertilization technique accomplishment

Metaphase II ovums were kept in the culture media plate, separately, under aqua paraffin oil at 37°C incubation for 1 hour (31). Meanwhile, to complete the IVF technique, male mice were

killed, a section of their caudaepididymides was cut out, and placed into the culture media.

Afterwards, with punching and gentle pushing, sperms were squeezed out of epididymides and entered the culture media by swim out technique. Then, sperm suspension with the concentration of 5×10^4 sperm/mL was prepared, mixed with metaphase II ovums and transferred to fresh culture media.

In the final step, fertilization culture media of experimental group were placed under emitting of specified ELF-PEMF exposure system, and was then kept in the incubator for 5 hours.

Fertilization culture media of control group was kept in same incubation condition, without companionship of such designed system.

After 5 hours, the status of fertilization was investigated under microscope. The criterion of fertilization was to detect two pronuclei zygotes (2PN). Unfertilized ovums were counted separately (Fig1).



Fig 1: Mouse ovum in different stages of maturation and after inoculation.

Pulsed electromagnetic field device set up

The PEMF exposure device was previously stated in study by Hannay et al. (17). Briefly, two separate coils with dimensions 150×100 mm which was made up of 50 turns of 0.51 mm diameter acrylic coated copper wire were connected together in series and placed 20 mm apart. Each coil produced a resistance of 2.3 Ω . For inducing a parallel-aligned electric field, a PEMF pulse generator that produced a pulsed magnetic field per-

pendicular to the cell monolayer were wired to the coils. The PEMF signal contain 20 pulses of 5 milliseconds burst that was repeating at 15 Hertz. It was creating an asymmetrical "quasisquare wave" voltage trace during each burst at a frequency of ~4 kilohertz. Peak coil current duration lasted for 204 milliseconds, producing a maximum magnetic flux of 1.3 millitesla. The electromagnetic field strength induced inside the plate was identical thoroughly and independent to distance from the center (Fig 2). The Effect of PEME on IVE Success Rate



Fig 2: ELF- PEMF device settings. A. Wave generator, B. Amplifier 30 W, C. The designed circuit



Fig3: Frequencies of metaphase II ovums.

100% Unfertiliized ovum 90% 19 12 80% Fertilized ovum 70% 60% 50% 39 40 40% 30% 20% 10% 0% 58 (30.05%) 52 ((30.36%) Control aroup Experimental group

Fig 4: Comparison of percentage of fertilized and unfertilized ovums between two groups.

Discussion

Due to undeniable negative social and economic consequences of infertility (32) and also failure in ART or other infertility therapies, it seems more attention should be devoted to improve the efficacy of methods.

The aim of this study was clarifying ELF-PEMF exposure effect on mice IVF success rate. The primary results showed an increase in forming couple pronuclei in ELF-PEMF exposed group, but that was a statistically insignificant increase. Consequently, we can say that exposing gametes to 50 Hertz pulsating electromagnetic field can influence the fertilization process.

Various hypotheses have been proposed for the mechanism of EMF action on cells. The most probable target for EMFs is the plasma membrane

Statistical analysis

To compare the obtained results between experimental and control groups, Statistical Package for the Social Sciences (SPSS) software 16th release was carried out, whilechi-Square and Fisher's exact test were applied to interpret the values. The p value under 0.05 was considered significant.

Results

Total number of collected ovums in the control and experimental groups was 191 and 173, respectively, from which, 58 (30.05%) and 52 (30.36%) ovums were collected in metaphase II, respectively. Forming two pronuclei, which was considered as IVF success rate, was 77% in ELF-PEMF exposed group (experimental), while 68% in control group (Table1, Fig 3, 4).

Table1: Comparison of obtained results between two groups		
Variable	Control group	Experimental group
Total number of ovums	191	173
Ovums in metaphase II	58 (30.05 %)	52 (30.36 %)
Fertilized ovums	39	40
Unfertilized ovums	19	12
IVF success rate	68%	77%

and transmembrane proteins rather than the cytoplasm. Gap junctions, specialized intercellular junctions, have been proposed as mediators of the EMF related cellular responses which change the cellular activity. One of the organelles responding to electromagnetic fields is microtubule, but this has not been proven yet (33, 34).

Despite numerous studies about the effects of ELF-EMF on reproduction system, a paradigm has remained. Investigation of ELF-EMF on mammalian sperm showed both negative (3, 4, 35) and positive (14, 16, 36) effects. This controversy is also coherent with female fertility issues in which exposure to EMF was accompanied (37, 38) or not accompanied (39, 40) by significant adverse effects such as congenital anomaly. Ryan et al. (39) and Ohnishi et al. (40) have demonstrated that exposure to pure, linearly polarized 60 Hartz and power-frequency magnetic fields has no major effects on reproduction and development in mice. Tomás et al. (41) have also showed an increase in reproductive investment by breeding adults exposed to EMFs as compared to those in the adjacent reference area.

These findings have opened a new perspective and a growing interest for us to clarify whether ELF EMF ameliorate the IVF success rate in mice. It seems that no previous studies have been done about electromagnetic field as a promoting factor in mice IVF.

According to our previous unpublished data, we observed increased proliferation and differentiation of leukemic lymphoblasts after being exposed to ELF EMF emanating from a system same as in current study.

It is difficult to compare different studies due to many factors such as variety of frequency, intensity, timing and other magnetic properties which may interfere in results. But the structure of magnetic producer and circuit seems to be the most important factor.

Conclusion

In this pilot animal study, we observed insignificantly improvement of IVF rate in NMRI mice. On the basis of vital effects of natural electromagnetic fields on cells and organisms, although our results were not significantly positive, we suggest investigating the effects of extremely low frequency PEMF (ELF-PEMF) on success rate of IVF on larger sample sizes of NMRI mice. We also strongly recommend more focusing on probable teratogens and other genetic disorders that may occur during ELF-PEMF exposure. One of our limitations in this study was small sample size.

Acknowledgements

This project was financially supported by Mashhad University of Medical Sciences, Vice Chancellor of Research. There is no conflict of interest to disclose.

References

- Dini L, Abbro L. Bioeffects of moderate-intensity static magnetic fields on cell cultures. Micron. 2005; 36(3): 195-217.
- Soeradi O, Tadjudin MK. Congenital anomalies in the offspring of rats after exposure of the testis to an electrostatic field. Int J Androl. 1986; 9(2): 152-160.
- Bernabo N, Tettamanti E, Pistilli MG, Nardinocchi D, Berardinelli P, Mattioli M, et al. Effects of 50 Hz extremely low frequency magnetic field on the morphology and function of boar spermatozoa capacitated in vitro. Theriogenology. 2007; 67(4): 801-815.
- Bernabo N, Tettamanti E, Russo V, Martelli A, Turriani M, Mattoli M, et al. Extremely low frequency electromagnetic field exposure affects fertilization outcome in swine animal model. Theriogenology. 2010; 73(9): 1293-1305.
- Al-Akhras MA, Elbetieha A, Hasan MK, Al-Omari I, Darmani H, Albiss B. Effects of extremely low frequency magnetic field on fertility of adult male and female rats. Bioelectromagnetics. 2001; 22(5): 340-344.
- Hisamitsu T, Narita K, Kasahara T, Seto A, Yu Y, Asano K. Induction of apoptosis in human leukemic cells by magnetic fields. Jpn J Physiol. 1997; 47(3): 307-310.
- Tenuzzo B, Chionna A, Panzarini E, Lanubile R, Tarantino P, Di Jeso B, et al. Biological effects of 6 mT static magnetic fields: a comparative study in different cell types. Bioelectromagnetics. 2006; 27(7): 560-577.
- Walleczek J, Liburdy RP. Nonthermal 60 Hz sinusoidal magnetic-field exposure enhances 45Ca2+ uptake in rat thymocytes: dependence on mitogen activation. FEBS lett. 1990; 271(1-2): 157-160.
- Scaiano JC, CozensFL, Mohtat N. Influence of combined AC-DC magnetic fields on free radicals in organized and biological systems. Development of a model and application of the radical pair mechanism to radicals in micelles. Photochem Photobiol. 1995; 62(5): 818-829.
- Simko M, Droste S, Kriehuber R, Weiss DG. Stimulation of phagocytosis and free radical production in murine macrophages by 50 Hz electromagnetic fields. Eur J Cell Biol. 2001; 80(8): 562-566.
- 11. Rollwitz J, Lupke M, Simko M. Fifty-hertz magnetic fields induce free radical formation in mouse bone marrow-derived promonocytes and macrophages. Biochim Biophys Acta. 2004; 1674(3): 231-238.
- Ivancsits S, Diem E, Pilger A, Rudiger HW, Jahn O. Induction of DNA strand breaks by intermittent exposure to extremely-low-frequency electromagnetic fields in human diploid fibroblasts. Mutat Res. 2002; 519(1-2): 1-13.
- 13. Wolf FI, Torsello A, Tedesco B, Fasanella S, Bonin-

CELL JOURNAL(Yakhteh), Vol 15, No 4, Winter 2014 314

segna A, D'Ascenzo M, et al. 50-Hz extremely low frequency electromagnetic fields enhance cell proliferation andDNA damage: possible involvement of a redox mechanism. Biochim Biophys Acta. 2005; 1743(1-2): 120-129.

- Iorio R, Scrimaglio R, Rantucci E, Delle Monache S, Di Gaetano A, Finetti N, et al. A preliminary study of oscillating electromagnetic field effects on human spermatozoon motility. Bioelectromagnetics. 2007; 28(1): 72-75.
- Delle Monache S, Alessandro R, Iorio R, Gualtieri G, Colonna R. Extremely low frequency electromagnetic fields (ELF-EMFs) induce in vitro angiogenesis process in human endothelial cells. Bioelectromagnetics. 2008; 29(8): 640-648.
- Iorio R, Delle Monache S, Bennato F, Di Bartolomeo C, Scrimaglio R, Cinque B, et al. Involvement of mitochondrial activity in mediating ELF-EMF stimulatory effect on human sperm motility. Bioelectromagnetics. 2011; 32(1): 15-27.
- Hannay G, Leavesley D, Pearcy M. Timing of pulsed electromagnetic field stimulation does not affect the promotion of bone cell development. Bioelectromagnetics. 2005; 26(8): 670-676.
- De Mattei M, Caruso A, Traina GC, Pezzetti F, Baroni T, Sollazzo V. Correlation between pulsed electromagnetic fields exposure time and cell proliferation increase in human osteosarcoma cell lines and human normal osteoblast cells in vitro. Bioelectromagnetics. 1999; 20(3): 177-182.
- Scott G, King JB. A prospective, double-blind trial of electrical capacitive coupling in the treatment of non-union of long bones. J Bone Joint Surg Am. 1994; 76(6): 820-826.
- Baker LL, Chambers R, DeMuthSK, Villar F. Effects of electrical stimulation on wound healing in patients with diabetic ulcers. Diabetes care. 1997; 20(3): 405-412.
- Trock DH, Bollet AJ, Dyer RH Jr, Fielding LP, Miner WK, Markoll R. A double-blind trial of the clinical effects of pulsed electromagnetic fields in osteoarthritis. J Rheumatol. 1993; 20(3): 456-460.
- Trock DH, Bollet AJ, Markoll R. The effect of pulsed electromagnetic fields in the treatment of osteoarthritis of the knee and cervical spine. Report of randomized, double blind, placebo controlled trials. J Rheumatol. 1994; 21(10): 1903-1911.
- Stiller MJ, Pak GH, Shupack JL, Thaler S, Kenny C, Jondreau L. A portable pulsed electromagnetic field (PEMF) device to enhance healing of recalcitrant venous ulcers: a double-blind, placebo-controlled clinical trial. Br J Dermatol. 1992; 127(2): 147-154.
- Lee EW, Maffulli N, Li CK, Chan KM. Pulsed magnetic and electromagnetic fields in experimental achilles tendonitis in the rat: a prospective randomized study. Arch Phys Med Rehabil. 1997; 78(4); 399-404.
- Kim SS, Shin HJ, Éom DW, Huh JR, Woo Y, Kim H, et al. Enhanced expression of neuronal nitric oxide synthase and phospholipase C-gamma1 in regenerating murine neuronal cells by pulsed electromagnetic field. Exp Mol Med. 2002; 34(1): 53-59.
- Gmitrov J, Ohkubo C, Okano H. Effect of 0.25 T static magnetic field on microcirculation in rabbits. Bioelectromagnetics. 2002; 23(3): 224-229.
- Okano H, Onmori R, Tomita N, Ikada Y. Effects of a moderate-intensity static magnetic field on VEGF-A stimulated endothelial capillary tubule formation in vitro. Bioelectro-

magnetics. 2006; 27(8): 628-640.

- Zare S, Hayatgeibi H, Alivandi S, Ebadi AG. Effects of whole-body magnetic field on changes of glucose and cortisol hormone in guinea pigs. Am J Biochemistry Biotechnol. 2005; 1(4): 217-219.
- Zegers-Hochschild F, Adamson GD, de Mouzon J, Ishihara O, Mansour R, Nygren K, et al. International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) revised glossary of ART terminology. Fertil Steril. 2009; 92(5): 1520-1524.
- Centers for Disease Control and Prevention, American Society for Reproductive Medicine, Society for Assisted Reproductive Technology. 2007 Assisted Reproductive Technology Success Rates: National Summary and Fertility Clinic Reports, Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2009.
- Nielsen HI. History and fundamentals of oocyte maturation in vitro. Cell J. 2007; 9 Supple 1.
- Ramezanzadeh F, Aghssa MM, Abedinia N, Zayeri F, Khanafshar N, Shariat M, et al. A survey of relationship between anxiety, depression and duration of infertility. BMC Womens Health. 2004; 4(1): 9.
- Piacentini R, Ripoli C, Mezzogori D, Azzena GB, Grassi C. Extremely low-frequency electromagnetic fields promote in vitro neurogenesis via upregulation of Ca(v)1-channel activity. J Cell Physiol. 2008; 215(1): 129-139.
- 34. Goodman R, Lin-Ye A, Geddis MS, Wickramaratne PJ, Hodge SE, Pantazatos SP, et al. Extremely low frequency electromagnetic fields activate the ERK cascade, increase hsp70 protein levels and promote regeneration in Planaria. Int J Radiat Biol. 2009; 85(10): 851-859.
- De Vita R, Cavallo D, Raganella L, Éleuteri P, Grollino MG, Calugi A. Effects of 50 Hz magnetic fields on mouse spermatogenesis monitored by flow cytometric analysis. Bioelectromagnetics. 1995; 16(5): 330-334.
- Roychoudhury S, Jedlicka J, Parkanyi V, Rafay J, Ondruska L, Massanyi P, et al. Influence of a 50 hz extra low frequency electromagnetic field on spermatozoa motility and fertilization rates in rabbits. J Environ Sci Health A Tox Hazard Subst Environ Eng. 2009; 44(10): 1041-1047.
- Huuskonen H, Juutilainen J, Komulainen H. Éffects of lowfrequency magnetic fields on fetal development in rats. Bioelectromagnetics. 1993; 14(3): 205-213.
- Mevissen M, Buntenkotter S, Loscher W. Effects of static and time-varying (50-Hz) magnetic fields on reproduction and fetal development in rats. Teratology. 1994; 50(3): 229-237.
- Ryan BM, Symanski RR, Pomeranz LE, Johnson TR, Gauger JR, McCormick DL. Multigeneration reproductive toxicity assessment of 60-Hz magnetic fields using a continuous breeding protocol in rats. Teratology. 1999; 59(3): 156-162.
- Ohnishi Y, MizunoF, Sato T, Yasui M, Kikuchi T, Ogawa M. Effects of power frequency alternating magnetic fields on reproduction and pre-natal development of mice. J Toxicol Sci. 2002; 27(3): 131-138.
- Tomás G, Barba E, Merino S, Martínez J. Clutch size and egg volume in great tits (Parus major) increase under low intensity electromagnetic fields: a long-term field study. Environ Res. 2012; 118: 40-46.