

The Effects of Levofloxacin on Testis Tissue and Spermatogenesis in Rat

Ramesh Ahmadi, Ph.D.¹, Mehdi Ahmadifar, M.Sc.^{2,3}, Elham Safarpour, Ph.D.⁴,
Nazila Vahidi-eyrisofla, M.Sc.⁵, Mehraneh Darab, Ph.D.⁶,
Ali Mohammad Eini, M.Sc.⁶, AliReza Alizadeh, Ph.D.^{7*}

1. Department of Biology, Qom Branch, Islamic Azad University, Qom, Iran
2. Department of Biology, College of Science, University of Science and Culture, Tehran, Iran
3. Department of Embryology, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran
4. The Toronto Institute for Reproductive Medicine, (Repromed), Toronto, Canada
5. Islamic Azad University, Damghan Branch, Damghan, Iran
6. Young Researchers and Elites Club, Science and Research Branch, Islamic Azad University, Tehran, Iran
7. Department of Animal Science, Saveh Branch, Islamic Azad University, Saveh, Iran

*Corresponding Address: P.O. Box: 39187/366, Department of Animal Science, Saveh Branch, Islamic Azad University, Saveh, Iran
Email: dr.alizadeh@iau-saveh.ac.ir

Received: 17/Mar/2015, Accepted: 30/Jul/2015

Abstract

Levofloxacin is one of the Fluroquinoline antibiotic groups, which affect on controlling infections, especially in reproductive organs. It has therapeutic use in numerous countries, but little information exists on the effects of Levofloxacin on spermatogenesis when it is used for infectious treatment. The current study was designed to determine whether Levofloxacin influences testis tissue and spermatogenesis in rats.

In this survey 50 male Wistar rats 6-8 weeks (250 ± 10 g) were used: normal saline as sham and control groups and 3 treatment groups (0.03, 0.06 and 0.08 mg Levofloxacin/kg body weight) during 60 days. The experimental groups were daily gavages. After 60 days, they were anesthetized with ether and testes were taken for histopathology studies, sperm parameters evaluation and several hormone concentrations.

Although testosterone concentration was not affected by Levofloxacin levels, follicle stimulating hormone (FSH) and luteinizing hormone (LH) concentration significantly increased by Levofloxacin consumption in 0.03 and 0.06 mg Levofloxacin/kg body weight groups ($P < 0.01$). Moreover, sperm concentration decreased linearly as Levofloxacin was increased (200, 192, 170, 128 and 75×10^6 sperm for control, sham, 0.03, 0.06 and 0.08 mg Levofloxacin/kg body weight, respectively, $P < 0.05$). Testis tissue cuts in experimental group when the amount dosage of Levofloxacin increased cells solidarity to the primary and secondary spermatogonia. Adding Levofloxacin linearly reduced spermatocyte cells and amount of all cells in semenifer pipes tube ($P < 0.05$).

Levofloxacin as an antibiotic has histopathology effects on the spermatocyte cells, especially in high dose. Therefore, it might reduce fertility in male that requires further studies.

Keywords: Levofloxacin, Spermatogenesis, Rat

Cell Journal (Yakhteh), Vol 18, No 1, Apr-Jun (Spring) 2016, Pages: 112-116

Citation: Ahmadi R, Ahmadifar M, Safarpour E, Vahidi-eyrisofla N, Darab M, Eini AM, Alizadeh AR. The effects of levofloxacin on testis tissue and spermatogenesis in rat. Cell J. 2016; 18(1): 112-116.

One of the main problems in the world of today's medicine is infertility effects. In other words, approximately 50% of infertility is related to problems in male. Multiple environmental factors are life style (nutrition, drugs, smoking and alcohol), diseases (systemic, genetic, and infectious) and

antibiotics. Antibiotics are one of the main factors which involved in male infertility (1, 2). The research indicates that antibiotics have effects on semen as well as known side effects on sperm mechanism during treatment period (3, 4). Composition of Cotrimoxazole induces the reduction of number

and movement of sperm (5). Sulfasalazine cause infertility (6) as well as sperm count reduction (7), oligospermia (8) and sperm immotility (9) and abnormal morphology (10).

Levofloxacin is one of the Fluroquinoline antibiotic groups, which have effects on controlling infections, especially in reproductive organs. It has therapeutic use in more than 100 countries, but little information exists on the impacts of Levofloxacin on spermatogenesis when it is used for infection treatment. Sometimes patients need to take antibiotics for long period for about 50-70 days, whereas spermatogenesis duration is 64 ± 8 and 48 ± 5 days in human and rat, respectively (11, 12). Researchers have shown that antibiotics, such as Amoxicillin, Erythromycin and Cotrimoxazol reduce the sperm concentration, too (13). Levofloxacin is so effective in treatment of chronic infectious diseases. In many sexually transmitted diseases infections of the urogenital tract and diseases such as tuberculosis and brucellosis need long term antibiotics in take for treatment that sometimes takes 10 to 50 days of treatment. However, to our knowledge, the question of whether using Levofloxacin affects spermatogenesis in male has not been addressed. The current study was designed to determine whether Levofloxacin influences testis tissue in rats.

This study received the approval of the Ethics Committee of Royan Institute. In this study, 50 male Wistar rats 6-8 weeks (250 ± 10 g) were used. Duration of the survey lasted 60 days. The rats were exposed to lightness and then darkness for 12 hours, respectively. The room temperature stood at $23-25^{\circ}\text{C}$ as its moisture was counted as 50-55%. The experimental groups are normal saline as sham and control group and 3 treatment groups (0.03, 0.06 and 0.08 mg Levofloxacin/kg body weight). That dose is in human but we mixed drug with distilled water with due consideration to the rat weight (2). After 60 days, rats were anesthetized. Blood was centrifuged (Spectrafuge, 16MLabnet IntUSA) at $14,000 \times g$ for 15 minutes at 10°C to obtain serum. Follicle stimulating hormone (FSH), luteinizing hormone (LH) and testosterone were measured by using laboratory procedures and commercial kits. Hematoxyline-eosin (H&E) several slides were studied by optical micro-

scope equipped with camera. In survey of spermatogenic cells (spermatogonia, primary spermatocyte, spermatid, sperm, and sertoli cells) compared the amount of each one of cells in experimental groups and control ones. The data generated from the study were analyzed by ANOVA by using the General Linear Model (GLM) procedure of SPSS. Differences between treatment means were tested by Duncan's Multiple Range test.

In current study gavages of several doses 0.03 and 0.06 mg/kg Levofloxacin did not affect cells and full sperms. But destruction in tubules as well as spermatogonia and primary spermatocytes were altered by 0.08 mg/ml Levofloxacin (Fig.1A-E, $P < 0.05$). Also sperms are not fully formed because of high dose of drug and there is accumulation of interstitial fluid in tubules. Although testosterone concentration was not affected by Levofloxacin, FSH and LH concentrations significantly increased by Levofloxacin consumption (Fig.2A-C). Sperm concentration decreased linearly as Levofloxacin was consumed ($200, 192, 170, 128$ and 75×10^6 sperm for control, sham, 0.03, 0.06 and 0.08 mg Levofloxacin/kg body weight, respectively).

The previous research reveals that Ciprofloxacin causes the expected death of cells by activating caspase3 which can play an important role in sperm reduction (14-18), sperm motility, and the increase of sliced DNA in sperm, and varicocele (19). Another former research done previously shows the substantial effect of Moxifloxacin on the number and motility of sperm, and the process spermatogenesis. This effect was so that it could reduce the number and motility of sperms and normal morphology (20-23).

One of these factors is to stay healthy in germinal cells create a balance between the number of live cells and dead ones both of which play a role in spermatogenesis and the rate of sperm production. The amount of germinal sexual cells has reduced because of different types of dead cells such as apoptosis (24). In recent years, some of conducted studies on animal laboratories have show the role of chemicals, radiation, infection, virus, cod, freeze and removal of pituitary gland from body when apoptosis occurred and virus cell death in testis tissue (12, 25, 26).




Fig.1: A. Microscopic image of the testis tissue of the control group, seminiferous tubules (St), and less Leydig cells (Ls) [Hematoxyline-eosin (H&E) stain $\times 200$], B. Microscopic image of the testis tissue of the sham group, ST, Ls (H&E stain $\times 200$), C. Microscopic image of the testis tissue of the 0.03 mg Levofloxacin/kg body weight, immature spermatozoa (IM), (H&E stain $\times 200$), D. Microscopic image of the testis tissue of the 0.06 mg Levofloxacin/kg body weight, IM, less seminiferous tubules (LF), less Ls, (H&E stain $\times 200$) and E. Microscopic image of the testis tissue of the 0.08 mg Levofloxacin/kg body weight, IM, LF, less Ls, (H&E stain $\times 200$).




Fig.2: A. Testosterone concentration in experimental groups, B. LH concentration in experimental groups and C. FSH concentration in experimental groups. ***; $P < 0.001$, **; $P < 0.01$, T1; Group treatment by 0.03 mg/kg dosage, T2; Group treatment by 0.06 mg/kg dosage, T3; Group treatment by 0.08 mg/kg dosage, FSH; Follicle serum human and LH; Lutheal hormone.

It seems that several antibiotics might affect reproductive hormones by several pathways. Although the negative significant repercussion of Ciprofloxacin on FSH and LH concentration were reported in previous studies (14, 27), time-dependent and dose-dependent responses may explain the Levofloxacin effects on hormones. Indeed, schedule sampling in all studies with antibiotics was conducted at the completion of the experiment and

the time of the maximum response has remained unclear in studies. Therefore, increased FSH and LH concentration in some experimental groups of these studies need more investigations.

Evaluation made on fluoroquinolones reveals these drugs have ripple effect on testis tissue and sperm parameters as well. It appears in artificial insemination (AI) process using Ciprofloxacin has priority to Gentamicin. The reason is that antibiotic in high dose is capable of destroying bacterial contamination of semen with less undesirable impacts on the functions of sperm. Further surveys are required so as to make an efficient comparison between fluoroquinolones and aminoglycosides. For evaluation of fluoroquinolones and aminoglycosides analogues parameters are applied. Moreover, it is proposed that further research on all drugs as part of aminoglycosides and fluoroquinolones be carried out on human beings. In a separate survey conducted on Gentamicin's toxicity on rat sperm, reduced caudal epididymal sperm reserves administration of Gentamicin. Ensuing Gentamicin administration sperm mobility decreased (28).

Sex hormones (LH and FSH) and spermatogenesis (sperm count, motility and viability) were significantly decreased in test group compared to those of controls ($P < 0.05$). Ciprofloxacin has some adverse effects on sperm related variables in 28 day period. It appears that dose-dependent response is important in this issue; because the testes cells in high dose groups try to compensate the destroyed cells by increasing FSH and LH concentration.

Altogether, the administration of therapeutic doses of Levofloxacin 0.08 mg/ml for 60 days can cause pathological changes such as atrophy. Seminiferous tubule and irreversible damages to the cells in the testis cause death to the spermatogonia cells and primary spermatocytes disrupt the normal cycle of spermatogenesis and caused hypo spermatogenesis and infertility in male rats. Thus, our results in high dose treatment support the theory that the use of some antibiotics such as Levofloxacin disturbs spermatogenesis, particularly Leydig cells, primary and secondary spermatocytes as well as Sertoli cells. Our results suggest that unlike other antibiotics this drug have dose-dependent effects on all parameters of spermatogenesis and have more histopathology effects.

Acknowledgments

This study was financed by a grant from Islamic Azad University, Qom Branch, Qom, Iran. None of the authors have any conflict of interest to declare.

References

1. Abd-Allah AR, Aly HA, Moustafa AM, Abdel-Aziz AA, Hamada FM. Adverse testicular effects of some quinolone members in rats. *Pharmacol Res.* 2000; 41(2): 211-219.
2. Huang K, Yang J, Zhang J, Ding Y, Chen L, Xu WY, et al. Determination of sitafloxacin in human plasma by liquid chromatography-tandem mass spectrometry method: application to a pharmacokinetic study. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2014; 957: 36-40.
3. Chapman JS, Georgopadakou NH. Routes of quinolone permeation in *Escherichia coli*. *Antimicrob Agents Chemother.* 1988; 32(4): 438-442.
4. Summers MC. A brief history of the development of the KSOM family of media. *J Assist Reprod Genet.* 2013; 30(8): 995-999.
5. Douros A, Grabowski K, Stahlmann R. Safety issues and drug-drug interactions with commonly used quinolones. *Expert Opin Drug Metab Toxicol.* 2015; 11(1): 25-39.
6. Womble PR, Linsell SM, Gao Y, Ye Z, Montie JE, Gandhi TN, et al. A statewide intervention to reduce hospitalizations after prostate biopsy. *J Urol.* 2015; 194(2): 403-409.
7. Velkov T, Bergen PJ, Lora-Tamayo J, Landersdorfer CB, Li J. PK/PD models in antibacterial development. *Curr Opin Microbiol.* 2013; 16(5): 573-579.
8. Takahara K, Ibuki N, Ghaffari M, Tearle H, Ong CJ, Azuma H, et al. The influence of growth hormone/insulin-like growth factor deficiency on prostatic dysplasia in *pARR2-Cre*, *PTEN* knockout mice. *Prostate Cancer Prostatic Dis.* 2013; 16(3): 239-247.
9. Bjarnsholt T, Ciofu O, Molin S, Givskov M, Høiby N. Applying insights from biofilm biology to drug development - can a new approach be developed? *Nat Rev Drug Discov.* 2013; 12(10): 791-808.
10. Norra C, Skobe E, Breuer C, Haase G, Hanrath P, Hoff P. Ciprofloxacin-induced acute psychosis in a patient with multidrug-resistant tuberculosis. *Eur Psychiatry.* 2003; 18(5): 262-263.
11. Villani P, Fresegna AM, Ranaldi R, Eleuteri P, Paris L, Pacchierotti F, et al. X-Ray induced DNA damage and repair in germ cells of *PARP1(-/-)* male mice. *Int J Mol Sci.* 2013; 14(9): 18078-18092.
12. O'shaughnessy PJ. Hormonal control of germ cell development and spermatogenesis. *Semin Cell Dev Biol.* 2014; 29: 55-65.
13. Abbitt B, Berndtson WE, Seidel GE Jr. Effect of dihydrostreptomycin or oxytetracycline on reproductive capacity of bulls. *Am J Vet Res.* 1984; 45(11): 2243-2246.
14. Elias A, Nelson B. Sub chronic evaluation of ciprofloxacin and perfloxacin on sperm parameters of male guinea pigs. *Asian J Exp Biol Sci.* 2012; 3: 595-601.
15. Vasquez MI, Garcia-Käufer M, Hapeshi E, Menz J, Kostarelos K, Fatta-Kassinou D, et al. Chronic ecotoxic effects to *Pseudomonas putida* and *Vibrio fischeri* and cytostatic and genotoxic effects to the hepatoma cell line (HepG2) of ofloxacin photo (cata) lytically treated solutions. *Sci Total Environ.* 2013; 450-451: 356-365.
16. Khaki A, Heidari M, Ghaffari Novin M, Khaki AA. Adverse effects of ciprofloxacin on testis apoptosis and sperm parameters in rats. *Iran J Reprod Med.* 2008; 6(2): 71-76.
17. Raz R, Naber KG, Raizenberg C, Rohana Y, Unamba-Oparah I, Korfman G, et al. Ciprofloxacin 250 mg twice daily versus ofloxacin 200 mg twice daily in the treatment of complicated urinary tract infections in women. *Eur J Clin Microbiol Infect Dis.* 2000; 19(5): 327-331.
18. Wang M, Tran JH, Jacoby GA, Zhang Y, Wang F, Hooper DC. Plasmid-mediated quinolone resistance in clinical isolates of *Escherichia coli* from Shanghai China. *Antimicrob Agents Chemother.* 2003; 47(7): 2242-2248.
19. Weyers AI, Ugnia LI, Garcia Ovando H, Gorla NB. Ciprofloxacin increases hepatic and renal lipid hydroperoxides levels in mice. *Biocell.* 2002; 26(5): 225-228.
20. Rodríguez-Martínez JM, Cano ME, Velasco C, Martínez-Martínez L, Pascual A. Plasmid-mediated quinolone resistance: an update. *J Infect Chemother.* 2011; 17(2): 149-182.
21. Ong-Tone L. Aqueous humor penetration of gatifloxacin and moxifloxacin eyedrops given by different methods before cataract surgery. *J Cataract Refract Surg.* 2007; 33(1): 59-62.
22. Lane SS, Osher RH, Masket S, Belani S. Evaluation of the safety of prophylactic intracameral moxifloxacin in cataract surgery. *J Cataract Refract Surg.* 2008; 34(9): 1451-1459.
23. O'Brien TP, Arshinoff SA, Mah FS. Perspectives on antibiotics for postoperative endophthalmitis prophylaxis: potential role of moxifloxacin. *J Cataract Refract Surg.* 2007; 33(10): 1790-1800.
24. Norra C, Skobel E, Breuer C, Haase G, Hanrath P, Hoff P. Ciprofloxacin-induced acute psychosis in a patient with multidrug-resistant tuberculosis. *Eur Psychiatry.* 2003; 18(5): 262-263.
25. Dutta HM, Misquitta D, Khan S. The effects of endosulfan on the testes of bluegill fish, *Lepomis macrochirus*: a histopathological study. *Arch Environ Contam Toxicol.* 2006; 51(1): 149-156.
26. Rajaei F, Karja NW, Agung B, Wongsrikeao P, Taniguchi M, Murakami M, et al. Analysis of DNA fragmentation of porcine embryos exposed to cryoprotectants. *Reprod Domest Anim.* 2005; 40(5): 429-432.
27. Blumberg HM, Rimland D, Carroll DJ, Terry P, Wachsmuth IK. Rapid development of ciprofloxacin resistance in methicillin-susceptible and -resistant *Staphylococcus aureus*. *J Infect Dis.* 2001; 163(6): 1279-1285.
28. Zahedi A, Khaki A, Ahmadi-Ashtiani HR, Rastegar H, Reza-zadeh Sh. Zingiber officinale protective effects on gentamicin's toxicity on sperm in rats. *J Med Plants.* 2010; 9(35): 93-98.
29. Khaki A. Effect of Ciprofloxacin on Sexual Hormones and Spermatoocytes in Rat. *MLJ.* 2014; 8 (2): 8-13.