Original Article

Effect of Fetal Mouse Lung Tissue Co-Culture on *In Vitro* Maturation of Mouse Immature Oocytes

Masomeh Belbasi, M.Sc.¹, Seyed Gholam Ali Jorsaraei, Ph.D.^{2*}, Maryam Gholamitabar tabari, M.Sc.^{2*}, Ramzan Khanbabaei, Ph.D.¹

1. Department of Biology, Qaemshahr Branch, Islamic Azad University, Qaemshahr, Iran 2. Infertility and Reproductive Health Research Center, Health Research Institute, Babol University of Medical Science, Babol, Iran

*Corresponding Address: P.O.Box: 4136747176, Infertility and Reproductive Health Research Center, Health Research Institute, Babol University of Medical Science, Babol, Iran Emails: alijorsara@yahoo.com, Maryam.tabari407@gmail.com

Received: 2/May/2015, Accepted: 3/Sep/2016

Abstract —

Objective: The aim of this study was to evaluate the fetal mouse lung tissue co-culture on *in vitro* maturation (IVM) of mouse immature oocytes.

Materials and Methods: In this experimental study, germinal vesicle (GV) oocytes from ovaries of a group of 25 female mice, 6-8 weeks of age, were dissected after being stimulated by 7.5 IU pregnant mare serum gonadotropin (PMSG) through an intraperitoneal (IP) injection. The fetal lung tissues were then prepared and cultured individually. A total number of 300 oocytes were cultured in the following three groups for 24 hours: control group (n=100) containing only base medium, group I (n=100) containing base medium co-cultured with 11.5- to 12.5-day old fetal mouse lung tissues, and group II (n=100) containing base medium co-cultured with 12.5- to 13.5-day old fetal mouse lung tissues. The proportion of GV and metaphase I (MI) oocytes matured into MII oocytes were compared among the three groups using analysis of variance (ANOVA). Correlation test were also used to evaluate the successful rate of IVM oocytes.

Results: The proportions of GV oocytes reaching MII stage were 46, 65, and 56%, in control, I and II groups, respectively (P<0.05). The percentage of the oocytes remaining at the GV stage were higher in control group as compared with two treatment groups (P<0.05).

Conclusion: This study indicated that fetal mouse lung tissue co-culture method increased the percentage of GV oocytes reaching MII stage.

Keywords: In Vitro Maturation, Lung Tissue, Co-Culture

Cell Journal(Yakhteh), Vol 19, No 3, Oct-Dec (Autumn) 2017, Pages: 476-481 _

Citation: Belbasi M, Jorsaraei SGA, Gholamitabar tabari M, Khanbabaei R. Effect of fetal mouse lung tissue co-culture on in vitro maturation of mouse immature oocytes. Cell J. 2017; 19(3): 476-481. doi: 10.22074/cellj.2017.3866.

Introduction

Following the first successful clinical application of *in vitro* fertilization (IVF), Cha et al. (1) reported the first live birth from *in vitro* maturation (IVM) a decade later. In this advanced experimental technique, immature oocytes are extracted from the ovaries without need of gonadotropin stimulation. Various factors such as nutrients, energy substrates, vitamins, hormones and growth factors help *in vitro* development of oocytes (2-4). Fibroblast growth factor (FGF) family play an important role in setting the initial follicle growth, oocyte survival, proliferation and differentiation of granulosa cells, corpus luteum (CL) formation, steroidogenesis and angiogenesis (5-7). Many paracrine-acting factors have ability to arrange oocyte maturation and follicular development. More than 22 genes encode FGFs as significant paracrine regulators of angiogenesis, morphogenesis luteinizing hormone and reproduction (8, 9). FGFs, especially FGF2 produced from granulosa cell, are involved in luteinizing hormone (LH) receptor expression, primordial follicle development and granulosa cell proliferation (10-13). FGF10 is a significant mediator of mesenchymal-epithelial signaling in various organs and tissues (14, 15). In addition to the several members of the FGFs family, FGF receptors (FGFRs) have been implicated in mouse lung development (16). FGF10 is expressed at high levels from the earliest stages of lung development (15). Use of fibroblast co-culture and activin A promoted the growth and maintenance of preantral follicles in mice (17). Also, in a study by Heidari et al. (18) on evaluation of the effect of fibroblast coculture system on the maturation and fertilization of oocytes from mouse preantral follicles, they have suggested that fibroblast co-culture increased the growth and survival rate of cultured preantral follicles in a significant manner by enhancement of granulosa cell proliferation. Retinoic acid is another factor playing a role on lung regeneration (19) and considered as an important modifier of vertebrate development, cell differentiation and tissue function (20). IVM with 9-cis-retinoic acid increased the developmental competence of the oocyte (21). Co-culturing is a method to support a long culturing period of IVM. In vitro co-culturing system has advantages in secreting trophic factors such as nutrients, substrates, growth factors, and cytokines as well as in eliminating toxins from the culture medium (22). We hypothesized that co-culturing with embryonic mouse lung tissue cells may promote immature follicle development due to presence of FGF and retinoic acid in early embryonic life of mouse life. The effect of embryonic lung tissue cells co-culture on IVM of immature follicles has not yet been studied in any species. Therefore, to improve culture conditions and develop appropriate culture systems with the idea of stimulating immature follicle growth, the present study aimed to investigate the effect of mouse embryonic lung tissue cells on in vitro growth of mouse immature follicles.

Materials and Methods

This is an experimental study that was approved by the Islamic Azad University, Science and Research Branch (Qaemshahr, Iran). All reagents were obtained from Gibco, UK, unless otherwise specified.

Male and female NMRI mice were obtained from Pasteur's Institute (Amol, Iran) and kept in the Central Animal House of Babol University of Medical Science (Babol, Iran) under a 12hour light/12- hour dark regime at 20-24°C, with adequate food ad libitum.

Culture medium condition

Base medium was consisted of alpha-Minimal Essential Medium (α -MEM, Gibco, UK), 100 µg/ml penicillin, 50 µg/ml streptomycin, and 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) supplemented with 10% fetal bovine serum (FBS, Gibco, UK). A total number of 300 oocytes were cultured in the following three groups for 24 hours: control group (n=100) containing only base medium, group I (n=100) containing base medium co-cultured with 11.5- to 12.5-day old fetal mouse lung tissues and group II (n=100) containing base medium co-cultured with 12.5- to 13.5-day old fetal mouse lung tissues.

Germinal vesicle follicle isolation

Mice were stimulated by 7.5 IU pregnant mare serum gonadotropin (PMSG, Hipra, Spain) through an intraperitoneal (IP) injection. The animals were killed by cervical dislocation 48 hours later, while the ovaries were removed and placed on dissection medium consisting of α -MEM supplemented with 10% FBS, 100 µg/ml penicillin and 50 µg/ml streptomycin under mineral oil (Irvine Scientific, Belgium) to prevent evaporation, severe pH and temperature fluctuation. The germinal vesicle (GV)-stage oocytes were released by puncturing ovarian follicles with a 27G needle under a stereomicroscope and immediately transferred into 50 ml droplets of culture medium. A small pipette (slightly larger than the diameter of the egg) was applied to denude oocyte through rapid pipetting cumulus-oocyte complex (COC). After being washed 3 times, about 5 oocytes were placed in a 30 µl droplet for IVM (Fig.1A).

Isolation of fetal mouse lung tissue

According to previous study, the second phase (days 11-14 of gestational age) involves accumulation of retinoic acid from maternal circulation (23). Thus, 3 female and 1 male mice were placed in a cage for mating. Observation of vaginal plug was considered as day 0 of gestational day. After 10 to 14 days of gestation, the pregnant mice were killed. After dissection, the uterine horns under sterile conditions was removed (Fig.1B). Lung tissues was separated from embryos (Fig.1C). After washing, individual lung tissue sample along with 5 immature oocytes were placed in a drop of 30 µl under mineral oil and incubated at 37° C in a humidified atmosphere of 5% CO₂ in air.

Assessment of oocyte maturation

Immature oocytes were classified as GV and GV breakdown (GVBD). Oocytes were cultured using inverted microscope (IX71, Olympus, Japan). Morphologic changes in the nucleus were evaluated with the release of the first polar body metaphase II (MII) as a measure of nuclear maturation of immature mouse oocytes.

Statistical analysis

The proportion of GV and GVBD oocytes reaching MII stage were compared among the three groups using analysis of variance (ANOVA). Correlation test were also used to evaluate the successful rate of IVM oocytes. All experiments were replicated 6 times. The statistical analysis was accomplished using the Statistical Package for the Social Sciences version 22 (SPSS, SPSS Inc., USA). A value of P<0.05 was considered statistically significant.

Results

As shown in Table 1, the proportion of GV oocytes (Fig.1C) reaching MII stage (Fig.1D) were 46, 65, and 56%, in control, I and II groups, respectively (P<0.05). The percentage of the oocytes remaining at the GV stage was higher in control group than the treatment groups (P < 0.05). There is no significant difference in percentage of GVBD and degenerated oocytes among three groups (P>0.05). However, there were differences in oocytes remaining at GV stage among three groups, indicating that the oocytes remaining at the GV stage of control group were more than those of I (P<0.000) and II (P=0.002) groups. There was a significant different between the control and II groups regarding percentage of oocytes reaching MII stage, meaning that the number of MII oocytes in the group II were more than the control group (P=0.04). Correlation analysis showed that there is a reverse relation between GV and MII oocytes (R=-0.4), suggesting that increasing the number of MII oocytes decreased the number of GV oocytes (P<0.000).



Fig.1: *In vitro* maturation (IVM) of immature oocytes and fetal mouse lung tissue co culture. **A.** Mouse embryos in the uterine horn oocytes at germinal vesicle (GV) stage, **B.** Immature oocytes before being co-cultured with the fetal mouse lung tissue, **C.** Immature oocytes with the fetal mouse lung tissue co-culture, and **D.** Oocytes at metaphase II (MII) stage.

Group	Cultured follicles	MII follicles	Degenerated follicles	GV follicles	GVBD follicles
	Number of immature follicles	n	n	n	n
Control	100	46*	4	29*,**	21
Ι	100	65*	2	11**	22
II	100	56	6	14*	24

Table 1: Effect of fetal mouse lung tissue co-	culture on in vitro maturation of mouse immature oocytes
--	--

GVBD; Germinal vesicle breakdown, MII; Metaphase II, *; P<0.05, and **; P<0.000.

Discussion

It has been indicated that some growth factors play an important role in the final follicular growth. Several studies have shown the effect of some growth factors on IVM and development of immature oocytes (24-26). Our results demonstrated that the percentage of oocyte reaching MII stage were significantly higher in both experimental groups as compared to the control group. This finding support this principle that several members of the FGF family and FGFRs are involved in early mouse lung development (15). It has been also shown that FGFs and FGFRs are expressed in the developing lung, indicating that they are the major regulators of lung growth and differentiation (27). FGF10 as a specific group of chain- polypeptides that is bonded to heparin has an essential role in cell growth and tissue repair. FGF10 also stimulates ovarian granulosa cell differentiation (28) that leads to the expression of LH receptors and proliferation of ovarian germ cells (29).

The establishment of an effective blood flow and formation of new blood vessels are necessary for ovarian tissue growth. Zeleznik et al. (30) have suggested that the selected follicles possess a more elaborate microvasculature than other follicles. In the studies by Yamashita et al. (31) and Schams et al. (32) they have indicated that preovulatory follicle benefits from FGF2 and vascular endothelial growth factor (VEGF), so an increase in precursor hormones and nutrients lead to the mature follicle. Since a similar study that examined co-culture system with lung tissue was not found, and since the fetal lung like FGFs has growth factors effect on oocyte maturation, we focused on the articles on growth factors or other somatic cells present in co-culture system. Previous studies have shown the effects of basic fibroblast growth factor (bFGF) and retinoic acid on follicle growth in vitro. In addition, 5 µM of retinol showing antioxidant effects during bovine embryo culture improved blastocyst development (33). Modified 3D culture system during 8 days of in vitro culture of human follicles showed that growth, survival and viability of follicles were improved by bFGF (25). Zhang et al. (34) have showed that exposure to 50 ng/ ml FGF10 supplementation affected bovine oocyte maturation and increased the percentage of MII oocytes as compared to the control group. Adding 50 ng/ml FGF10 increased the proportion of oocytes with compressed chromatin after 6 hours. In a study by Nilsson et al. (13), a prepubertal 4-day-old rat ovary was cultured and treated with 40 ng/ml bFGF for 14 days and their results have showed varying degrees of follicle development, but the total number of follicles persection and the size of the ovary did not change. Their findings have also indicated that ovaries cultured with bFGF showed a significant decrease in the number of primordial follicles to $15 \pm$ 2% that was followed by a corresponding increase to $85 \pm 3\%$ as compared to the control group.

In another study, Matos et al. (35) have been shown the maintained morphological integrity of caprine preantral follicles after 5-day treatment with 50 ng/ml FGF2, stimulation of primordial follicles, and the growth of activated follicles. However, in a study by Nandi et al. (36), they have showed that oocyte maturation was affected by FGF alone as compared to the control, epidermal growth factor (EGF) and FGF+EGF groups. The application of the somatic cell co-culture system used in this study was also mentioned in some other studies. In a study by Duszewska et al. (22), they have demonstrated that the cleavage of bovine premature oocytes cultured with Vero cells were more than the control group, greater percentage of fertilized bovine oocytes developed into blastocyst, as well as greater percentage of bovine oocytes achieved inner cell mass layer. However, oocyte maturation in vitro is a dose-dependent growth factor and completely affected by culture conditions. Nandi et al. (36) have also reported that 20 ng/ml FGF increased the rates of nuclear maturation, cleavage and blastocyst formation in culture conditions of buffalo oocytes. At the 2 and 4 µM concentrations of all-trans retinoic acid, the rate of GV oocytes reaching MII stage were higher than the control group without any effect on the development of the fertilized oocyte on 2-cell stage (37). The purpose of choosing the fetal mouse lung tissue in this study was the presence of FGF and retinoic acid in early embryonic life. The maturation rates in the both treatment groups were more than the control group. Also, in group I, fetal mouse lung tissue co-culture showed a better response to oocyte maturation as compared to group II. It may be caused by the effect of FGFinduced secreted from fetal mouse lung in the early embryonic day. Accordingly, the stage of organogenesis occurred on days 10 to 14, while the creation of fetal mouse lungs starting from the day 9 was affected by FGFs secreted from lung tissue that continued to day 17. High percentage of MI and MII oocytes in experimental groups as compared to the control group may be attributed to fetal lung tissue due to the abundance of FGF secreted (38).

Conclusion

This study suggests that use of co-culture system with fetal mouse lung tissue increased the percentage of GV oocytes reaching MII stage. Comparison of the two treatment groups revealed that the group with the earliest stages of lung development had better response than the other treatment group. The evaluation of fetal mouse lung tissue co-culture system on IVM of mouse immature oocytes was verified for the first time. However, more research is necessary for the sanitation of IVM of immature follicles.

Acknowledgments

We expressed our appreciation to Dr. Evangeline Foronda, Babol University of Medical Science (Babol, Iran) for the English editing. This project was financially supported by Islamic Azad University (Qaemshahr, Iran). There is no conflict of interest in this study.

References

- Cha KY, Koo JJ, Ko JJ, Choi DH, Han SY, Yoon TK. Pregnancy after in vitro fertilization of human follicular oocytes collected from nonstimulated cycles, their culture in vitro and their transfer in a donor oocyte program. Fertil Steril. 1991; 55(1): 109-113.
- Gilchrist RB, Thompson JG. Oocyte maturation: emerging concepts and technologies to improve developmental potential in vitro. Theriogenology. 2007; 67(1): 6-15.
- Eppig JJ, O'Brien MJ. Development in vitro of mouse oocytes from primordial follicles. Biol Reprod. 1996; 54(1): 197-207.
- Mahmoudi R, Subhani A, Salehnia M, Etesam F, Pasbakhsh P, Abolhasani F, et al. The effects of cumulus cells on in vitro maturation of mouse germinal vesicle stage oocytes. Iranian J Reprod Med. 2005; 3(2): 74-78.
- Martin GR. The roles of FGFs in the early development of vertebrate limbs. Genes Dev. 1998; 12(11): 1571-1576.
- Chaves RN, de Matos MH, Buratini JJr, de Figueiredo JR. The fibroblast growth factor family: involvement in the regulation of folliculogenesis. Reprod Fertil Dev. 2012; 24(7): 905-915.
- Berisha B, Sinowatz F, Schams D. Expression and localization of fibroblast growth factor (FGF) family members during the final growth of bovine ovarian follicles. Mol Reprod Dev. 2004; 67(2): 162-171.
- Ornitz DM, Itoh N. Fibroblast growth factors. Genome Biol. 2001; 2(3): REVIEWS3005.
- Itoh N. The Fgf families in humans, mice, and zebrafish: their evolutional processes and roles in development,

metabolism, and disease. Biol Pharm Bull. 2007; 30(10): 1819-1825.

- Knee RS, Pitcher SE, Murphy PR. Basic fibroblast growth factor sense (FGF) and antisense (gfg) rna transcripts are expressed in unfertilized human oocytes and in differentiated adult tissues. Biochem Biophys Res Commun. 1994; 205(1): 577-583.
- Ben-Haroush A, Abir R, Ao A, Jin S, Kessler-Icekson G, Feldberg D, et al. Expression of basic fibroblast growth factor and its receptors in human ovarian follicles from adults and fetuses. Fertil Steril. 2005; 84 Suppl 2: 1257-1268.
- Garor R, Abir R, Erman A, Felz C, Nitke S, Fisch B. Effects of basic fibroblast growth factor on in vitro development of human ovarian primordial follicles. Fertil Steril. 2009; 91(5 Suppl): 1967-1975.
- Nilsson E, Parrott JA, Skinner MK. Basic fibroblast growth factor induces primordial follicle development and initiates folliculogenesis. Mol Cell Endocrinol. 2001; 175(1-2): 123-130.
- Bazer FW, Spencer TE, Johnson GA, Burghardt RC, WuG. Comparative aspects of implantation. Reproduction. 2009; 138(2): 195-209.
- Bellusci S, Grindley J, Emoto H, Itoh N, Hogan BL. Fibroblast growth factor 10 (FGF10) and branching morphogenesis in the embryonic mouse lung. Development. 1997; 124(23): 4867-4878.
- Peters K, Werner S, Liao X, Wert S, Whitsett J, Williams L. Targeted expression of a dominant negative fgf receptor blocks branching morphogenesis and epithelial differentiation of the mouse lung. EMBO J. 1994; 13(14): 3296-3301.
- Karimpour Malekshah A, Heidari M, Parivar K, Azami NS. The effects of fibroblast co-culture and activin a on in vitro growth of mouse preantral follicles. Iran Biomed J. 2014; 18(1): 49-54.
- Heidari M, Malekshah AK, Parivar K, Khanbabaei R, Rafiei A. Effect of fibroblast co-culture on in vitro maturation and fertilization of mouse preantral follicles. Int J Fertil Steril. 2011; 5(1): 1-8.
- Maden M. The role of retinoic acid in embryonic and postembryonic development. Proc Nutr Soc. 2000; 59(1): 65-73.
- Morriss-Kay GM ,Ward SJ. Retinoids and mammalian development. Int Rev Cytol. 1999; 188: 73-131.
- Hidalgo CO, Díez C, Duque P, Facal N, Gómez E. Pregnancies and improved early embryonic development with bovine oocytes matured in vitro with 9-cis-retinoic acid. Reproduction. 2003; 125(3): 409-416.
- Duszewska AM, Reklewski Z, Pieńkowski M, Karasiewicz J, Modliński JA. Development of bovine embryos on Vero/ BRL cell monolayers (mixed co-culture). Theriogenology. 2000; 54(8): 1239-1247.
- Chytil F. Retinoids in lung development. FASEB J. 1996; 10(9): 986-992.
- Ghasemian F, Bahadori MH, Azarnia M, Soltani B, Nasiri E, Ahmadi Jalali Moghaddam M. The beneficial effect of fibroblastic growth factor on in vitro maturation and development ability of immature mouse oocytes. Open Reprod Sci J. 2010; 3: 1-5.
- Wang TR, Yan LY, Yan J, Lu CL, Xia X, Yin TL, et al. Basic fibroblast growth factor promotes the development of human ovarian early follicles during growth in vitro. Hum Reprod. 2014; 29(3): 568-576.
- Toori MA, Mosavi E, Nikseresht M, Barmak MJ, Mahmoudi R. Influence of insulin-like growth factor-i on maturation and fertilization rate of immature oocyte and embryo development in NMRI mouse with TCM199 and α-MEM medium. J Clin Diagn Res. 2014; 8(12): AC05-08.

www.SID.ir

Effect of Fetal Mouse Lung Tissue Co-Culture on IVM

- Cardoso WV, Itoh A, Nogawa H, Mason I, Brody JS. FGF-1 and FGF-7 induce distinct patterns of growth and differentiation in embryonic lung epithelium. Dev Dyn. 1997; 208(3): 398-405.
- Adashi EY, Resnick CE, Croft CS, May JV, Gospodarowicz D. Basic fibroblast growth factor as a regulator of ovarian granulosa cell differentiation: a novel non-mitigenic role. Mol Cell Endocrinol. 1988; 55(1): 7-14.
- Gospodarowicz D, Plouët J, Fujii DK. Ovarian germinal epithelial cells respond to basic fibroblast growth factor and express its gene: implications for early folliculogenesis. Endocrinology. 1989; 125(3): 1266-1276
- Zeleznik AJ, Schuler HM, Reichert LE Jr. Gonadotropinbinding sites in the rhesus monkey ovary: role of the vasculature in the selective distribution of human chorionic gonadotropin to the preovulatory follicle. Endocrinology.1981; 109(2): 356-362.
- Yamashita H, Kamada D, Shirasuna K, Matsui M, Shimizu T, Kida K, et al. Effect of local neutralization of basic fibroblast growth factor or vascular endothelial growth factor by a specific antibody on the development of the corpus luteum in the cow. Mol Reprod Dev. 2008; 75(9): 1449-1456.
- Schams D, Steinberg V, Steffl M, Meyer HH, Berisha B. Expression and possible role of fibroblast growth factor family members in porcine antral follicles during final mat-

uration. Reproduction. 2009; 138(1): 141-149.

- Livingston T, Eberhardt D, Edwards JL, Godkin J. Retinol improves bovine embryonic development in vitro. Reprod Biol Endocrinol. 2004; 2: 83.
- Zhang K, Hansen PJ, Ealy AD. Fibroblast growth factor 10 enhances bovine oocyte maturation and developmental competence in vitro. Reproduction. 2010; 140(6): 815-826.
- Matos MH, van den Hurk R, Lima-Verde IB, Luque MC, Santos KD, Martins F, et al. Effects of fibroblast growth factor-2 on the in vitro culture of caprine preantral follicles. Cells Tissues Organs. 2007; 186(2): 112-120.
- Nandi S, Ravindranatha BM, Gupta PS, Raghu HM, Sarma PV. Developmental competence and post-thaw survivability of buffalo embryos produced in vitro: effect of growth factors in oocyte maturation medium and of embryo culture system. Theriogenology. 2003; 60(9): 1621-1631.
- Eimani H, Tahaei LAS, Parivar K, Rezazadeh Valojerdi M, Kazemi Ashtiani S, Shahverdi AH, et al. The effect of granulesa cell co-culthre and retinoic acid on maturation and development of immature mouse oocyte in vitro. Anat Sci J. 2007; 5(19-20): 137-146.
- Heine UI, Munoz EF, Flanders KC, Roberts AB, Sporn MB. Colocalization of TGF-beta 1 and collagen I and III, fibronectin and glycosaminoglycans during lung branching morphogenesis. Development. 1990; 109(1): 29-36.