The Protein Extract of *Chlorella minutissima* Inhibits The Expression of *MMP-1*, *MMP-2* and *MMP-9* in Cancer Cells through Upregulation of *TIMP-3* and Down Regulation of *c-Jun*

Mugdha Kunte, M.Sc.¹, Krutika Desai, Ph.D.^{2*}

1. Department of Biological Sciences, NMIMS University, Vile Parle (W), Mumbai, India 2. Department of Microbiology, Mithibai College, Vile Parle (W), Mumbai, India

*Corresponding Address: Department of Microbiology, Mithibai College, Vile Parle (W), Mumbai 400056, India Email: krutika.desai@mithibai.ac.in

Received: 8/Jun/2017, Accepted: 16/Aug/2017

Abstract — Objective: Considering the bioactivities exhibited by microalgae, the effect of protein extract of *Chlorella minutissimma* (CP extract) was investigated on the expression of human matrix metalloproteinases-1 (*MMP-1*) in the breast cancer cell line MDA-MB231, and that of *MMP-2* and -9 in hepatocellular cancer cell line HepG2 at different expression levels. The study aimed identification and analysis of inhibitory activity of microalgal components extracted from *Chlorella minutissima* against human MMPs.

Materials and Methods: In this experimental study, we analysed the effect of *Chlorella* extracts on *MMP-1, -2*, and -9 expression at various levels. Gelatin zymography was performed to study the inhibitory effect of *Chlorella* exracts on human gelatinases at the activity level, followed by western blotting to analyse the expression of all three MMPs at the protein level. The similar effect at the mRNA level along with the probable mechanism underlying inhibition of MMPs was assessed using real-time polymerase chain reaction (PCR).

Results: The results reveal that the treatment with CP extract decreased the mRNA expression of *MMP-1*, *MMP-2*, and *MMP-9* by 0.26-, 0.29-, and 0.40-fold, respectively, at 20 µg/ml concentration as well as inhibited the activity of *MMP-2* and MMP-9 by 37.56 and 42.64%, respectively, at 15 µg/ml concentration. Additionally, upregulated mRNA expression of tissue inhibitor of metalloproteinases-3 (*TIMP-3*) by 1.68-fold was seen in HepG2 cells at 20 µg/ml concentration treatment group. However, CP extract did not induce any change in the mRNA expression of the *TIMP-1*, -2 and -4 in HepG2 and *TIMP-1*, -2, -3 and -4 in MDA-MB231 cells. Activator protein-1 (AP-1)-dependent *c-Jun*-mediated transcriptional regulation of *MMP-1*, -2, and -9 was also studied to elucidate the appropriate mechanism involved in the inhibition of MMPs.

Conclusion: The CP extract successfully inhibited *MMP-1, -2*, and -9 at different expression levels through *TIMP-3* upregulation and *c-Jun* downregulation.

Keywords: Chlorella spp., Matrix Metalloproteinases, Microalgae, Tissue Inhibitor of Metalloproteinases, Transcription Factor AP-1

Cell Journal (Yakhteh), Vol 20, No 2, Jul-Sep (Summer) 2018, Pages: 211-219 _

Citation: Kunte M, Desai K. The protein extract of chlorella minutissima inhibits the expression of MMP-1, MMP-2 and MMP-9 in cancer cells through upregulation of TIMP-3 and down regulation of c-Jun. Cell J. 2018; 20(2): 211-219. doi: 10.22074/cellj.2018.5277.

Introduction

Matrix metalloproteinases (MMPs) are the essential matrix proteases involved in migration and proliferation of cells by degradation of matrix components. The tightly regulated expression of MMPs is essential for organ development and tissue remodelling. Besides their positive roles in the normal physiology of cells, they have been identified as significant prognostic markers in diseases, such as cancer (1, 2). Hence, MMPs were identified as potential targets to control invasive and metastatic cancers (3, 4). Regulation of MMPs in pathogenic conditions using MMP inhibitors (MMPIs) derived from synthetic as well as a natural sources (5, 6) has been widely studied previously.

In the last decade, the focus of algal research has

shifted from the use of algae a superfoods to that as fuelproducing bio-resources. Besides their notable role in the bio-fuel production industry, their biological activities in medicine and pharmaceutics also have received increasing attention in recent years. *Chlorella* species were explored for various activities such as anti-inflammatory, antiangiogenic anti-proliferative, anti-ageing, anti-oxidant, apoptosis inducing activities (7-13).

Chlorella extracts were studied previously for their inhibitory activity on *MMP-1* expression induced by ultra violet β rays (UVB) exposure in skin fibroblasts (7). However, *Chlorella* extract has not been studied previously for its effect on MMPs from other classes. Hence, the current study aimed examination of inhibitory activity of protein extracts of *Chlorella minutissima*

(CP extracts) on human *MMP-2* and *MMP-9* at mRNA, protein, as well as activity level in HepG2 cells, and that of *MMP-1* at protein and mRNA expression levels in MDA-MB231 cells.

Additionally, the inhibition of gelatinase activity was studied with respect to change in expression of endogenous MMPIs (*TIMP-1*, -2, -3, and -4) in HepG2 cells. Additionally, the activator protein-1 (AP-1)-dependent transcriptional regulation of *MMP-1*, -2, and -9 was investigated using real-time polymerase chain reaction (PCR) analysis.

Materials and Methods

Maintenance of microalgae

In this experimental study, *C. minutissima* strain was obtained from Indian Agricultural Research Institute (IARI), New Delhi, India. The strain was cultured in BG-11 medium at 28°C in continuous light with 8 kLux intensity and maintained under similar conditions on BG-11 agar. The growing culture was harvested every alternate day to monitor its growth using parameters, such as wet cell biomass and protein content, for 15 days.

Preparation of Chlorella protein extract

C. minutissima was cultured in BG-11 medium for 15 days and then the wetbiomass was harvested and used for the further studies. The wet biomass obtained was soaked in MilliQ water and heated at a constant temperature of 60°C for 30 minutes. Further, it was incubated in 0.1 N NaOH for 15 minutes under similar temperature conditions. After alkali treatment cell debris were removed by centrifugation at 5000 \times g at 4°C and the cell free extract of proteins was used for further studies. The protein extract was fractionated using ammonium sulphate precipitation and the proteins were isolated into different fractions, such as 0-30%, 30-60%, and 60-90%. These precipitates were dialysed at 4°C using phosphate buffered saline pH=8. These dialysed fractions were evaluated using gelatin zymography for their activity against MMP-2 and -9 (data not shown).

The results revealed that the activity was retained in the crude protein extract; however, the isolated fractions lost the activity. Therefore, the crude protein extract was selected for further experiments in the current study. The extract obtained was filtered through 0.22-µm membrane filter, freeze dried and stored at -20°C until further use. The total protein concentration of the cell free extract (CP extract) was determined using Lowry's method.

Maintenance and culturing of human cell lines

The MDA-MB 231 and HepG2 cells were obtained from (authenticated and maintained by) National Centre for Cell Science (NCCS), Pune, India and was cultured in Dulbecco's modified Eagle's medium (DMEM, Sigma-Aldrich, USA) supplemented with heat-inactivated fetal bovine serum (FBS, 10%, Genetix, India) and antibiotic solution of penicillin streptomycin (Sigma-Aldrich, USA). Trypsin-EDTA (Gibco, India) was used for trypsinisation of cells. All materials used were of cell culture grade.

Cytotoxicity assay

The cytotoxicity of CP extract was estimated using the methyl thiazoltetrazolium (MTT) assay. MDA-MB231 cells $(7 \times 10^3/0.1 \text{ ml})$ and HepG2 cells $(3 \times 10^3/0.1 \text{ ml})$ were seeded in each well of a 96-well plate and exposed to different concentrations (5-40 µg/0.1 ml) of CP extract. The untreated cells were maintained as control. After 24 hours of incubation at 37°C, the cells were exposed to 20 μ l of MTT (2 mg/ml) for 3 hours. Supernatant from each well was discarded and dimethyl sulphoxide (DMSO, 100 µl/well) was added to dissolve formazan crystals formed during the incubation step. The absorbance of all treated and untreated samples was measured using a microplate reader (Bio-Rad) at 570 nm. The data is represented as mean \pm SD of three independent experiments in terms of percentage cellular viability of cells as compared to that of untreated controls.

Non reducing gelatin zymography

Gelatin zymography was performed to analyse the gelatinolytic activity of metalloproteinases (MMP-2 and -9) in HepG2 cells. The cells $(1 \times 10^{6}/\text{well})$ seeded in 6-well plates were exposed to different concentrations of CP extract (10, 15, 20, and 25 µg/ml) in serum free medium for 24 hours. The untreated cells were maintained as control. After incubation, cell-free supernatants were collected, aliquoted, and lyophilized, till further use. The samples were reconstituted in loading buffer and loaded on 10% sodium dodecyl sulfate (SDS)-polyacrylamide gel containing 1% gelatin. The MMPs resolved during electrophoresis were incubated in 25% Triton-X100 for 1hour, followed by incubation in substrate buffer [50 mM Tris-Cl (pH=7.6), 5 mM CaCl₂] at 37°C overnight followed by staining and destaining for visual observation of a clear band against the blue-colored background along with respective protein markers. The band intensities were quantified with Image J software 5.0.

Western blotting

The expression of *MMP-1* (in MDA-MB 231 cells), and *MMP-2* and -9 (in HepG2 cells) at the protein level was determined by western blotting. MDA-MB231 and HepG2 cells (1×10^{6} /well) were cultured in 6 well plates and exposed to CP extract (10, 15, 20 and 25 µg/ml) in FBS free medium. The untreated cell control was maintained. The monolayer of cells was removed using mechanical scraping and incubated in lysis buffer, at 4°C. Protein samples (20 µg) obtained were loaded on 10% reducing and denaturing polyacrylamide gel for separation. The proteins resolved on the gel were transferred to nitrocellulose membrane (Amersham Biosciences, UK) by electroblotting at 18 mA for 16 hours at 4°C.

The membrane was blocked with 5% BSA to avoid nonspecific binding followed by probing with rabbit monoclonal MMP-1, -2 and -9 antibodies (1:1000, Abcam, UK) on respective blots. After subsequent washing the blots were later probed with (1:2000, Merck, India). Goat anti-rabbit HRP-conjugated polyclonal secondary antibody. The chemiluminescence immerged through enzyme conjugated antibodies was captured on X-ray films. The mouse monoclonal β -actin antibody (Sigma-Aldrich, USA) was maintained as an internal and loading control. The band intensities were quantified with Image J software 5.0.

Real time polymerase chain reaction

The expressions of *MMP-1* (inMDA-MB231 cells), and MMP-2 and -9 (in HepG2 cells) at the mRNA level were analysed using real-time PCR. MDA-MB231 and HepG2 cells $(1 \times 10^{6}/\text{well})$ were cultured in a 6-well plates and exposed to different concentrations of CP extract (15 and 20 µg/ml) in FBS-free medium. The untreated cell control was maintained during the experiment. After incubation total RNA was extracted using TRI-reagent (Sigma-Aldrich) and followed with complementary DNA synthesis using TAKARA, India cDNA synthesis kit. The expression of MMP (MMP-1, -2, and -9), TIMPs (TIMP-1, -2, -3, and -4), and transcription factor AP-1 (*c-Jun* and *c-Fos*) at the mRNA level before and after the treatment was assessed using real time PCR on Applied Biosystems Step One Plus thermocycler.

The primer sequences used for amplification of genes were:

MMP-1-

F: 5'GAGCAAACACACTGACCTACAGGA3' R: 5'TTGTCCCGATGATCTCCCCTGACA3' *MMP-2-F:* 5'CAAGGACCGGTTTATTTGGC3'

R: 5'ATTCCCTGCGAAGAACACAGC3' MMP-9-

F: 5'TGGGCTACGTGACCTATGAC3' R: 5'CAAAGGTGAGAAGAGAGGGC3' *TIMP-1-*

*F: 5'*ACTTCCACAGGTCCCACAAC3' R: 5'CACTGTGCATTCCTCACAGC3' *TIMP-2-*

F: 5'ATGCACATCACCCTCTGTGA3' R: 5'CTCTGTGACCCAGTCCATCC3' *TIMP-3-*

F: 5'ACCGAGGCTTCACCAAGATG3' R: 5'CATCATAGACGCGACCTGTCA3' *TIMP-4-*

F: 5'TACCAGGCTCAGCATTAT3' R: 5'CCACTTGGCACTTCTTATT3' *c-Jun-*

F: 5'GGATCAAGGCGGAGAGGAAG3' R: 5'GCGTTAGCATGAGTTGGAAC3' *c-Fos-*F: 5'GGGGCAAGGTGGAACAGTTA3' R: 5'TTGGTCTGTCTCCGCTTGGA3' The *GAPDH* with a primer sequence of F: 5'CTCTCTGCTCCTCCTGTTCG 3' R: 5'ACGACCAAATCCGTTGACTC3' was used as an endogenous internal control.

The expression level of MMPs, TIMPs, *c-Jun* and *c-Fos* genes was normalised with *GAPDH*. An appropriate set of primers mentioned above were used to amplify respective genes using the following cycling conditions: 94° C for 5 minutes; followed by 35 cycles at 94° C for 30 seconds, 60° C for 30 seconds (for *MMP-1*, *-2*, *-9*, *TIMP-1*, *-2*, and *c-Jun* and *c-Fos*) and 58° C for 30 seconds (for *TIMP-3*, *-4*), and 72° C for 30 seconds extension. Melting curve was determined for the samples to verify change in products as per its specific melting temperature (T_m). The data is represented in the form of fold change calculated using $2^{-\Delta\Delta Ct}$ method in treated samples with respect to that in the untreated control. All sets of reactions were performed in triplicates.

Statistical analysis

All the experiments were performed in triplicates and results are expressed in terms of mean \pm SD. The data were analysed statistically using one-way ANOVA and Dunnett'spost test on GraphPad Prism software, (GraphPad software, Inc), P<0.05 was considered significant.

Results

Extraction and identification of *Chlorella* protein extract

The hot water extraction method was used to extract and isolate proteins from the wet cell biomass of C. *minutissima*. The total protein content of the CP extract was determined to be 5.086 mg/g of wet cell biomass.

Effect of *Chlorella* protein extract on cancer cell viability

Cell viability was determined by MTT assay. The cytotoxic concentrations of the CP extract in HepG2 and MDA-MB231 cells were assessed by this assay, and the observed MMP inhibition was confirmed to not due to cellular toxicity. The results are represented in terms of percentage cell viability. Treatment with CP extract did not induce a significant change in the viability of HepG2 or MDA-MB231 cells in the concentration range of 5-25 μ g/100 μ l. At 35 μ g/100 μ l concentration, the viability of HepG2 cells was significantly reduced to 85% (P<0.001, Fig.1A), and at 30 μ g/100 μ l concentration, viability of MDA-MB231 cells decreased to 77% (P<0.001, Fig.1B). Therefore, the concentrations between 10-25 μ g/100 μ l were selected for further study and precisely lower concentration was avoided to reduce technical errors.



Fig.1: Cytotoxicity induced by *Chlorella* protein (CP) extract in HepG2 and MDA-MB231 cells. The percentage of cell viability of **A.** HepG2 cells and **B.** MDA-MB231 cells after CP extract treatment at different concentrations. The results were analysed by one-way ANOVA and Dunnett'spost test. *; P<0.05, **; P<0.01, and ***; P<0.001.

Chlorella protein extract treatment reduced *MMP-9*, -2, and -1 expression at the protein level

All treated samples were analysed and expressed as percentage inhibition of MMP expression at the protein level in respective cancer cells, based on the intensity of the protein band. The band intensity of *MMP-1* and *MMP-9* were significantly (P<0.001) inhibited by 69.89 and 51.82% at 20 µg/ml concentration of CP extract. At 15 µg concentration of CP extract, band intensities of *MMP-2* was inhibited by 45.98% (Fig.2A-E). No further change was observed in band intensities of all three MMPs at the protein level. The data obtained from three individual experiments indicates that the expression of *MMP-1*, -2, and -9 was successfully reduced at the protein level after CP extract treatment.





Fig.2: Effect of *Chlorella* protein (CP) extract treatment on the expression of *MMP-9* and -2 in HepG2 cells and MMP-1 in MDA-MB 231 cells at the protein level. The effect of CP extract on **A.** MMP-9, MMP-2 in HepG2 cells and on **B.** MMP-1 in MDA-MB231 cells was observed by western blotting. The densitometric evaluation of **C.** MMP-9, **D.** MMP-2, and **E.** MMP-1 in terms of percentage band intensities depicted as mean ± SD of three individual experiments. The results were analysed by one-way ANOVA and Dunnett'spost test. ***; P<0.001, UC; Untreated control, and AUC; Area under curve.

Chlorella protein extract treatment inhibits enzymatic activity of *MMP-9* and *-2*

CP extract treatment causes inhibition of enzymatic activity of MMP-2 and -9 in HepG2 cells. The inhibition of enzymatic activity is expressed in terms of percentage inhibition of band intensities for MMP-2 and -9. CP extract at a concentration of 15 µg/ml induced inhibition of MMP-2 (~64 kDa) by 37.52% (P<0.001) and MMP-9 (~84 kDa) by 42.13% (P<0.001). CP extract at 20 µg/ml concentration inhibits MMP-2 activity by 45% (P<0.001) and MMP-9 activity by 52% (P<0.001). No further change was observed in the percentage of band intensities with respect to MMP-9 with increase in CP extract concentration. While, MMP-2 was further inhibited up to 60% at CP extract concentration of 25 µg/ml (Fig.3A-C). The results obtained from three independent experiments have confirmed that CP extract successfully inhibits enzymatic activity of both the gelatinases in HepG2 cells without inducing cytotoxicity.



Chlorella protein extract treatment downregulates mRNA expression of *MMP-1*, *-2*, and *-9*

The effect of CP extract on mRNA expression of *MMP-1*, *MMP-2* and *MMP-9* was studied using real time PCR. The analysis was done at 20 and 25 μ g/ml concentrations of CP extract which gave promising results in gelatin zymography and western blotting. The CP extract significantly reduced, *MMP-9* expression by 0.49- and 0.27-fold (Fig.4A) *MMP-2* by 0.29- and 0.20-fold (Fig.4B) and *MMP-1* by 0.26- and 0.24-fold (Fig.4C) at 20 and 25 μ g/ml concentrations, respectively. The study has confirmed that CP extract inhibits all three MMPs at the mRNA level of expression.



Fig.3: Effect of *Chlorella* protein (CP) extract on the gelatinolytic activity of *MMP-9* and -2 in HepG2 cells. **A.** Band of clearance in gelatin zymogram. Densitometric analysis of the gelatinolytic activity of **B.** *MMP-9*, and **C.** *MMP-2*, respectively, with respect to untreated control (UC). The results were analysed by one-way ANOVA and Dunnett'spost test. **; P<0.01, ***; P<0.001, and AUC; Area under curve.

Fig.4: Effect of *Chlorella* protein (CP) extract on mRNA expression of *MMP-1*, -2 and -9. The change in mRNA expression of *MMP-2* and -9 after CP extract treatment was studied using real-time polymerase chain reaction (PCR). Fold change in mRNA expression of **A**. *MMP-9* in HepG2 cells, **B**. *MMP-2* in HepG2 cells, and **C**. *MMP-1* in MDA-MB231 cells with respect to untreated control (UC). The results were analysed by one-way ANOVA and Dunnett'spost test. ***; P<0.001.

Chlorella protein extract treatment upregulates *TIMP-3* expression in HepG2 cells

The effect of CP extract on mRNA transcription from all the four TIMPs was studied using real-time PCR. Expression of TIMPs and that of MMPs is negatively correlated in cancer cells. Here, we investigated the change in transcription of TIMPs in HepG2 cells. CP extract treatment affected *TIMP-3* expression, although the expression of *TIMP-1*, -2 and -4 remained unaffected. Similarly, no change in expression of TIMPs was seen in MDA-MB231 cells (Fig. 5A-D). *TIMP-3* expression was upregulated by 1.68-fold and 2.44-fold at 20 and 25 µg/ ml concentration of CP extract respectively. This suggests that *TIMP-3* upregulation may have a positive impact on downregulation of *MMP-2* and -9 at the enzyme activity level in HepG2 cells.





Fig.5: Effect of *Chlorella* protein (CP) extract on mRNA expression of *TIMP-1, -2, -3,* and -4. The change in mRNA expression of TIMPs after CP extract treatment was studied using real-time polymerase chain reaction (PCR). Fold change in mRNA expression of A. *TIMP-1,* B. *TIMP-2,* C. *TIMP-3,* and D. *TIMP-4* with respect to untreated control (UC). The results were analysed by one-way ANOVA and Dunnett's post test. ***; P<0.001.

Effect of *Chlorella* protein extract on *c-Jun* and *c-Fos* expression in HepG2 and MDA-MB231 cells

The inhibitory effect of CP extract on mRNA transcription of the homodimers of the AP-1 transcription factor (*c-Jun* and *c-Fos*) was assessed by real-time PCR. Treatment with 20 µg/ ml of CP extract significantly (P<0.001) reduced the mRNA transcription of *c-Jun* by 0.43-fold in HepG2 cells and 0.3fold in MDA-MB231 cells, and the expression decreased further with increase in CP extract concentration (Fig.6A, B). However, the mRNA transcription of *c-Fos* was unchanged following a similar treatment (Fig.6C, D).



Cell J, Vol 20, No 2, Jul-Sep (Summer) 2018



Fig.6: Effect of *Chlorella* protein (CP) extract on mRNA expression of *c-Jun* and *c-Fos*. The change in mRNA expression of *c-Jun* and *c-Fos* after CP extract treatment was studied using real-time polymerase chain reaction (PCR). Fold change in mRNA expression of *c-Jun* in **A**. HepG2 cells, **B**. MDA-MB231 and *c-Fos* in, **C**. HepG2 cells, and **D**. MDA-MB231 with respect to untreated control (UC). The results were analysed by one-way ANOVA and Dunnett'spost test. ***; P<0.001.

Discussion

MMPs are present in several species ranging from viruses and bacteria to humans along with a conserved sequence at the active site motif. They are zinc binding proteases having structural similarities within the class. The MMPs discovered in humans until now are classified into sub-classes depending on their substrate specificity and structural arrangement. The class of gelatinases contains *MMP-2* and *MMP-9*. *MMP-1* falls into a group of collagenases possessing a capacity to cleave all types of fibrillar collagen proteins, such as type I, II, III, VII,VIII, and X. Structurally, collagenases have all the same subunits of gelatinases, except the fibronectin domain (4).

Several studies have revealed that the expression of MMP-1, -2, and -9 in cancers is regarded as a remarkable prognostic marker because activated MMP enzymes are almost undetectable under normal physiological conditions. MMP-2 and -9 are potential targets linked to migration and invasion of tumor cells (14-19). Similarly, over expression of MMP-1 in breast cancer (20, 21) is linked to their metastatic and invasive potential. These studies reported the unregulation of

MMP-2 and *MMP-9* in the HepG2 cell line and *MMP-1* in the MDA-MB231 cell lines. This makes these cell lines an important model to investigate the expression of *MMP-2*, -9 and -1 respectively.

MMP expression should be modulated to avoid uncontrolled degradation of matrix components and to minimize the severity of cancer. The regulation of MMPs has been studied before using various MMPIs (22). Several of them failed due to the lack of a "targeted approach" towards unregulated MMPs under pathogenic conditions. In the last few years, MMPI studies have improved with newer insights (23, 24). Considering this background, in the current study, we attempted to analyse the inhibitory potential of a natural component extract against human MMPs at different levels of regulation, and to analyse the specificity of the extracts towards MMPs from two distinct classes.

The enzymatic activity of MMPs is controlled by naturally expressed endogenous TIMPs. As compared to other compatible inhibitors found in other mammals, TIMPs found in the human genome are the most extensively studied class of proteins. TIMPs not only target MMPs, but also have a disintegrin and metalloproteinase with thrombospondin domains (ADAMs) as their non-MMP targets. Broadly, TIMPs are distributed as *TIMP-1*, -2, -3, and -4 according to their sequence simililarity, affinity for protein targets, and dynamic function in cell signaling cascades, excluding the inhibitory roles (25). Induction of TIMP expression in cancer cells to overcome MMP over expression has been widely studied in the past (26, 27).

In the current study, we investigated MMP-2 and -9 inhibition in their active forms with respect to the induction of mRNA transcription of TIMPs due to CP extract treatment. TIMP expression is negatively correlated to MMP expression in cancer cells. Here, we investigated the change in transcriptional expression of TIMPs in HepG2 and MDA-MB231 cells. Due to CP extract treatment TIMP-3 expression was observed to be affected only in HepG2 cells, although the expression of TIMP-1, -2, and -4 was changed in both the cell lines. Interestingly, TIMP-3 expression in HepG2 cells was dramatically upregulated in accordance with the increase in CP extract concentration. This suggests that there may be a positive impact from TIMP-3 upregulation on and MMP-2 and -9 downregulation, at the enzyme activity level, in HepG2 cells.

According to previous studies, *TIMP-3* is a broad spectrum inhibitor of MMPs. Mino et al. have reported that inibition of mRNA expression of *MMP-2* and -9 is linked to upregulation of *TIMP-3* expression (28). It has been reported that *TIMP-3* suppresses angiogenesis in lung cancer by downregulating *MMP-2* (29). A study conducted by Neill et al. (30) reported that induction of *TIMP-3* due to decorin (proteoglycan) certainly inhibits pro-angiogenic proteases, *MMP-2* and -9, in MDA-MB231 breast cancer cell lines. Several sources of evidence

suggest that the induction of *TIMP-3* may inhibit *MMP-2* and *MMP-9* activity in cancer cells. Hence, collectively, we can conclude that the dramatic increase in *TIMP-3* expression in HepG2 cells might be one of the causes for inhibition of the activity of *MMP-2* and *MMP-9*.

Activator protein-1 (AP-1) is a known positive regulator of MMPs. It has been indicated that, AP-1 regulates basal as well as transactivated (through external stimuli, such as phorbolmyristate acetate, cytolines, and growth factors) expression of MMPs (31). Several biological components have been studied previously for trascriptional regulation of MMPs through AP-1-activated pathways.

Chrysin, a chemical extracted from a plant source inhibited MMP-9 expression through inhibition of the AP-1 trasneription factor in gastric cancer cells (32). Berberine, a Chinese medicinal herb, inhibited migration of human smooth muscle cells by reducing mRNA transcription of MMP-2 and -9 by interrupting AP-1 and NF-κB signalling pathways (33). Accordingly, in the current study, we have investigated the change in expression of the AP-1 transcription factor due to CP extract treatment. AP-1 is a heterodimer consisting of proteins from the c-Jun and c-Fos families. MMP transcription can be regulated through activation, phosphorylation, and translocation of c-Jun and c-Fos family members. Additionally, c-Jun activation is regulated by c-Jun N-terminal kinases (JNKs)mediated pathway (34), and the extracellular signalregulated kinases (ERKs) phosphorylate and activate the c-Fos protein family (35).

Based on this background information, we investigated the change in expression of the members belonging to the AP-1 family. The CP extract treatment down regulated mRNA transcription of the *c-Jun* homodimer. However, *c-Fos* expression was not affected in either of the cell lines. This may indicate that *MMP-1*, *-2*, and *-9* were suppressed at the gene level by down regulation of *c-Jun* through a JNK-mediated pathway, although other members of this pathway should also be studied to understand the complete mechanism of action of CP extract.

Conclusion

Collectively, the results of this study confirmed that CP extract successfully inhibits *MMP-1*, *-2*, and *-9* expression at the mRNA as well as the protein level and inhibits *MMP-2* and *-9* in their active forms. This indicates that CP extract is not selective for gelatinases or *MMP-1* in terms of its inhibitory activity, although further studies are required to conclude its inhibitory activity towards MMPs from a distinct class. CP extract triggered MMP inhibition without inducing any cytotoxic effect. The current study also states that the activity of *MMP-2* and *MMP-9* may have been inhibited due to *TIMP-3* upregulation. Additionally, CP extract treatment also suppressed mRNA transcription of *c-Jun*, in turn suppressing *MMP-1*, *-2*, and *-9* at the gene level, although the complete underlying

mechanism is still to be investigated in the future. The current research would form a basis for conducting further studies on the identification and isolation of specific MMPIs from a natural source, which will help researchers achieve milestones in this field.

Acknowledgements

This study was conducted at Sunandan Divatia School of Science, NMIMS University, Mumbai. No separate funding bodies supported this work. The authors declare that they have no conflicts of interest.

Author's Contributions

M.K., K.D.; Contributed to conception and design, were responsible for overall supervision. M.K.; Contributed to all experimental work, data and statistical analysis, and interpretation of data, drafted the manuscript, which was revised by K.D. All authors have read and approved the final version of the manuscript.

References

- Mohammed MA, Seleim MF, Abdalla MS, Sharada HM, Abdel Wahab AH. Urinary high molecular weight matrix metalloproteinases as non-invasive biomarker for detection of bladder cancer. BMC Urol. 2013; 13: 25.
- Ren F, Tang R, Zhang X, Madushi WM, Luo D, Dang Y, et al. Overexpression of MMP family members functions as prognostic biomarker for breast cancer patients: a systematic review and metaanalysis. PLoS One. 2015; 10(8): e0135544.
- Vihinen P, Ala-aho R, Kähäri VM. Matrix metalloproteinases as therapeutic targets in cancer. Curr Cancer Drug Targets. 2005; 5(3): 203-220.
- Cathcart J, PulkoskiGross A, Cao J. Targeting matrix metalloproteinases in cancer: Bringing new life to old ideas. Genes Dis. 2015; 2(1): 26-34.
- Nuti E, Tuccinardi T, Rossello A. Matrix metalloproteinase inhibitors: new challenges in the era of post broad-spectrum inhibitors. Curr Pharm Des. 2007; 13(20): 2087-2100.
- Mannello F. Natural bio-drugs as matrix metalloproteinase inhibitors: new perspectives on the horizon? Recent Pat Anticancer Drug Discov. 2006; 1(1): 91-103.
- Chen CL, Liou SF, Chen SJ, Shih MF. Protective effects of Chlorella-derived peptide on UVB-induced production of MMP-1 and degradation of procollagen genes in human skin fibroblasts. Regul Toxicol Pharmacol. 2011; 60(1): 112-119.
- Sibi G, Rabina S. Inhibition of Pro-inflammatory mediators and cytokines by CXhlorella vulgaris extracts. Pharmacognosy Res. 2016; 8(2): 118-122.
- Kyadari M, Fatma T, Azad R, Velpandian T. Evaluation of antiangiogenic and antiproliferative potential of the organic extract of green algae Chlorella pyrenoidosa. Indian J Pharmacol. 2013; 45(6): 569-574.
- Yusof YA, Saad SM, Makpol S, Shamaan NA, Ngah WZ. Hot water extract of Chlorella vulgaris induced DNA damage and apoptosis. Clinics (Sao Paulo). 2010; 65(12): 1371-1377.
- Renju GL, Muraleedhara Kurup G, Saritha Kumari CH. Anti-inflammatory activity of lycopene isolated from Chlorella marina on Type II Collagen induced arthritis in Sprague Dawley rats. Immunopharmacol Immunotoxicol. 2013; 35(2): 282-891.
- Makpol S, Yaacob N, Zainuddin A, Yusof YA, Ngah WZ. Chlorella vulgaris modulates hydrogen peroxide-induced DNA damage and telomere shortening of human fibroblasts derived from different aged individuals. Afr J Tradit Complement Altern Med. 2009; 6(4): 560-572.
- Nakashima Y, Ohsawa I, Nishimaki K, Kumamoto S, Maruyama I, Suzuki Y, et al. Preventive effects of Chlorella on skeletal muscle atrophy in muscle-specific mitochondrial aldehyde dehydrogenase 2 activity-deficient mice. BMC Complement Altern Med. 2014; 14: 390.
- 14. Jang M, Lee ST, Kim JW, Yang JH, Yoon JK, Park JC, et al. A

feeder-free, defined three-dimensional polyethylene glycol-based extracellular matrix niche for culture of human embryonic stem cells. Biomaterials. 2013; 34(14): 3571-3580.

- Lee H, Baek SH, Lee JH, Kim C, Ko JH, Lee SG, et al. Isorhynchophylline, a potent plant alkaloid, induces apoptotic and anti-metastatic effects in human hepatocellular carcinoma cells through the modulation of diverse cell signaling cascades. Int J Mol Sci. 2017; 18(5): Pii: E 1095.
- Kunte M, Desai K. The Inhibitory effect of c-phycocyanin containing protein extract (C-PC Extract) on human matrix metalloproteinases (MMP-2 and MMP-9) in hepatocellular cancer cell line (HepG2). Protein J. 2017; 36(3): 186-195.
- Liang J, Wu S, Lo H, Hsiang C, Ho T. Vanillin inhibits matrix metalloproteinase-9 expression through down-regulation of nuclear factor-kappaB signaling pathway in human hepatocellular carcinoma cells. Mol Pharmacol. 2009; 75(1): 151-157.
- Dou CY, Cao CJ, Wang Z, Zhang RH, Huang L, Lian JY, et al. EFEMP1 inhibits migration of hepatocellular carcinoma by regulating MMP2 and MMP9 via ERK1/2 activity. Oncol Rep. 2016; 35(6): 3489-3495.
- Zhao X, Yang W, Pei F, Ma W, Wang Y. Downregulation of matrix metalloproteinases contributes to the inhibition of cell migration and invasion in HepG2 cells by sodium valproate. Oncol Lett. 2015; 10(1): 531-535.
- NgoMatip M, Pieme CA, AzabjiKenfack M, Moukette BM, Korosky E, Stefanini P, et al. Impact of daily supplementation of Spirulina platensis on the immune system of naïve HIV-1 patients in Cameroon: a 12-months single blind, randomized, multicenter trial. Nutr J. 2015; 14: 70.
- Singh LP, Mishra A, Saha D, Swarnakar S. Doxycycline blocks gastric ulcer by regulating matrix metalloproteinase-2 activity and oxidative stress. World J Gastroenterol. 2011; 17(28): 3310-3321.
- Zucker S, Cao J, Chen WT. Critical appraisal of the use of matrix metalloproteinase inhibitors in cancer treatment. Oncogene. 2000; 19(56): 6642-6650.
- Zhang C, Kim SK. Matrix metalloproteinase inhibitors (MMPIs) from marine natural products: the current situation and future prospects. Mar Drugs. 2009; 7(2): 71-84.
- Overall CM, López-Otín C. Strategies for MMP inhibition in cancer: innovations for the post-trial era. Nat Rev Cancer. 2002; 2(9): 657-672.

rch

- Brew K, Nagase H. The tissue inhibitors of metalloproteinases (TIMPs): an ancient family with structural and functional diversity. Biochim Biophys Acta; 2010; 1803(1): 55-71.
- Halder SK, Osteen KG, AlHendy A. Vitamin D3 inhibits expression and activities of matrix metalloproteinase-2 and -9 in human uterine fibroid cells. Hum Reprod. 2013; 28(9): 2407-2416.
- Hassan ZK, Daghestani MH. Curcumin effect on MMPs and TIMPs genes in a breast cancer cell line. Asian Pacific J Cancer Prev. 2012; 13(7): 3259-3264.
- Mino N, Takenaka K, Sonobe M, Miyahara R, Yanagihara K, Otake Y, et al. Expression of tissue inhibitor of metalloproteinase-3 (TIMP-3) and its prognostic significance in resected non-small cell lung cancer. J Surg Oncol. 2007; 9(3): 250-257.
- Miranda MS, Cintra RG, Barros SB, Mancini Filho J. Antioxidant activity of the microalga Spirulina maxima. Braz J Med Biol Res. 1998; 31(8): 1075-1079.
- Neill T, Painter H, Buraschi S, Owens RT, Lisanti MP, Schaefer L, et al. Decorin antagonizes the angiogenic network: concurrent inhibition of Met, hypoxia inducible factor 1α, vascular endothelial growth factor A, and induction of thrombospondin-1 and TIMP3. J Biol Chem. 2012; 287(8): 5492-5506.
- Benbow U, Brinckerhoff CE. The AP-1 site and MMP gene regulation: what is all the fuss about? Matrix Biol. 1997; 15(8-9): 519-526.
- Xia Y, Lian S, Khoi PN, Yoon HJ, Joo YE, Chay KO, et al. Chrysin inhibits tumor promoter-induced MMP-9 expression by blocking AP-1 via suppression of ERK and JNK pathways in gastric cancer cells. PLoS One. 2015; 10(4): e0124007.
- Liu SJ, Yin CX, Ding MC, Xia SY, Shen QM, Wu JD. Berberine suppresses in vitro migration of human aortic smooth muscle cells through the inhibitions of MMP-2/9, u-PA, AP-1, and NF-κB. BMB Rep. 2014; 47(7): 388-392.
- Davies C, Tournier C. Exploring the function of the JNK (c-Jun Nterminal kinase) signalling pathway in physiological and pathological processes to design novel therapeutic strategies. Biochem Soc Trans. 2012; 40(1): 85-89.
- Shaul YD, Seger R. The MEK/ERK cascade: From signaling specificity to diverse functions. Biochim Biophys Acta. 2007; 1773(8): 1213-1226.