The Role of Surgery for Local Recurrence of Renal Ewing's Sarcoma A Case Report

Abbas Basiri, Mahmoud Parvin, Navid eza Simaei, Amir Haji-Mohammadmehdi-Arbab

Keywords: bladder stones, percutaneous cystolithotripsy, children

Urol J (Tehran). 2006;4:250-2. www.uj.unrc.ir

INTRODUCTION

Primitive neuroectodermal tumor (PNET) is a rare malignancy mainly developed in the central nervous system and soft tissues of the children; however, primary occurrence of this tumor in the kidney has been recently reported.^(1,2) Histological diagnosis of such cases is challenging. We report a case with recurrence of left renal PNET that was treated successfully by mere tumor resection.

CASE REPORT

In February 2003, a 22-year-old man presented to our center with the

chief complaint of gross hematuria and pain in the left flank. Medical history and physical examination were unremarkable. Except for a microscopic hematuria, all laboratory tests had normal results. Ultrasonography revealed a 145 × 106-mm heterogeneous mass in the upper pole of the left kidney without the evidence of the metastatic disease. The findings were confirmed by CT scan with mild enhancement after contrast injection (Figure 1A). No tumoral tissue was detected in the inferior vena cava and the renal veins by magnetic resonance imaging. Chest radiography was also normal.



Figure 1. Diagnostic and follow-up CT scans. A, Primary renal tumor. B, Local recurrence. C, Last follow-up.

www.SID.ir

Urology and Nephrology Research Center, Shaheed Beheshti University of Medical Sciences, Tehran, Iran

> Corresponding Author: Abbas Basiri, MD No 44, 9th Boustan, Pasdaran Tehran 1666679951, Iran Tel: +98 21 2256 7222 Fax: +98 21 2256 7282 E-mail: basiri@unrc.ir

> > Received June 2006 Accepted September 2006

With the probable diagnosis of primary left kidney tumor, radical nephrectomy was performed. Histopathological assessment demonstrated uniform, small, round cells compatible with a stage II blastemal-type Wilms tumor. The surgical margins were free of tumor.

The patient underwent chemotherapy afterwards. But due to its complications, he refused to continue the treatment after 3 courses of chemotherapy. He was only followed with biochemistry, chest radiography, and abdominal and pelvic CT scan every 3 months. At the 12th postoperative month, CT scan revealed a 40 \times 50-mm mass with central calcification in the left renal fossa (Figure 1B). Since chemotherapy was refused by the patient, resection of the mass was performed. It was excised with a margin of the psoas muscle and the colon mesentery to which it was adhered (Figure 2A).

Pathologic findings were identical to the primary



kidney tumors such as a small round-cell sarcoma with extensive necrosis. Although all margins were tumor-free, immunohistochemical assessment of the specimen was performed due to the rosette formation (Figure 2B). The patient had a periodic acid-Schiff staining as focal fine granular cytoplasmic depositions. On immunohistochemistry, the neoplastic cells showed cytoplasmic staining of synaptophysin and cell membrane staining for microneme protein 2, but no reactivity was noted for desmin, actin, terminal deoxynucleotidyl transferase, and Wilms tumor suppressor (WT1). We additionally performed fluorescence in situ hybridization analysis which showed a split signal with DNA probes against the *EWS* gene, indicating *EWS* rearrangement.

Overall, histology and immunohistochemistry studies confirmed the diagnosis of Ewing's sarcoma (Figure 2C). The tumor was qualified as a renal PNET. Again, the patient refused further chemotherapy and he remained disease-free (within 16 months of followup after the recurrence). On the last follow-up visit, CT scan of the abdomen and pelvis was negative for metastatic disease or local recurrence (Figure 1C) and the patient was in a good general condition.

DISCUSSION

Primitive neuroectodermal tumor should be differentiated from other small round cell tumors of the kidney including lymphoma, small cell carcinoma, renal carcinoid tumor, neuroblastoma, rhabdomyosarcoma, blastema-predominant Wilms tumor, synovial sarcoma, and desmoplastic round cell tumor.⁽³⁾



Figure 2. A, Excised recurrent tumor. B, Sheets of small cells with rosette formation (hematoxylin-eosin, × 400). C, The cell membrane staining of MIC2 in the neoplastic cells (× 400)

Renal PNET was first reported by Mor and colleagues.⁽⁴⁾ Thereafter, a large number of renal Ewing's sarcoma/primitive neuroectodermal tumor (ES/PNET) cases have been reported in the literature.^(1,2,5,6) Most of the reported cases have been found among the young adults (average age, 28 years) with a slight male predominance (male-female ratio, 1.5 to 1).⁽⁶⁾

Radiotherapy or chemotherapy is not effective and the prognosis is poor in most cases with a reported 5-year survival rate of 45% to 55%.⁽⁷⁾ Current chemotherapy protocols used in the treatment of the children with ES/PNET is composed of induction and consolidation cycles with vincristine, doxorubicin, cyclophosphamide, ifosfamide, and etoposide.⁽⁶⁾ Further investigation of these regimens in patients with renal ES/PNET is warranted to determine their precise impact on the survival.

These tumors have a great tendency to develop metastases and subsequently a poorer prognosis is expected.^(2,5,6) Although there is a report of spontaneous regression of pulmonary metastasis following nephrectomy,⁽⁸⁾ most patients die within an average duration of 16 months despite chemotherapy.⁽⁶⁾

To our best knowledge, this is the first report in which mere resection of the recurrent tumor in the kidney could yield at least 16 months of being free of tumor or metastasis. Thus, re-exploration and removal of the recurrent tumor in cases of renal PNET may have a decisive role in the treatment of local recurrences of renal Ewing's sarcoma.

CONFLICT OF INTEREST

None declared.

REFERENCES

- Kuroda M, Urano M, Abe M, et al. Primary primitive neuroectodermal tumor of the kidney. Pathol Int. 2000;50:967-72.
- Thomas JC, Sebek BA, Krishnamurthi V. Primitive neuroectodermal tumor of the kidney with inferior vena cava and atrial tumor thrombus. J Urol. 2002;168: 1486-7.
- Parham DM, Roloson GJ, Feely M, Green DM, Bridge JA, Beckwith JB. Primary malignant neuroepithelial tumors of the kidney: a clinicopathologic analysis of 146 adult and pediatric cases from the National Wilms' Tumor Study Group Pathology Center. Am J Surg Pathol. 2001;25:133-46.
- 4. Mor Y, Nass D, Raviv G, Neumann Y, Nativ O, Goldwasser B. Malignant peripheral primitive neuroectodermal tumor (PNET) of the kidney. Med Pediatr Oncol. 1994;23:437-40.
- Karnes RJ, Gettman MT, Anderson PM, Lager DJ, Blute ML. Primitive neuroectodermal tumor (extraskeletal Ewing's sarcoma) of the kidney with vena caval tumor thrombus. J Urol. 2000;164:772.
- Jimenez RE, Folpe AL, Lapham RL, et al. Primary Ewing's Sarcoma/Primitive Neuroectodermal Tumor of the Kidney: A Clinicopathologic and Immunohistochemical Analysis of 11 Cases. 2002;26: 320-7.
- Casella R, Moch H, Rochlitz C, et al. Metastatic primitive neuroectodermal tumor of the kidney in adults. Eur Urol. 2001;39:613-7.
- Wada Y, Yamaguchi T, Kuwahara T, Sugiyama Y, Kikukawa H, Ueda S. Primitive neuroectodermal tumour of the kidney with spontaneous regression of pulmonary metastases after nephrectomy. BJU Int. 2003;91:121-2.