

## Pulmonary Aspergillosis in Ostrich

### Case Report

Sasani<sup>\*1</sup>, F., Khosravi, A.R.,<sup>2</sup> Dordari, S.,<sup>3</sup> Rajabi Moghadam, M.<sup>3</sup> and Hajibabaie, A.<sup>3</sup>

1. Veterinary Pathology Dept., Faculty of Veterinary Medicine, University of Tehran, Iran,  
P.O.Box:14155-6453

2. Mycology Dept., Faculty of Veterinary Medicine, University of Tehran

3. Private veterinarian

Received 26 Nov 2002; accepted 19 Mar 2003

Birds are routinely exposed to this fungus and only rarely become pathogenic and the lower respiratory tract is most severely affected by inhalation rout. There have been some recent reports about respiratory aspergillosis in ostrich (Kyoung 2001, Marks *et al* 1994). *Aspergillus* spp usually cause disease under condition of stress, immunosuppression, and prolonged treatment with antibiotics or massive exposure to the microorganism. Clinical signs of aspergillosis in ostriches are weight loss, lethargy and dyspnea (Kyoung 2001). Other less common forms of aspergillosis in birds are encephalitis, ophthalmitis, osteomyelitis, dermatitis and systemic form (Fitzgerald & Moisan (1995). *A.fumigatus* was isolated in some cases of respiratory aspergillosis (Campbell (1986). It has been reported that aspergillosis was observed in ostriches in the late 19th and early 20th centuries, but is relatively uncommon today (Terzich & Vanhooser (1993). This is the first case report of aspergillosis due to *A.fumigatus* in Iran. The ostriches of this report had received chloramphenicol and enrofloxacin for a long time (45 days) so these drugs may have been the cause of the pulmonary aspergillosis.

---

\* Author for correspondence. E.mail: fsasani@yahoo.com

### **Case history**

In a flock with 165 ostriches (*Struthio Camelus*) in Tehran province, two four-month-old male ostriches with respiratory distress and dyspnea, open mouth breathing, loss of appetite, weakness and retardation of growth were observed and finally died. Four weeks before appearing of clinical findings, enrofloxacin and chloramphenicol had been administered for 45 days.

Macroscopically, there were white and gray nodules about 1 to 4mm in diameter on the lungs and air sacs (Figure 1). In direct examination active form of mycelia were seen by DMSO and KOH. The mycelia were visible with routine hematoxylin and eosin as well as with the periodic acid-schiff (PAS) reaction with branched and septated form. (Figure 2).




Figure 1. White nodules of Aspergillosis on lungs and air sacs of ostriches




Figure 2. *Granulomatous pneumonia with mycelium of A.fumigatus (PAS) (400×)*

Microscopically many granulomatous nodules showed central necrosis with mycelia, mononuclear cells, giant cells and fibrosis. The nodules were cultured in Sabouraud dextrose agar (Merck) and incubated at 37°C for 48-72h and the causative agent media was diagnosed as *A.fumigatus* (Figures 3 and 4).




Figure 3. *Colony of A. fumigatus in Sabouraud culture media*




Figure 4. Mycelia of *A. fumigatus* (400×)

### References

- Campbell, T.N. (1986). Mycotic Disease. In: G. J.Harrison, L.R. Harrison (Eds.). *Clinical Avian Medicine and Surgery*. Pp: 464-466. Philadelphia, WB Saunders Co.
- Fitzgerald, S.D., Moisan, P.G. (1995). Mycotic rhinitis in an ostrich. *Avian Diseases* 39:194-196.
- Kyoung, O.H. (2001). Aspergillosis in an ostrich (*Struthio Camelus*). *Journal of Veterinary Clinics* 18:174-177.
- Marks, S.L., Stauber, E.H. and Ernstrom, S.B. (1994). Aspergillosis in an ostrich. *Journal of American Veterinary Medicine Association (JAVMA)* 204:784-785.
- Terzich, M., Vanhooser, S. (1993). Postmortem findings of ostriches submitted to the Oklahoma Animal Disease Diagnostic Laboratory. *Avian Diseases* 37:1136-1141.