



PCR detection of *Escherichia coli* O157:H7 directed from slaughtered cattle in Shiraz, Iran

Tahamtan^{*1}, Y.E., Pourbakhsh², S.A., Shekarforoush¹, S.S.

1. Food Hygiene Department of Veterinary Medicine, University of Shiraz, Shiraz, Iran

2. Department of Avian Disease Research & Diagnosis, Razi Vaccine and serum Research Institute, Karadj, Iran

Received 15 Aug 2005; accepted 7 Jan 2006

ABSTRACT

Escherichia coli O157:H7 lives in the intestines of healthy cattle, and can contaminate meat during slaughtering practices. Detection of the low infectious dosage of bacterium requires a sensitive method. We developed polymerase chain reaction (PCR) assays to detect the gene Stx₂ irrespective of the bacterial serotype. In this study, the detection limit of the PCR protocol in detecting Stx₂ in *E. coli* O157:H7 without prior enrichment was 10³ CFU/ml, and with 4 h prior enrichment in modified tryptic soy broth was 10² CFU/ml. During a period of 7 months (December 2004 to June 2005), 154 slaughtered cattle at Shiraz slaughterhouse, were randomly selected and examined for surface carriage of *E. coli* O157:H7 via detection the Stx₂ gene of verotoxigenic *E. coli* by PCR technique. *E. coli* O157:H7 was found in 14 of 145 (9.65%) feedlot cattle and in 1 of 9 (11.1%) dairy cull cows. This is the first report of the presence of *E. coli* O157:H7 in cattle from Iran.

Keywords: *Escherichia coli* O157: H7, Stx₂, Cattle, PCR

INTRODUCTION

E. coli O157:H7 is a worldwide threat to public health is implicated in hemorrhagic colitis, hemolytic uremic syndrome, and thrombotic thrombocytopenic purpura (Wells *et al* 1991, Currie *et al* 1999, Blackburn *et al* 2000, McCarthy *et al* 1998 & Nicole *et al* 2001). Approximately 52% of outbreaks have been associated with bovine products (Chapman *et al* 2001, Chinen *et al* 2001). Most illness has been associated with eating undercooked, contaminated

ground beef (Spika *et al* 1986, Carter *et al* 1987), suggesting that cattle are one of the most important sources of *E. coli* O157:H7 infection. An estimated 73,000 cases of infection and 61 deaths occur in the United States each year (Hancock *et al* 1994, Faith *et al* 1996). In the United States, the prevalence in cattle is reported to range from below 2% to around 45%, depending on the used culture techniques, the age of the animals, and the season in which samples are collected (Wells *et al* 1991, Hancock *et al* 1994, Zhao *et al* 1995, Faith *et al* 1996, Elder *et al* 2000). Animal prevalence between less than 1% and 13% have been reported in European countries

* Author for correspondence. E-mail: tahamtan@shirazu.ac.ir

(Garber *et al* 1995, Albihn *et al* 1997, Heuvelink *et al* 1999, Bonardia *et al* 1999, Andersen *et al* 2000). Isolation rates of bacterium from ground beef in Asia have so far been reported in Japan, India and China. Other Asian countries (like Iran) do not seem to be exempt from the *E. coli* O157:H7 infection (Son *et al* 1998, Durso *et al* 2004).

Most detection methods for verotoxigenic *E. coli* (VTEC) are based on the cultivation of serotype O157 (Gilgen *et al* 1998). The low infectious dosage of *E. coli* O157:H7 requires a sensitive detection method (Armstrong *et al* 1996). *E. coli* O157:H7 have two verotoxin genes (VT), shiga toxins (Stx₁ and Stx₂). Polymerase chain reaction (PCR) technique not only allows detection of VTECs irrespective of their serotype, but also detects all different variants of VT (Gilgen *et al* 1998).

The aim of the present study was to identification of *E. coli* O157:H7 from slaughtered cattle at Shiraz slaughterhouse, via the detection of Stx₂ gene using PCR. We were carried out this research for the first time in Iran.

MATERIALS AND METHODS

Sample collection. During a period of 7 months (December 2004 to June 2005), 145 feedlot cattle (reared intensively inside lots, weight 400-700 kg) and 9 dairy cull cows, slaughtered at Shiraz slaughterhouse, were selected randomly and examined for *E. coli* O157:H7. At the end of slaughtering practice, surface swabs were taken from the abdominal and lower parts of the carcasses with sterile wet cotton swabs. The samples were placed in sterile universal bottle and were transferred on ice to the laboratory within less than 2 h of collection. In the laboratory, 20 ml of tryptic soy broth (TSB, Oxoid) plus cefexime and tellurite (modified TSB) was added to each sample, which was then agitated vigorously by hand for 30 s. The suspensions were incubated over night at 37 °C prior to DNA extraction.

Identification. We used modified in-house enrichment method for DNA extraction as described by Holland *et al* (2000). The PCR protocols used in this study are referred to as Gannon *et al* 1992 and Holland *et al* 2000. Oligonucleotides primers used for PCR were synthesized by Roche, Germany, based on published data (Gannon *et al* 1992, Meyer *et al* 1992).

Amplification of Stx₂ sequences, including genetic variants, was performed using the primer pairs II-F (CCATGACAACGGACAGCAGTT) and II-R (CCTGTCAACTGAGCACTTTG), producing an amplicon of 779 base pairs. PCR was done with some modifications to the protocol described by Holland *et al* (2000) and in 25 µl reactions containing: 0.2 µM of each primers Stx₂, 0.2 µM of each deoxynucleotide triphosphates, 1X PCR buffer (10 mM Tris-HCl [pH 8.3], 50 mM KCl, 0.001% gelatin), 1.5 mM MgCl₂, 2.5 U of Taq DNA polymerase (Roche Molecular system), and 2 µl DNA template. Cycling condition provided by Techne Thermocycler (England) was as follows: 94 °C for 2 min, 30 cycle of 94 °C for 30 s, 57.5 °C for 45 s, and 72 °C for 30 s and a final extension at 72 °C for 10 min. PCR amplicons were run on 1% agarose gel as well ethidium bromide and visualized under UV illumination.

Sensitivity of the PCR. To assay the sensitivity of the PCR, 10-fold serial dilutions of *E. coli* O157:H7 (ATCC43895) in 0.1% peptone-water were prepared and the suspensions containing 10¹ to 10⁸ colony forming unit (CFU)/ml were amplified without prior enrichment and also 4 h of enrichment. The PCR procedure was done as described by Uyttendaele *et al* (1999) and Holland *et al* (2000) and the PCR products were run on 1% agarose gel.

Specificity of the PCR. To determine the specificity of the PCR assay, 7 different bacterial strains included *E. coli* serotypes O1, O2, O6 and O55, *Salmonella* spp., *Klebsiella* spp. and *Listeria* spp. were tested by PCR with conditions describe above.

RESULTS

A total number of 154 samples obtained from 154 animals were analyzed by selective enrichment in mTSB followed by PCR with Stx₂-specific gene primers.

E. coli O157:H7 was detected in 15 of 154 (9.74%) samples. Figure 1 shows the PCR protocol for some isolated *E. coli* O157:H7.




Figure 1. Products of PCR on some *E. coli* isolates. Lane 1: 100 bp ladder marker, Lane 2: Negative control, Lane 3-12: 779 bp positive samples.




Figure 2. PCR products of a 10-fold serial dilution of *E. coli* O157:H7. Lane 1: 100 bp ladder marker, Lane 2,3: Negative control, Lane 4-11: Bacterial suspensions containing 10¹ to 10⁹ CFU/ml, respectively.

The detection limit of the PCR protocol for Stx₂ in *E. coli* O157:H7 without prior enrichment was 10³ CFU/ml, and with 4 h prior enrichment in mTSB, was 10² CFU/ml (Figure 2).

Using primer set Stx₂ for amplification of the 779 bp, amplification products were observed for standard strain of *E. coli* O157:H7 but not observed for the other *E. coli* serotypes (4 strains), *Salmonella*, *Klebsiella* and *Listeria* (Figure 3).




Figure 3. PCR products of 8 bacterial strains. Lane 1: 100 bp ladder marker, Lane 2: Negative control, Lane 3: *E. coli* O157:H7, lanes 4-7: *E. coli* serotype O1, O2, O6 and O55, lane 8: *Salmonella* spp, Lane 9: *Klebsiella* spp, Lane 10: *Listeria* spp.

DISCUSSION

This study was designed to identify and to estimate the prevalence of *E. coli* O157: H7 in slaughtered cattle at Shiraz slaughterhouse, and to establish the sensitivity and specificity of PCR protocol. The PCR protocol, that we have described here, permits rapid, accurate, and inexpensive detection of these important toxin genes in clinical materials and possibly in contaminated foods. We identified *E. coli* O157:H7 on 9.74% of the examined cattle. We recovered the organism from 11.11% of healthy dairy cow and 9.65% of healthy feedlot carcasses. Other studies have shown similar rates of isolation. Wells *et al* (1991) presented Shiga-like-toxin-producing *E. coli* (SLTEC) in 8.4% of the adult cows in Wisconsin and Washington states. Similar results were reported by Canadian study of randomly selected cattle at slaughter. SLTEC was isolated

from 10.5% of beef cattle and 19.5% of dairy cows (Clarke *et al* 1998). In a study on healthy cattle in Germany, SLTEC was recovered from 17% of dairy cows and 9.4% of bulls (Montenegro *et al* 1990, Wells *et al* 1991). The prevalence rates of VTEC in cattle reported from other European countries were from less than 1% to 13% (Blanco *et al* 1996, Albihn *et al* 1997, Elder *et al* 2000, Andersen *et al* 2000). Some studies from the United States indicate that more than 45% of the cattle are carriers of *E. coli* O157:H7 (Chapman *et al* 2001, Dargatz *et al* 1997, Heuvelink *et al* 1999, Bonardi *et al* 1999, Elder *et al* 2000).

The rates of VTEC isolation in our study are very similar to those reported by Son *et al* (1998) in Malaysia, Rivas *et al* (2001) in Argentina and Terajima *et al* (2000) in Japan. For detection of lower numbers of the pathogen, prior enrichment will be necessary (Uyttendaele *et al* 1999). In this study, we could easily detect the Stx₂ genes in as little as 10³ CFU/ml without prior enrichment and 10² CFU/ml after enrichment of bacterial suspension. This result is in consistent with Pollard *et al* (1990). Results for positive controls of the above described experiment confirmed this detection limit.

In conclusion, this investigation, that was the first epidemiological survey on the occurrence of *E. coli* O157:H7 in healthy cattle in Iran, confirms that in Iran, as well as in other countries, cattle represent an important reservoir of this pathogen and the prevalence of the bacterium appears to be similar to those to the other parts of Asia (Albihn *et al* 1997, Gro *et al* 2001). Further studies are now needed to identify the vehicles of infection most relevant to our epidemiologic situation and the most appropriate measures to prevent the spread of this life-threatening food borne disease.

Acknowledgments

We thank Vice Chancellor for Research Affairs of Shiraz University for financial support of the project

No. 83-VE-1724-2280. The authors thank Dr. Shahsavandi, Department of Poultry Diseases Diagnosis, Razi Vaccine and Serum Research Institute, Dr. A. Hosseini, Department of Pathobiology, School of Veterinary Medicine, Shiraz University, Mrs. M. Masoudian and Mr. G.A. Niknia, School of Veterinary Medicine, Shiraz University, for technical assistance. We also thank Dr. S.M.H. Hosseini, Department of Immunology, Razi Vaccine and Serum Research Institute, Shiraz.

References

- Albihn, A., Zimmerman, U., Rehbinder, V., Jansson, C., Tysen, E. and Engvall, A. (1997). Enterohaemorrhagic *E. coli* (EHEC) a nation-wide Swedish survey of bovine faeces. *Epidemiological Sante Animal* 31(32): 131-133.
- Andersen, J.K., Nielsen, E.M. and Tegtmeier, C. (2000). Influence of age, sex and herd characteristics on VTEC O157 excretion in cattle from Danish dairy herds. VTEC 2000 *The 4th International Symposium and Workshop on Shiga Toxin Verocytotoxin*.
- Armstrong, G.L., Hollingsworth, J., and Morris, J.G. (1996). Emerging foodborne pathogens: *Escherichia coli* O157:H7 as a model of entry of a new pathogen into the food supply of the developed world. *Epidemiology Review* 18: 29-51.
- Blanco, M., Blanco, J.E., Blanco, J., Gonzalez, E.A., Mora, A., Prado, C., Fernandez, L., Rio, M., Ramos, J. and Alonso, M.P. (1996). Prevalence and characteristics of *Escherichia coli* serotype O157:H7 and other verotoxin-producing *E. coli* in healthy cattle. *Epidemiological Infection* 117: 251-257.
- Blackburn, C.W. and McCarthy, J.D. (2000). Modifications to methods for the enumeration and detection of injured *Escherichia coli* O157:H7 in foods. *International Journal of Food Microbiology* 55: 285-290.
- Bonardia, S.E., Maggia, A., Bottarella, M.L., Pacciarinib, A., Ansuinib, G., Vellinic, Morabitod, S. and Capriolid, A. (1999). Isolation of Verocytotoxin-producing *Escherichia coli* O157: H7 from cattle at slaughter in Italy. *Veterinary Microbiology* 67: 203-211.
- Carter, A.O, Borczyk, A.A. and Carlson, J.A.K. (1987). A severe outbreak of *Escherichia coli* O157:H7 associated

- hemorrhagic colitis in a nursing home. *New England Journal of Medicine* 317: 1496-1500.
- Chapman, P.A., Ellin, M., Ashton, R. and Shafique, W. (2001). Comparison of culture, PCR and immunoassays for detecting *Escherichia coli* O157 following enrichment culture and immunomagnetic separation performed on naturally contaminated raw meat products. *International Journal of Food Microbiology* 68: 11-20.
- Chinen, I., Jose, D., Taniel, T., Elizabeth, M., Liliana, H., Lound, G., Chillemi, S., Ledri, A., Baschkier, M., Scarpin, E. and Manfredi, M.R. (2001). Isolation and Characterization of *Escherichia coli* O157:H7 from Retail Meats in Argentina. *Journal of Food Protection* 64: 1346-1351.
- Clarke, R.S., McEwen, N., Harnett, H., Lior and C., Gyles (1988). *Abstract Annual Meeting American Society Microbiology* (No:48). Pp: 282.
- Currie, C.G., and Poxton I.R. (1999). The lipopolysaccharide core type of *Escherichia coli* O157:H7 and other non-O157 verotoxin-producing *E. coli* FEMS. *Immunology and Medical Microbiology* 24: 57-62.
- Dargatz, D.A., Wells, S.J., Thomas, L.A., Hancock, D.D. and Garber, L.P. (1997). Factors associated with the presence of *Escherichia coli* O157 in feces of feedlot cattle. *Journal of Food Protection* 60: 466-470.
- Durso, L.M., David, Smith, and Robert, W.H. (2004). Measurements of Fitness and Competition in Commensal *Escherichia coli* and *E. coli* O157:H7 strains. *Applied and Environmental Microbiology* 70: 6466-6472.
- Elder, R.O., Keen, K.J., Siragusa, G.R., Barkocy-Gallagher, G.A., Koohmaraie, M. and Laegreid, W.W. (2000). Correlation of enterohemorrhagic *Escherichia coli* O157 prevalence in feces, hides, and carcasses of beef cattle during processing. *Proceeding National of Academic Science U.S.A* 28: 2999-3003.
- Faith, N.G., Shere, J.A., Brosch, R., Arnold, K.W., Ansay, S.E., Lee, M.S. and Luchansky, J.B. and Kaspar, C.W. (1996). Prevalence and clonal nature of *Escherichia coli* O157:H7 on dairy farms in Wisconsin. *Applied and Environmental Microbiology* 62: 1519-1525.
- Gannon, P.J., King, Robin, K., Kim, Jong, Y. and Golsteyn, Thomas, E.J. (1992). Rapid and sensitive method for detection of Shiga-Like Toxin-Producing *Escherichia coli* in ground beef using the polymerase chain reaction. *Applied and Environmental Microbiology* 58: 3809-3815.
- Garber, L.P., Wells, D.D., Hancock, M.P., Doyle, J., Tuttle, J. A., and Zhao, T. (1995). Risk factors for fecal shedding of *Escherichia coli* O157:H7 in dairy calves. *Journal American Veterinary Medicine Association* 207: 46-49.
- Gilgen, M., Hubner, P., Hofelein, C., Luthy, J. and Candrin, U. (1998). PCR-based detection of verotoxin-producing *Escherichia coli* (VTEC) in ground beef. *Research Microbiology* 149: 145-154.
- Gro, J., Yngvild, W., Even, H., Odd I.B. and Hallgeir, H. (2001). *Escherichia coli* O157:H7 in faeces from cattle, sheep and pigs in the southwest part of Norway during 1998 and 1999. *International Journal of Food Microbiology* 65: 193-200.
- Hancock, D.D., Besser, T.E., Kinsel, M.L., Tarr, P.I., Rice, D.H. and Paros, M.G. (1994). The prevalence of *Escherichia coli* O157:H7 in dairy and beef cattle in Washington State. *Epidemiology Infection* 113: 199-207.
- Heuvelink, A.E., Zwartkruis-Nahuis, J.T.M., van den Biggelaar, F.L.A.M., van Leeuwen, W.J. and deBoer, E. (1999). Isolation and characterization of verocytotoxin-producing *Escherichia coli* O157 from slaughter pigs and poultry. *International Journal of Food Microbiology* 52: 67-75.
- Holland, J.L., Louie, L., Simor, A.E., and Louie M. (2000). PCR detection of *Escherichia coli* O157:H7 directly from stools: Evaluation of commercial extraction methods for purifying fecal DNA. *Journal of Clinical Microbiology* 38: 4108-4113.
- McCarthy, J., Holbrook, R. and Stephens, P.J. (1998). An improved direct plate method for the enumeration of stressed *Escherichia coli* O157:H7 from food. *Journal of Food Protection* 61: 1093- 1097.
- Meyer, T., Karch, H., Hacker, J., Bocklage, H., and Hesemann, J. (1992). Cloning and sequencing of a shiga like toxin-II-related gene from *Escherichia coli* O157:H7 strain 7279. *Zentralblatt Bakteriologie Mikrobiologie und Hygiene* 276: 176-188.
- Montenegro, M.M., Bulte, M., Trumpf, T., Aleksic, S., Reuter, G., Bulling, E. and Helmuth, R. (1990). Detection and characterization of fecal verotoxin-producing *Escherichia coli* from healthy cattle. *Journal of Clinical Microbiology* 28: 1417-1421.
- Nicole, T.P., Guy, P., Valerie, B., Bob, M., Jeremy, D.G., Debra, J.R., George, F.M., Peter, S.E., Jason, G.R., Heather, A.K., Gyorgy, P., Jeremiah, H., Sara, K., Adam, B., Ying, S., Leslie, M., Erik, J.G.N., Wayne, D., Alex, L., Eileen, T.D., Konstantinos, D.P., Jennifer, A.,

- Thomas, S.A., Jieyi, L., Galex, Y., David, C.S., Rodney, A.W. and Frederick, R.B. (2001). Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* 409: 529-533.
- Pollard, D.R., Johnson, W.M., Lior, H., Tyler, S.D. and Roze, K.R. (1990). Rapid and specific detection of verotoxin genes in *Escherichia coli* by the polymerase chain reaction. *Journal of Clinical Microbiology* 28: 540-545.
- Rivas, M., Isabil, C., Jose, D., Taniel, T., Miliwebsky, Liliana, E., Haydee, L., German, C., Ledri, S. Ariela, B., Scarpin M., and Eduardo, M. (2001). *Journal of Food Protection* 64: 1346-1351.
- Son, R., Shahilah, A.M.G.R., Zainori, A., Tadaaki, M., and Mitsuki, N. (1998). Detection on *Escherichia coli* O157:H7 in the beef market in Malaysia. *Applied and Environmental Microbiology* 64: 1153-1156.
- Spika, J.S., Parsons, J.E., Nordenberg, D., Wells, J.G., Gunn, R.A. and Blake, P.A. (1986). Hemolytic uremia syndrome and diarrhea associated with *Escherichia coli* O157:H7 in a day care center. *Journal of Pediatrics* 109: 287-291.
- Trajima, J., Izumiya, H., Wada, A., Tamura, K. and Watanabe, H. (2000). Molecular epidemiological investigation of enterohaemorrhagic *Escherichia coli* isolates in Japan. *Journal of Applied Microbiology Symposium Supplement* 88: 99S-105S.
- Uyttendaele, M., Boxstael, S. van and Debevere, J. (1999). PCR assay for detection of the *E. coli* O157:H7 *eae*-gene and effect of the sample preparation method on PCR detection of heat-killed *E. coli* O157:H7 in ground beef. *International Journal of Food Microbiology* 52: 85-95.
- Wells, J.G., Shipman, L.D., Greene, K.D., Sowers, E.G., Green, J.H., Cameron, D.N., Downes, F.P., Martin, M.L., Griffin, P.M., Ostroff, S.M., Potter, M.E., Tauxe, R.V. and Wachsmuth, I.K. (1991). Isolation of *Escherichia coli* serotype O157:H7 and other Shiga-like-toxin-producing *E. coli* from dairy cattle. *Journal of Clinical Microbiology* 29: 985-989.
- Zhao, T., Doyle, M.P., Shere, J. and Garber, L. (1995). Prevalence of enterohemorrhagic *Escherichia coli* O157:H7 in a survey of dairy herds. *Applied and Environmental Microbiology* 61: 1290-1293.