



## Aflatoxin M<sub>1</sub> contamination of raw and pasteurized milk produced in Sanandaj, Iran

Mohammadian<sup>\*1</sup>, B., Khezri<sup>1</sup>, M., Ghasemipour<sup>2</sup>, N., Mafakheri<sup>1</sup>, Sh., Poorghafour Langroudi<sup>3</sup>, P.

1. Agricultural and Natural Resources Research Center of Kurdistan Province, Sanandaj, Iran

2. Agricultural and Natural Resources Research Center of Kermanshah, Kermanshah, Iran

3. Agricultural and Natural Resources Research Center of Golestan, Gorgan, Iran

Received 20 Jul 2010; accepted 25 Sep 2010

---

### ABSTRACT

This study was conducted to evaluate and compare the levels of aflatoxin M<sub>1</sub> in raw and pasteurized milk samples during different seasons by Enzyme- Linked Immuno Sorbent Assay in Sanandaj, Iran. In 257 (94.49%) out of 272 milk samples the presence of aflatoxin M<sub>1</sub> was detected in concentrations ranging between 0.007 and 115.930 ng/l. AFM<sub>1</sub> level in 12 (4.4%) of positive samples were higher than the maximum tolerance limit (50 ng/l) accepted by Iran and European Union countries. Statistical evaluations showed that the differences between raw and pasteurized samples were not significant (p<0.05). There was no significant difference between spring and summer but the differences between other seasons were statistically significant. Winter samples with 22.35 ng/l and summer samples with 5.14 ng/l had the highest and lowest concentration, respectively (p<0.05). Since contamination of milk with aflatoxin is a potential risk for human health, milk and milk products should be controlled periodically for Aflatoxin contamination.

**Keywords:** Milk, Aflatoxin, measurement, ELISA

---

### INTRODUCTION

Aflatoxins are toxic by-products produced by the mold fungus varieties *Aspergillus flavus* and *Aspergillus parasiticus* (Harper 2003). AFB<sub>1</sub> is the most potent of the group and has been shown to be a potent carcinogen. According to the International Agency Research on Cancer (IARC), AFB<sub>1</sub> and AFM<sub>1</sub> are categorized as class 1A and class 2B carcinogens, respectively (International Agency for

Research on Cancer 1993, Maurice 2002). Many researchers have reported that there was a linear relationship between the amount of AFM<sub>1</sub> in milk and AFB<sub>1</sub> in feed consumed by animals (Bakirci 2001, Dragacci *et al* 1995). When animals eat foodstuff containing AFB<sub>1</sub>, it will be metabolized and excreted as AFM<sub>1</sub> in the urine and also in milk (Bakirci 2001, Gurbay *et al* 2006, Lopez *et al* 2003). Apart from this, exposure to aflatoxin can be through ingestion of contaminated milk containing AFM<sub>1</sub>. Occupational exposure to aflatoxins in

---


\*Author for correspondence. Email: mohammadian.baharak@gmail.com

agricultural workers, people working in oil mills and granaries have been reported (Ramdell & Eaton 1990). The parameters affecting levels of AFM<sub>1</sub> contamination in milk are the sources of animal feeds, ecologic and economic factors on the farm, and also farm management. It seems that the kind of animal feed and the harvesting time and temperature could be effective parameters in this regard (Kuiper 1999). According to the European Union and Codex Alimentarius the maximum level of AFM<sub>1</sub> in liquid milk and dried or processed milk products should not exceed 50 ng/kg (Codex Alimentarius Commissions 2001) but based on the US regulations it should not be higher than 500 ng/kg (Stoloff *et al* 1991, Tajik *et al* 2007). Susceptibility to aflatoxin is greatest in the young, and there are very significant differences between species, individuals, and gender. The toxicity of aflatoxin also varies according to many nutritional factors and recovery from protein malnutrition is delayed by exposure to aflatoxin. With increasing levels of aflatoxin in the diet, reduction in feed intake and growth of rate become severe. If aflatoxin levels are high enough, liver damage can occur (Williams *et al* 2004). The results of the numerous studies on the effect of milk processing on the concentration of AFM<sub>1</sub> are variable. It is resistant to thermal inactivation, pasteurization, autoclaving (Pittet 1998, Deshpande 2002, Creppy 2002, Park 2002, Soha *et al* 2006, Galvano *et al* 1996) and production of yoghurt, cheese, cream, milk powder, or butter does not lead to loss of AFM<sub>1</sub>, although it is redistributed differentially into the products resulting from these processes (Kamkar 2005, Food and Agriculture Organization/World Health Organization). In this work, data on the natural occurrence of AFM<sub>1</sub> in milk produced in Sanandaj is presented for the first time in order to determine the presence and levels of AFM<sub>1</sub> in milk during different seasons and to compare the results with maximum tolerable limits in milk that may be accepted by the European Union.

## MATERIALS AND METHODS

**Sampling.** During the years 2006-2007, 240 raw and 32 pasteurized milk samples were obtained from Sanandaj Dairy and Milk Company. Samples of raw milk were taken with raw milk jar samplers from raw milk tankers, arriving directly from traditional dairy farm and after pasteurization process, samples of pasteurized milk were taken from pasteurized milk tankers. All of milk samples were collected based on Iranian National Standards (INS) milk sampling method No.419 (Institute of Standard and Industrial Research of I.R. 1965). The samples were prepared for analysis of AFM<sub>1</sub> with the competitive enzyme-linked immunosorbent assay (ELISA) method described by R-Biopharm AG, Darmstadt, Germany (Aflatoxin M130/15 Kit) by ELISA Reader ELX800, USA.

**Samples Preparation.** Milk samples were centrifuged for degreasing at 3500 G for 10 min at 10°C. The upper cream layer was removed by aspirating through a Pasteur pipette. The skimmed milk was used directly in the test (100 µl per well).



**Figure 1.** Level of Aflatoxin M1 in analyzed milk samples.

ELISA test procedure: 100 µl standard solutions and prepared samples in separate wells were added and incubated for 30 min at room temperature in the dark. After the washing steps, 100 µl of the enzyme conjugate was added to each well and incubated for 15 min at room temperature in the dark. Again the washing steps were done, then 100 µl of substrate/chromogen were added to each well and incubated

for 15 min at room temperature in the dark. 100 µl of the stop solution was added to each well, mixed gently and the absorbance measured at 450 nm against an air blank within 15 min after addition of stop solution.

**Statistical methods.** sequential differences among means were calculated at the level of  $p < 0.05$ , using SPSS 12 software.

## RESULTS

In general 12 (4.67%) of positive samples, 10 (4.17%) of raw milk and 2 (6.25%) of pasteurized milk samples, was exceeded the legal level of AFM<sub>1</sub> accepted by Iran and European countries (Codex Alimentarius Commissions 2001, Institute of Standard and Industrial Research of I.R.2002).

**Table 1.** Aflatoxin M<sub>1</sub> concentration of raw and pasteurized milk samples.

Type	No.of Samples	Negative	Positive	Mean Concentration (ng/l)	SD
Raw	240	14	226	12.65 <sup>a</sup>	17.76
Pasteurized	32	1	31	12.43 <sup>a</sup>	17.53
Total	272	15	257	12.63	17.70

**Table 2.** Distribution of Aflatoxin M<sub>1</sub> concentration by season.

Seasons	No.of Samples	Mean Concentration (ng/l)	SD	Maximum	Minimum
Spring	49	9.40 <sup>a</sup>	13.09	52.24	0.000
Summer	65	5.14 <sup>ba</sup>	5.14	33.50	0.000
Autumn	79	11.01 <sup>c</sup>	12.39	60.31	0.000
Winter	79	22.35 <sup>d</sup>	25.59	115.93	0.410
Total	272	12.60	17.67	115.93	0.000

According to Table 1 Aflatoxin M<sub>1</sub> was found in 257 (94.49%) of the examined samples with no significant differences between raw and pasteurized milk samples. The levels of aflatoxin M<sub>1</sub> were different between seasons. The overall contamination levels in winter were higher than other seasons.

## DISCUSSION

There are few literature data on the occurrence of AFM<sub>1</sub> levels in milk and milk products in Iran (Tajik *et al* 2007, Kamkar 2005, Alborzi *et al* 2006, Tajkarimi *et al* 2007, Karim *et al* 1998, Oveisi *et al* 2007, Sefidgar *et al* 2008). In one study in 1998, out of 73 milk samples delivered to Tehran milk pasteurization plants, 60(82.2%) were contaminated. All contaminated samples had a level of AFM<sub>1</sub> above 50ng/l (Karim *et al* 1998). The incidence of the AFM<sub>1</sub> level in 328 milk and infant milk products in Tehran were 96.3%. The presence of AFM<sub>1</sub> in pasteurized liquid milk (n=128), infant formula (n=120) and milk-based cereal weaning food was 72.2±23.5, 7.3±3.9 and 16.8±12.5 ng/kg, respectively (Oveisi *et al* 2007). In Shiraz (Southern Iran) 624 pasteurized milk samples were analyzed. AFM<sub>1</sub> was found in 100% of the examined samples, 390 of them with contamination less than 45 ng/kg, 123 samples (19.7%) contained 45-50 ng/kg, 94 samples(15.1%) had 50-80 ng/kg and 17 (2.7%) of samples contained more than 80 ng/kg (Alborzi *et al* 2006). Tajik *et al.* (2007) found a high incidence rate of AFM<sub>1</sub> (100%) in 144 raw and pasteurized milk samples in Urmia, Iran of which the level in 6.25% of samples was higher than 50 ng/l (Tajik *et al* 2007). In 85 of 111 raw milk samples (76.6%) examined in Sarab city of Iran, AFM<sub>1</sub> was detected in concentrations ranging between 0.015 and 0.28 µg/l and in 40% of positive samples, its level was higher than 50 ng/l (Kamkar 2005). Lopez *et al.* (2003) suggested that levels of AFM<sub>1</sub> in samples of milk produced in Argentina were found to be very

low and in no case did the levels exceed the recommended limits for milk products (0.05µg/l).

**Table 3.** The prevalence of aflatoxin M<sub>1</sub> contamination in raw milk samples

Countries	No. of Raw Samples	Percent of Samples > 50 ng/l	References	
<b>Iran</b>	Sanandaj	240	4.17	
	Urmia	72	12.5	Tajik et al. (2007)
	Sarab	111	40	Kamkar (2005)
	Tehran	73	82.2	Oveisi et al. (2006)
	Babol	120	56.7	Sefidgar et al. (2008)
<b>Greece</b>	81	0	Markaki et al. (1997)	
<b>Argentina</b>	56	0	Lopez et al. (2003)	
<b>Italy</b>	296	1.7	Nachtmann et al. (2007)	

The incidence of AFM<sub>1</sub> contamination in raw milk analyzed in Portugal was 80.6%, 17 samples (54.8%) contained 0.005-0.010 µg/l, two samples (6.5%) had 0.011-0.02 µg/l, and six samples (19.3%) contained 0.021-0.050 µg/l (Martins *et al* 2000). Out of 85 milk samples in Ankara, Turkey, 75 samples (88.23%) were contaminated with AFM<sub>1</sub>; of which 27 samples (36%) were contaminated with concentrations <50 ng/l and 48 samples (64%) exceeded the legal level of Aflatoxin M<sub>1</sub> in milk (Çelik *et al* 2005). During 2005, out of 45 raw milk samples analyzed in Italy, only one sample was positive to the ELISA test, but the confirmation test values (0.035 µg/l) were below the legal limit (Decastelli *et al* 2007). In Korea, the incidence of AFM<sub>1</sub> in liquid milk was 76% with a mean concentration of 18 pg/g (Kim *et al* 2000). In this study the level of AFM<sub>1</sub> in raw and pasteurized milk samples showed no significant differences (p<0.05). As compared to other studies the contamination level of raw samples of Sanandaj were lower than those of Urmia (Tajik *et al* 2007), Sarab (Kamkar 2005), Babol (Sefidgar *et al* 2008) and Tehran (Oveisi *et al* 2007) in Iran and it was higher than Argentina (Lopez 2003) and Greece

(Markaki 1997) where in no case did the levels exceed the recommended limits. The results of the AFM<sub>1</sub> level derived from pasteurized samples were lower compared with other studies conducted in Shiraz (Alborzi *et al* 2006), Tehran (Oveisi *et al* 2007), Tabriz (Movassagh Ghazani 2009) India (Rastogi *et al* 2004), Brazil (Oliveira *et al* 2007), Turkey (Unusan 2006) and United State of America (Cathey *et al* 1994) and higher than samples examined in Urmia (Tajik *et al* 2007), Mashhad (Karimi *et al* 2007), Italy (Nachtmann *et al* 2007) and Argentina(Lopez *et al* 2003). The disagreement among the findings can be a result of differences in the level of contamination of consumed food stuffs. Seasonal evaluation of the data indicate meaningful differences in the contamination level (p<0.05).

**Table 4.** The prevalence of aflatoxin M<sub>1</sub> contamination in pasteurized milk samples.

Countries	No. of Pasteurized Samples	Percent of Samples > 50 ng/l	References	
<b>Iran</b>	Sanandaj	32	6.25	Tajik et al. (2007)
	Urmia	72	0.0	Alborzi et al.(2006)
	Shiraz	624	17.8	Oveisi et al. (2006)
	Tehran	128	78	Movassagh Ghazani(2009)
	Tabriz	50	62	Karimi et al.(2007)
	Mashhad	110	5.4	
<b>Turkey(Anatoli)</b>	129	47	Unusan (2006)	
<b>USA</b>	85	64	Cathey et al. (1994)	
<b>India</b>	12	33	Rastogi et al. (2004)	
<b>Turkey(Ankara)</b>	27	3.7	Gurbay et al. (2006)	
<b>Argentina</b>	16	0	Lopez et al. (2003)	
<b>Brazil</b>	57	7.4	Oliveira et al. (2007)	
<b>Italy</b>	316	0.6	Nachtmann et al. (2007)	

Some other studies also indicate a significant difference between winter and other seasons (Ghiasian *et al* 2007, Tajkarimi *et al* 2007, Kamkar 2005). This variation can be due to the fact that grass, pasture, weed, and rough feeds were found

more commonly in spring and summer than in winter. Therefore, obtained results were in agreement with prior studies. In order to achieve a low level of AFM<sub>1</sub>, it is essential to conduct training programs for producers about the toxicity potential of aflatoxins, reduce the concentration of AFB<sub>1</sub> in animal feed by good manufacturing and storage and integrating it with Hazard Analysis and Critical Control Points (HACCP) based safety program and doing regular monitoring of milk and milk products.

## References

- Alborzi, S., Pourabbas, B., Rashidi, M., Astaneh, B. (2006) Aflatoxin M<sub>1</sub> contamination in pasteurized milk in Shiraz (south of Iran). *Food Control* 17 (7): 582-584
- Bakirci, I. (2001). A study on the occurrence of aflatoxin M<sub>1</sub> in milk and milk products produced in Van province of Turkey. *Food Control* 12(1): 47-51
- Cathey, C.G., Huang, Z.G., Sarr, A.B., Clement, B.A. and Phillips, T.D. (1994) Development and evaluation of a minicolumn assay for the detection of aflatoxin M<sub>1</sub> in milk. *Journal of Dairy Science* 77: 1223-1231
- Çelik, T.H., Sarımehtetoğlu, B. and Küplülü, Ö. (2005). Aflatoxin M<sub>1</sub> contaminations in pasteurized milk. *Veterinarski Arhiv* 75(1): 57-65
- Codex Alimentarius Commissions. Comments submitted on the draft maximum level for aflatoxin M<sub>1</sub> in milk. Codex Committee on Food Additives and Contaminant (2001) 33rd sessions, Hauge, The Netherlands [ftp://ftp.fao.org/codex/ccfac33/fa0120e.pdf](http://ftp.fao.org/codex/ccfac33/fa0120e.pdf)
- Creppy, E.E. (2002). Update of survey, regulation and toxic effect of mycotoxins in Europe. *Toxicology Letters* 127(1-3), 19-28
- Decastelli, L., Lai, J., Gramaglia, M., Monaco, A., Nachtmann, C. and Orlando, F. (2007). Aflatoxins occurrence in milk and feed in Northern Italy during 2004–2005. *Food Control* 18: 1263-1266.
- Deshpande, S.S. (2002). Fungal toxins. In: S.S. Deshpande Editor, Handbook of Food Toxicology, Marcel Decker, New York. 387-456 ISBN: 0-203-90896
- Tajik, H., Razavi Rohani, S.M., and Moradi, M., Detection of Aflatoxin M<sub>1</sub> in Raw and Commercial Pasteurized Milk in Urmia, Iran. *Pakistan Journal of Biological Science* 10(22): 4103-4107
- Dragacci, S., Gleizes, E., Fremi, J.M. and Candlish, A.A.G. (1995). Use of immunoaffinity chromatography as a purification step for the determination of aflatoxin M<sub>1</sub> in cheeses. *Food Additives and Contaminants* 12(1): 59-65
- Food and Agriculture Organization/World Health Organization. Evaluation of certain mycotoxins in food. 56th report of the joint. (2002) FAO/WHO exports committee on Food Additives. WHO technical report. series no.906.
- Galvano, F., Galorafo, V., and Galvano, G. (1996). Occurrence and stability of aflatoxin M<sub>1</sub> in milk and milk products: a worldwide review. *Journal of Food Protection* 59(10): 1076-1090
- Ghiasiian, S.A., Maghsood, A.H., Neyestani, T.R. and Mirhendi, S.H. (2007). Occurrence of aflatoxin M<sub>1</sub> in raw milk during the summer and winter seasons in Hamedan, Iran. *Food Safety* 27:188-198
- Gurbay, A., Aydin, S., Girgin, G., Engin, A.B. and Sahin, G. (2006). Assessment of aflatoxin M<sub>1</sub> levels in milk in Akara, Turkey. *Food Control* 17: 1-4
- Harper, A. (2003). An occasional contaminant of feed grains. *LivestockUpdate*. Virginia Cooperative Extension. <http://en.engormix.com/MA>.
- Institute of Standard and Industrial Research of I.R. Milk Sampling methods. (1965). National Standard No.419.
- Institute of Standard and Industrial Research of I.R. Iran. Maximum Tolerated Limits of Mycotoxins in Foods and Feeds. (2002), National Standard No. 5925.
- International Agency for Research on Cancer. Monographs on the evaluation of carcinogenic risks to humans. (1993) 56: 245-395. <http://monographs.iarc.fr/ENG/Monographs/vol56/index.php>
- Kamkar, A. (2005). A study on the occurrence of aflatoxin M<sub>1</sub> in raw milk produced in Sarab city of Iran. *Food Control*. 6(7): 593-599
- Karim, G., Bokaiee, S. and Khorasani, K. (1998). Study on the contamination of raw bulk milk with aflatoxin M<sub>1</sub> in Tehran area using ELISA method. *Pajoohesh and Sazandeghi* 40- 42: 163-165
- Karimi, G., Hassanzadeh, M., Teimuri, M., Nazari, F. and Nili, A. (2007). Aflatoxin M<sub>1</sub> contamination in pasteurized milk in Mashhad, Iran. *Iranian Journal of Pharmaceutical Science* 3(3): 153-156. [www.ijps.ir](http://www.ijps.ir)
- Kim, E.K., Shon, D.H., Ryu, D., Park, J.W., Hwang, H.J. and Kim, Y.B. (2000). Occurrence of aflatoxin M<sub>1</sub> in Korean dairy products determined by ELISA and

- HPLC. *Food Additives and Contaminants* 17: 59-64 (6).
- Kuiper, G.T. (1999). Approaches to the risk analysis of mycotoxins in the food supply. 3<sup>rd</sup> Joint FAO/WHO/UNEP International Conference on Mycotoxins. Tunis, Tunisia Myc-Conf/99/7a, Pp 4-6.
- Lopez, C.E., Ramos, L.L., Ramadan, S.S. and Bulacio, L.C. (2003). Presence of aflatoxin M<sub>1</sub> in milk for human consumption in Argentina. *Food control* 14 (1): 31-34.
- Markaki, P. and Melissari, E. (1997). occurrence of Aflatoxin M<sub>1</sub> in commercial pasteurized milk determined with ELISA and HPLC. *Food Additives and contaminants* 14: 451-456.
- Martins, M.L. and Martins, H.M. (2000). Aflatoxin M<sub>1</sub> in raw and ultra high temperature treated milk commercial in portugal. *Food Additives and contaminants* 17: 871-874.
- Maurice, O.M. (2002). Risk assessment for aflatoxins in foodstuffs *International Biodeterioration and Biodegradation* 50(3-4): 137-142.
- Movassagh Ghazani, M.H. (2009). Aflatoxin M<sub>1</sub> contamination in UHT milk in Tabriz (northwest of Iran). *Toxicology Letters* 189: S 228.
- Nachtmann, C.A., Gallina, M., Ferro, G.L. and Decastelli, L. (2007). Regional monitoring plan regarding the presence of aflatoxin M<sub>1</sub> in pasteurized and HT milk in Italy. *Food Control* 18: 623-629
- Oliveira, C.A.F. and Ferraz, J.C.O. (2007). Occurrence of aflatoxin M<sub>1</sub> in pasteurised, UHT milk and milk powder from goat origin. *Food Control* 18(4): 375-378.
- Oveisi, M.R., Jannat, B., Sadeghi, N., Hajimahmoodi, M. and Nikzad, A. (2007). Presence of aflatoxin M<sub>1</sub> in milk and infant milk products in Tehran, Iran. *Food Control* 18: 1216-1218.
- Park, D.L. (2002). Effect of processing on Aflatoxin. *Advances in experimental medicine and biology* 504: 173-179.
- Pittet, A. (1998). Natural occurrence of mycotoxins in foods and feeds: An updated review. *Revue de Medicine Veterinaire* 149(6): 479-492.
- Ramdell, H.S. & Eaton, D.L. (1990). Species susceptibility to Aflatoxin B<sub>1</sub> carcinogenesis. *Cancer Research* 50(3): 615-620.
- Rastogi, S., Dwivedi, D.P., Khanna, K.s. and Das, M. (2004). Detection of Aflatoxin M<sub>1</sub> contamination in milk and infant milk products from Indian Markets by ELISA. *Food Control* 15: 287-290.
- Sefidgar, S.A.A., Gholampour Azizi, I., Khosravi, A.R. and Roudbar Mohammadi, S. (2008). Presence of Aflatoxin M<sub>1</sub> in Raw Milk at Cattle Farms in Babol, Iran. *Pakistan Journal of Biological Science* 11(3):484-486.
- Soha, S., Mazloumi, M.T. and Borji, M. (2006). Reduction of Aflatoxin M<sub>1</sub> residue in milk utilizing chemisorption compounds and its effect on quality of milk. *Journal of Arab Neonatology Forum* 3(2).
- Stoloff, L., Van Egmond, H.P. and Parks, D.L. (1991). Rationales for the Establishment of limits and regulations for mycotoxins. *Food Additives and contaminants* 8(2): 213-222.
- Tajkarimi, M., Shojaee Aliabadi, F., Salah Nejad, M., Pursoltani, H., Motallebi, A.A. and Mahdavi, H. (2007). Seasonal study of aflatoxin M<sub>1</sub> contamination in milk in five regions in Iran, *International Journal of Food and Microbiology* 116(3): 346-349.
- Unusan, N. (2006). Occurrence of aflatoxin M<sub>1</sub> in UHT milk in Turkey, *Food and Chemical Toxicology* 44: 1897-1900.
- Williams, J.H., Phillips, T.D., Jolly, P.E., Stiles, J.K., Jolly, C.M. and Aggarwal, D. (2004). Human aflatoxicosis in developing countries: a review of toxicology, exposure, potential health consequences and interventions. *American Journal of Clinical Nutrition* 80(5) :1106-1122.