

Full Article

Virulence assessment of a *Neospora caninum* isolate for inbred C57BL/6 mouse

kargar^{*1}, M., Sayari¹, M., Namavari³, M.M., Lotfi², M., Kamalzadeh², M.

1. Department of Pathobiology, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

2. Razi Vaccine and Serum Research Institute Karaj, Iran

3. Razi Vaccine and Serum Research Institute, Shiraz, Iran

Received 18 May 2011; accepted 01 Oct 2011

ABSTRACT

Neospora caninum (*N. caninum*) is an apicomplexan parasite and causes abortion and congenital neosporosis in cattle worldwide. In this study, we evaluate the virulence of a *N. caninum* isolate on mouse strain C57BL/6. Six groups of five mice C57BL/6 were intraperitoneally inoculated with 1×10^7 , 1.5×10^7 , 2×10^7 , 3×10^7 and 4×10^7 tachyzoites and a control inoculum of DMEM, respectively. Clinical signs and mortality rate were recorded and confirmed by histopathological findings and molecular method. The results of this study indicated that LD50 was 2.5×10^7 tachyzoites of *N. caninum* per C57BL/6 mouse. This can be used as a lethal challenge model in vaccine development studies.

Keywords: *Neospora caninum*, C57BL/6 mouse model, LD50

INTRODUCTION

Neospora caninum (*N. caninum*) is a cyst-forming coccidian parasite closely related to *Toxoplasma gondii* species. It has been shown to be a serious cause of abortion in cattle (Dubey & Schares 2006). Considerable economic losses are attributed to *N. caninum* in the farming industry, including the costs of still birth and neonatal mortality, born clinically normal but with chronic infection, increased calving interval resulting from early fetal death and reduced milk production (Dubey *et al* 1996, Trees *et al* 1999, Dubey 2003, Botelho *et al* 2007). The dairy and beef industries in several parts of the world are severely affected by this disease (Dubey 1999, Wouda 2000, Ramamoorthy

et al 2007). Reproductive losses and abortion due to *N. caninum* infection has been reported in cattles in Iran (Habibi *et al* 2005, Malmasi *et al* 2007, Razmi *et al* 2006 and 2007). Mice have been commonly used as a model in experiments conducted to study the biology of *N. caninum* and develop vaccine against this infection (Lindsay *et al* 1990, 1999, Cannas *et al* 2003, Bartley *et al* 2006). The C57BL/6 mice are very well suited for *N. caninum* studies (Long *et al* 1998, Botelho *et al* 2007, Ramamoorthy *et al* 2007).

LD50 is one of the pathogen characters and a widely accepted tool in vaccine and toxicity studies. It is of particular use to manufacturers as it is prescribed in most international pharmacopeias as a test to measure the efficacy of vaccines and drugs (Rabouhans 1986, Botham 2002, Ramamoorthy *et al* 2005). On other hand pervious study demonstrated that the number of

* Author for correspondence. Email: mkargar16@yahoo.com

passages of *N. caninum* in vitro could affect its virulence (Bartley et al 2006). The present study determined the LD50 of *N. caninum* strain NC-1 isolated at Razi vaccine and serum research institute, for the mouse strain C57BL/6.

MATERIALS AND METHODS

Parasite culture. Vero cells were cultured in DMEM Medium (Sigma Co., USA), containing 2% fetal calf serum, penicillin/ml (10,000 U), streptomycin/ml (100 µg) and amphotericin (25 µg, Invitrogen; USA) with 5% CO₂ in tissue culture, kept at 37 °C in flasks and were trypsinized 1 to 3 times a week. *N. caninum* (strain NC-1) tachyzoites were cultured on the monolayer spread of the Vero cells (Dubey et al 1988, Dubey 1999, Dubey & Schares 2006). Intracellular parasites were harvested by trypsinization of infected Vero cells about 4-5 days post infection, after lysis of the Vero cells (60-80%). Free tachyzoites were removed from the tissue culture flasks by collecting the medium supernatant. For preparation of the necessary inoculum doses (1×10^7 , 1.5×10^7 , 2×10^7 , 3×10^7 and 4×10^7), parasites were pelleted by centrifugation (2,000 g for 10 min) concentration set with a Neubauer chamber and diluted in DMEM medium.

Infection of Mice. Thirty female C57BL/6 mice were divided into 6 groups of five 6–8 week old mice, that each was inoculated intraperitoneally (IP) with 1×10^7 , 1.5×10^7 , 2×10^7 , 3×10^7 and 4×10^7 tachyzoites and a control inoculum of DMEM, respectively (Table 1). Mice were observed and monitored daily for 28 days for clinical signs and mortality.

Histopathologic and PCR examination. At necropsy, 20 mg of the brain tissue was weighed into sterile microtubes and stored frozen at -20 °C until for PCR assay. The rest of the brain material was fixed in 10% buffered formalin and processed for routine paraffin embedding and haematoxylin and eosin (H&E) staining of 5µm thin sections. The used protocol for DNA extraction and PCR assay of brains samples was

as the method previously described by Yamage et al (1996).

Statistical analysis. The LD50 was calculated by the Reed and Muench method (Pizzi 1950).

Table 1. Experimental design.

Group	Number of mice	Intraperitoneally (IP) inoculum
		NC tachyzoite
1	5	1×10^7
2	5	1.5×10^7
3	5	2×10^7
4	5	3×10^7
5	5	4×10^7
6	5	PBS

Table 2. LD50 titration of *N. caninum* tachyzoites

Mouse number	Dose of tachyzoites				
	1×10^7	1.5×10^7	2×10^7	3×10^7	4×10^7
1	K28	K28	K28	D6	D3
2	K28	K28	K28	D6	D3
3	K28	K28	K28	D7	D4
4	K28	K28	K28	D9	D5
5	K28	K28	K28	D9	D5

D: Died; K: killed; number: number of days to death or euthanasia

RESULTS

All the mice in the 2×10^7 group survived and none of the mice in the 3×10^7 group survived. Data from LD50 titration is represented in Table 2. The LD50 as determined by the Reed and Muench method was 2.5×10^7 tachyzoites per mouse. In our study, all the mice administered 3×10^7 and 4×10^7 tachyzoites died between days 3-9 PI., while mice administered 1×10^7 , 1.5×10^7 and 2×10^7 were survived. Most of the mice from groups 2 and 3 showed clinical signs of disease that they were reluctant to move and displayed rough hair coats. Also, at day 9, two mice from group 3 showed other clinical symptoms including hunching and pelvic limb weakness. Further, no clinical signs have been observed in mice infected with 1×10^7 tachyzoites. The control group showed no clinical symptoms or mortality.

The most significant lesions found on histopathological examination of brain tissues revealed mild to moderate lymphoplasmacellular meningoencephalitis, lymphoplasmacellular perivascular cuffing (PVC) (Figure 1) and focal glial cell activation. In the PCR assay brain samples from these infected mice produced a single product of 328 bp similar to those of the positive control while negative control was PCR negative (Figure 2). Histopathological examination of brain samples from mice that died from 3×10^7 and 4×10^7 tachyzoites inoculum did not show any pathological changes in the brain.




Figure 1. Histopathological lesions of brain in a mouse experimentally infected with 2×10^7 tachyzoites. Histopathological lesions of brain in a mouse experimentally infected with 2×10^7 tachyzoites. A: gliosis (arrow), B: perivascular cuffing (PVC, arrow) by monocellular cells (H and E, scale bar: 85 μ m).

DISCUSSION

Determining the LD50 value eliminates the need for a long waiting period for the analysis of tissue samples by histological or PCR methods (Rabouhans 1986, Botham 2002, Ramamoorthy *et al* 2005). In this study LD50 was determined to be 2.5×10^7 tachyzoites per C57BL/6 mouse that shows lower virulence compared with other study which the LD50 was calculated as 1.5×10^7 tachyzoites of NC1 per mouse (Ramamoorthy *et al* 2007). The NC-1 is the most common laboratory strain of *N. caninum*, so this difference between the LD50 are not entirely surprising since the passage number, condition of culture and even the

concentration of antibiotics affects on the virulence of tachyzoite and the possibility of difference in resistance of C57BL/6 mice (Daneshvar *et al* 2003, Bartley *et al* 2006). In this study, the most common clinical signs of disease in mice were related to central nervous system and include reluctant to move, hunching and pelvic limb weakness. On other hand, the main histopathological changes, found on histopathological examination of brain tissues of the animals of 3 group, were focal glial cell activation, lymphoplasmacellular meningoencephalitis and perivascular cuffing (PVC). Similar histological observations to those described in our study were seen by Long *et al* (1998), Lindsay *et al* (1995, 1999), Lundén *et al* (2002), Bartley *et al* (2006) and Ramamoorthy *et al* (2007), when examining CNS tissue from infected mice.




Figure 2. Detection of *N. caninum*-specific DNA in infected mice brains. Agarose gel electrophoresis of *N. caninum*-specific PCR products, amplified from DNA samples collected from brains of group 3 infected mice with 2×10^7 tachyzoites (Lane 2 and 3). Negative (Lane 4; brain from uninfected mice) and positive (Lane 1) control samples and DNA ladder standards (M) were also shown.

Histopathological evaluation of brain tissues from the animals of 3 and 4 group did not show any pathological changes, possibly because the time from inoculation to death was short and death occurred shortly, before brain lesions could develop. Similar findings have been

reported by using PCR to quantify infection intensitie in brain tissue (Eperon *et al* 1999, Gottstein *et al* 2001).

To our knowledge, C57BL/6 mice exhibit clinical signs and mortality due to Neosporosis and are, therefore, an invaluable tool in for determine the LD50 of *N. caninum* in mice.

References

- Bartley, P.M., Wright, S., Sales, J., Chianini, F., Buxton, D. and Innes, E.A. (2006). Long-term passage of tachyzoites in tissue culture can attenuate virulence of *Neospora caninum* in vivo. *Parasitology* 135: 421–432.
- Botham, P.A. (2002). Acute system toxicity. National Research Council, Institute of Laboratory Animal. 43 (Suppl.), S27- S30.
- Botelho, A.S., Teixeira, L., Correia-da-Costa, J.M., Faustino, A.M., Castro, A.G. and Vilanova, M. (2007). *Neospora caninum*: high susceptibility to the parasite in C57BL/10ScCr mice. *Experimental Parasitology* 115(1): 68-75.
- Cannas, A., Naguleswaran, A., Muller, N., Eperon, S., Gottstein, B. and Hemphill, A. (2003). Vaccination of mice against experimental *Neospora caninum* infection using NcSAG1- and NcSRS2-based recombinant antigens and DNA vaccines. *Parasitology* 126: 303-312.
- Daneshvar, H., Coombs, G.H., Hagen, P. and Philips, S. (2003). *Leishmania mexicana* and *Leishmania major* attenuation of wild type parasites and vaccination with the attenuated lines. *Journal of Infectious Diseases* 187: 1662–1668.
- Dubey, J.P. (1999). Neosporosis in cattle: biology and economic impact. *Journal of the American Veterinary Medical Association* 214: 1160–1163.
- Dubey, J.P. (2003). Review of *Neospora caninum* and neosporosis in animals. *Korean Journal of Parasitology* 41: 1–16.
- Dubey, J.P., Hattel, A.L., Lindsay, D.S. and Topper, M.J. (1988). Neonatal *Neospora caninum* infection in dogs: isolation of the causative agent and experimental transmission. *Journal of the American Veterinary Medical Association* 193: 1259–1263.
- Dubey, J.P. and Lindsay, D.S. (1996). A review of *Neospora caninum* and neosporosis. *Veterinary Parasitology* 67: 1-59.
- Dubey J.P. and Schares G. (2006). Diagnosis of bovine neosporosis. *Veterinary Parasitology* 140(1-2): 1-34.
- Eperon, S., Bronnimann, K., Hemphill, A. and Gottstein, B. (1999). Susceptibility of B-cell deficient C57BL/6 (μ MT) mice to *Neospora caninum* infection. *Parasite Immunology* 21: 225–236.
- Gottstein, B., Eperon, S., Dai, W.J., Cannas, A., Hemphill, A. and Greif, G. (2001). Efficacy of toltrazuril and ponazuril against experimental *Neospora caninum* infection in mice. *Parasitology Research* 87: 43–48.
- Habibi, G.R., Hashemi-Fesharki, R., Sadrebazzaz A., Bozorgi, S. and Bordbar, N. (2005). Seminested PCR for Diagnosis of *Neospora caninum* Infection in Cattle. *Archives of Razi Institute* 59: 55-64.
- Haddad, J.P.A., Dohoo, I.R. and VanLeewen, J.A. (2005). A review of *Neospora caninum* in dairy and beef cattle, a Canadian perspective. *The Canadian Veterinary Journal* 46: 230–243.
- Lindsay, D.S., Blagburn, B.L. and Dubey, J.P. (1990). Infection of mice with *Neospora caninum* (Protozoa: Apicomplexa) does not protect against challenge with *Toxoplasma gondii*. *Infection and Immunity* 58: 2699-2700.
- Lindsay, D.S., Lenz, S.D., Blagburn, B.L. and Brake, D.A. (1999). Characterization of temperature-sensitive strains of *Neospora caninum* in mice. *Journal of Parasitology* 85: 64-67.
- Lindsay, D.S., Lenz, S.D., Cole, R.A., Dubey, J.P. and Blagburn, B.L. (1995.) Mouse model for central nervous system *Neospora caninum* infections. *Journal of Parasitology* 81: 313-315.
- Long, M.T., Baszler, T.V. and Mathison, B.A. (1998). Comparison of intracerebral parasite load, lesion development and systemic cytokines in mouse strains infected with *Neospora caninum*. *Journal of Parasitology* 84: 316–320.
- Lunde'n, A., Wright, S., Allen, J.E. and Buxton, D. (2002). Immunisation of mice against neosporosis. *International Journal for Parasitology* 32: 867–876.
- Malmasi, A., Hosseinejad, M., Haddadzadeh, H., Badii, A. and Bahonar, A. (2007). Serologic study of anti-*Neospora caninum* antibodies in household dogs and dogs living in dairy and beef cattle farms in Tehran, Iran. *Parasitology Research* 100(5): 1143-1145.
- Pizzi, M. (1950). Sampling variation of the fifty percent endpoint, determined by the Reed-270 Muench (Behrens) method. *Human Biology* 22: 151-190.
- Rabouhans, M.L. (1986). Reduction of animal usage: British Pharmacopoeia Commission policy. *Developments in Biological Standardization* 64: 11–16.

- Ramamoorthy, S., Duncan, R., Lindsay, D.S. and Sriranganathan, N. (2007). Optimization of the use of C57BL/6 mice as a laboratory animal model for *Neospora caninum* vaccine studies. *Veterinary Parasitology* 145: 253-259.
- Ramamoorthy S., Sriranganathan N. and Lindsay D.S. (2005). Gerbil model of acute neosporosis. *Veterinary Parasitology* 127: 111-114.
- Razmi, G.R., Maleki, M., Farzaneh, N., Talebkhan Garoussi M. and Fallah, A.H. (2007). First report of *Neospora caninum*-associated bovine abortion in Mashhad area, Iran. *Parasitology Research* 100(4): 755-757.
- Razmi, G.R., Mohammadi, G.R., Garrosi, T., Farzaneh, N., Fallah, A.H. and Maleki, M. (2006). Seroepidemiology of *Neospora caninum* infection in dairy cattle herds in Mashhad area, Iran. *Veterinary Parasitology* 135(2): 187-189.
- Trees, A.J., Davidson, H.C., Innes, E.A. and Wastling, J.M. (1999). Towards evaluating the economic impact of bovine neosporosis. *International Journal for Parasitology* 29: 1195-1200.
- Wouda, W. (2000). Diagnosis and epidemiology of bovine neosporosis: a review. *The Veterinary quarterly* 22: 71-74.
- Yamaga, M., Flechtner, O., and Gottstein, B. (1996). *Neospora caninum*: specific oligonucleotide primers for the detection of brain "cyst" DNA of experimentally infected nude mice by the polymerase chain reaction (PCR). *Journal of Parasitology* 82: 272-279.

Archive of SID