



## Reducing nitrogen fertilization application in Cucumber by mycorrhiza colonization of the plant

M. Haghghi\*, B. Abdolahi-pour

Department of Horticulture, College of Agriculture, Isfahan University of Technology, Isfahan I. R. Iran

\* Corresponding Author: mhaghghi@cc.iut.ac.ir  
DOI:10.22099/iar.2019.5212

### ARTICLE INFO

#### Article history:

Received 21 May 2018

Accepted 28 February 2019

Available online 8 June 2018

#### Keywords:

Antioxidant activity  
Cucumber  
Mycorrhiza inoculation  
Nitrogen  
Nitrate reductase activity

**ABSTRACT-** The effect of different nitrogen levels on plant growth, phenol content, antioxidant and Nitrate reductase (NR) activity of cucumber (*Cucumis sativus* cv. Super) inoculated with mycorrhiza was studied. A factorial experiment based on a completely randomized design (CRD) with 6 replicates was designed. Treatments were three levels of nitrogen (NO<sub>3</sub>-50, NO<sub>3</sub>-75 and NO<sub>3</sub>-100) and two mycorrhiza inocula (1000 spore (AM1), 2000 spore (AM2) and no mycorrhiza treatment (AM0) as the control. Results showed that the root fresh weight increased by mycorrhiza inoculation at all nitrogen levels, while it was unaffected by nitrogen levels and mycorrhiza inoculation. FV/FM was higher in NO<sub>3</sub>-75 mycorrhizal inoculated plants compared to that of AM0 ones. Antioxidant activity of plants increased due to mycorrhiza symbiosis in nitrogen deficiency treatments, so that AM1 and AM2 increased antioxidant activity in NO<sub>3</sub>-50 and NO<sub>3</sub>-75 treatments, respectively, as compared to the AM0 plants. The highest NR activity was observed in the NO<sub>3</sub>-50 treatment. However, mycorrhiza inoculation decreased NR activity of the plants at all nitrogen levels. Therefore, it can be concluded that mycorrhiza inoculation, especially the 1000 spores treatment (AM1), and a moderate level of NO<sub>3</sub> (NO<sub>3</sub>-75) can be used and these application levels can be very effective for greenhouse cucumber production.

### INTRODUCTION

Nitrogen (N), as an essential nutrient element, which consists of about 3-4% of the dry matter of plants, often becomes a limiting factor for plant growth (Makhziah et al., 2013). Proteins increase vegetative and reproductive growth of plants. The production of proteins, in turn, depends on the supply of nitrogen (Lawlor, 2008). Nitrogen is also important for plant growth because it is one of the basic elements for the synthesis of chlorophyll and several enzymes and it is involved in cell division and growth (Liu et al., 2013).

Since N plays an important role in crop yield and quality (Chen et al., 2004), its application in agricultural practices is continuously increasing. Nitrate is the major source of N for higher plants. Uptake of nitrate by plants results in high nitrate accumulation in plants, especially in vegetable crops (Chen et al., 2004). On the other hand, application of nitrogen to produce more yields has become increasingly a potential pollutant (Zand-Parsa et al., 2006). Application of nitrogen fertilizers is more than what the plant needs and the nitrogen use efficiency is less than 10% (Dai et al., 2011); therefore, large amounts of unused N in soils lead to many environmental problems (Gollany et al., 2004).

Azcon et al. (1992) reported that mycorrhizal fungi are important in ecological agriculture as biofertilizers, biological protectors, and biological control agents. The positive effect of mycorrhiza on plant growth, especially

under nutrient limited conditions, has been established by some researchers (Mortimer et al., 2009; Goss and de Varennes, 2002). Smith and Read (1997) demonstrated that *Arbuscular mycorrhiza* hyphae spread in the soil around the root and provided the surface area, and caused absorption of nutritional elements such as phosphorus, nitrogen, zinc, and copper for the host plant. Symbiosis relation between the mycorrhizal fungi and their host plants can also improve transferring large amounts of nitrogen in different plants (He et al., 2007; Mortimer et al., 2009; Li et al., 2006). Also, it has been reported that mycorrhiza fungi increased phosphorous content of plants and positively affected nitrogen uptake by plants (Tufenkci et al., 2005).

The two most important nitrogen sources for plant and mycorrhiza are nitrate and ammonium (Masoumeh et al., 2009). The colonization and growth of mycorrhiza is correlated with the nitrogen status in the soil and it is inhibited by high nitrogen levels (Azcon et al., 1992). The transfer and metabolism of nitrogen is dependent on the biosynthesis and flux of carbon by host plant to mycorrhizal fungi (Liu et al., 2013).

N deficiency and excession is a nutritional stress on plant which activates defense plant system, such as antioxidant and non-antioxidant activities (Chen et al. 2013). On the other hand, it has been shown that mycorrhiza inoculation increased total phenol content leaves and flowers in artichoke plants (Ceccarelli et al.,

2010). Total phenol content of grape significantly increased by all strains of mycorrhiza inoculation compared to the control plants (Eftekhari et al., 2012).

The objective of the present research was to study the effect of i) reduction of nitrogen rates and ii) beneficial effects of mycorrhiza inoculation to compensate the reduction of nitrogen rates on plant growth, antioxidant amount, and nitrate reductase activity of cucumber (*Cucumis sativus* cv. Super N3) under greenhouse condition.

## MATERIALS AND METHODS

This experiment was conducted in a plastic greenhouse in the Department of Horticulture Science of Isfahan University of Technology, Isfahan, Iran. The temperature in greenhouse was 30-35 °C and relative humidity was 30-35% between July and October 2013. The study was carried out as a factorial experiment based on CRD with 6 replications. The inoculation dosages of mycorrhiza were 1000 spore (AM1; 50 g), 2000 spore (AM2; 100 g) and non-mycorrhizal inoculated plants (AM0) as controls. Nitrogen treatments were 52.5 (50% of Johnson nutrient solution= NO<sub>3</sub>-50), 78.75 (75% of Johnson nutrient solution= NO<sub>3</sub>-75) and 105 (100% of Johnson nutrient solution= NO<sub>3</sub>-100) mg L<sup>-1</sup> of Johnson nutrient solution (Jones, 1930). The components of Johnson nutrient solution are introduced in Table 1.

Seeds of cucumber (*Cucumis sativus* cv. Super N3) were germinated in peat at 25 °C for 30 days. The seedlings with uniform sizes were transplanted to 3-L pots containing sand which had received mycorrhiza inoculum of *Glomus mosseae* according to mycorrhizal treatments from Touran Biotech Company (Shahroud, Iran). The interaction of mycorrhizal and root was observed with light microscopic through examination of all roots, stained with Chlorazol Black E at the root surface. The pots were irrigated daily with about 50 ml Johnson nutrient solution with defined treatments.

Chlorophyll content was measured using chlorophyll meter (SPAD-502, Minolta Corp., Ramsey, NJ, USA) and FV/FM was measured by chlorophyll fluorescence (OS-30, USA) after 3 weeks. Nitrate reductase (NR) activity was measured according to Sagi et al. (1997) after one month.

In order to determine total phenol content in 1 g of the leaves, McDonald et al.'s (2001) procedure was followed. Briefly, each leaf sample was mixed with 5 ml folin ciocaltea and 4 ml aqueous Na<sub>2</sub>CO<sub>3</sub> separately. The phenol concentration was determined at 765 nm as gallic acid equivalents per gram (mg GAE g<sup>-1</sup> DW) by a spectrophotometer.

Antioxidant activity of cucumber leaves was determined following Koleva et al.'s (2002) procedure. Briefly, 3 mg of samples were dissolved in 5 ml methanol stock and 1.4 ml of this solution was blended with 0.6 ml of DPPH solution. After 30 min, the absorbance of the solution was recorded at 515 nm by a spectrophotometer (V-530, JASCO, Japan) against methanol as a blank. The 0.2 mM of DPPH solution in

methanol was used as a stock of DPPH for determination of the free radical scavenging activity of the samples.

The antiradical activity was calculated by the following equation:

$$\text{Antioxidant activity} = \frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \times 100$$

A<sub>control</sub>: absorbance of the DPPH solution

A<sub>sample</sub>: absorbance of the DPPH solution after the addition of the sample

At the end of the experiment when plants started anthesis (30 days after transplanting), plants were harvested, fresh weights of roots and shoots were measured and then, all the samples were oven dried at 70°C for 3 days and the dry weights were measured by an analytical balance (to 0.001 decimal places).

Mycorrhizal dependency for shoot (MDS) and root (MDR), which shows the dependency of yield to mycorrhiza inoculation (M1 and M2) compared with the non-inoculated treatment, was also calculated using the following equation (Planchette et al., 1983), which shows the dependency of yield to mycorrhiza inoculation (M1 and M2) compared with the non-inoculation media:

$$\text{MDS} = \frac{\text{FWS}_{\text{mycorrhiza}} - \text{FWS}_{\text{non-mycorrhiza}}}{\text{FWS}_{\text{mycorrhiza}}}$$

FWS= Fresh weight of shoot

$$\text{MDR} = \frac{\text{FWR}_{\text{mycorrhiza}} - \text{FWR}_{\text{non-mycorrhiza}}}{\text{FWR}_{\text{mycorrhiza}}}$$

FWR= Fresh weight of Root

Nitrogen dependency for shoot (NDS) and root (NDR) was calculated by the following modified formula:

$$\text{NDS} = \frac{\text{FWS}_{\text{full nitrogen}} - \text{FWS}_{\text{nitrogen deficit}}}{\text{FWS}_{\text{full nitrogen}}}$$

FWS= Fresh weight of shoot

$$\text{NDR} = \frac{\text{FWR}_{\text{full nitrogen}} - \text{FWR}_{\text{nitrogen deficit}}}{\text{FWR}_{\text{full nitrogen}}}$$

FWR= Fresh weight of Root

Shoot nutrient efficiency (SNE) was calculated following Sepeher et al.'s (2012) procedure, using the following equation

$$\text{SNE} = (\text{SDW}_{\text{nitrogen deficit}} / \text{SDW}_{\text{full nitrogen}}) \times 100$$

SDW=shoot dry weight

$$\text{RNE} = (\text{RDW}_{\text{nitrogen deficit}} / \text{RDW}_{\text{full nitrogen}}) \times 100$$

RDW=Root dry weight

Stress tolerance indexes were calculated using the following modified equation (Farshadfar et al., 2013):

$$\text{Stress Tolerance; TOL} = \text{FWS}_{\text{full nitrogen}} - \text{FWS}_{\text{deficit nitrogen}}$$

FWS= Fresh weight of shoot

The present study was arranged in a factorial design with 6 replications. Data were analyzed using Statistix 8 (Tallahassee FL, USA). All data were subjected to two-way ANOVA and the means were compared for significance by the least significant difference (LSD) test at  $P < 0.05$ .

## RESULTS AND DISCUSSION

Shoot and root fresh weights, shoot dry weight, chlorophyll content, and antioxidant activity increased with both AM1 and AM2 applications compared to non-mycorrhizal treatment (AM0). Root dry weight did not change significantly with AM application. FV/FM decreased with AM1 and AM2 applications compared to AM0. Total phenol content increased in AM1 and AM2 compared to AM0 and was the highest in AM2 treatment. Conversely, NR activity decreased in AM1 and AM2 compared to AM0 and was the lowest in AM2 treatment (Table 2).

By increasing NO<sub>3</sub>-50 level to NO<sub>3</sub>-75 and NO<sub>3</sub>-100, shoot and root fresh weights, shoot dry weight, and chlorophyll content increased significantly. However, root dry weight and FV/FM did not change significantly.

Antioxidant activity and NR activity increased in both NO<sub>3</sub>-75 NO<sub>3</sub>-100 treatments compared to NO<sub>3</sub>-50 treatment and was the highest in NO<sub>3</sub>-100 treatment. Total phenol content decreased by increasing NO<sub>3</sub> to NO<sub>3</sub>-100 level compared to NO<sub>3</sub>-50 to NO<sub>3</sub>-75 levels (Table 2).

Mycorrhizal dependency of root was higher in AM1 than AM2 treatment, and the same results were observed for MDS. Nitrogen dependency for shoot and root was higher in NO<sub>3</sub>-50 treatment than NO<sub>3</sub>-75 treatment. Shoot nutrient efficiency was higher in NO<sub>3</sub>-75 treatment than NO<sub>3</sub>-50 treatment. Conversely, Stress Tolerance to N deficiency was lower in NO<sub>3</sub>-75 treatment than NO<sub>3</sub>-50 treatment. Root nutrient efficiency was not affected by NO<sub>3</sub> content (Table 4).

**Table 1.** The components of Johnson nutrient solution

Compound	M.W	Stock conc. M	Stock conc. gL <sup>-1</sup>	ml stock per liter final sol.	Element	Final conc. ppm	50% ppm	75% ppm
KNO3	101.10	1.0	101.10	6.0	N	224	112	168
Ca(NO3)2.4H2O	236.16	1.0	236.16	4.0	K	235	117.5	176.25
NH4H2PO4	115.08	1.0	115.08	2.0	Ca	160	80	120
MgSO4.7H2O	246.49	1.0	246.49	1.0	P	62	62	62
					S	32	32	32
					mg	24	24	24
KCl	74.55	50	3.72	1.0	Cl	1.77	1.77	1.77
H3BO3	61.84	25	1.54	1.0	B	0.27	0.27	0.27
MnSO4.H2O	169.01	2.0	0.33	1.0	Mn	0.11	0.11	0.11
ZnSO4.7H2O	287.55	2.0	0.57	1.0	Zn	0.13	0.13	0.13
CuSO4.5H2O	249.71	0.5	0.12	1.0	Cu	0.03	0.03	0.03
H2MoO4	161.97	0.5	0.081	1.0	Mo	0.05	0.05	0.05
Fe-EDTA	346.08	50	21.53	1.0	Fe	2.80	2.80	2.80

Changes in potassium and calcium elements were adjusted by chloride salts.

**Table 2.** Main effect of *Arbuscular mycorrhiza* (AM) on some characteristics of cucumber

<i>Arbuscular mycorrhiza</i> treatment	Shoot fresh weight/plant (g)	Root fresh weight/plant (g)	Shoot dry weight/plant (g)	Root dry weight/plant (g)	Chlorophyll content (SPAD value)	FV/FM	Antioxidant activity (% inhibition)	Total phenol content (mg GAE g <sup>-1</sup> FW)	NR activity (mg g <sup>-1</sup> FW)
AM0	144.1 b	19.2 b	11.0 b	16.7 a	7.05 b	15.08 a	0.15 b	0.22 c	0.07 a
AM1	160.4 a	34.1 a	14.3 a	15.7 a	13.53 a	0.03 b	0.48 a	87.31 b	0.06 b
AM2	158.0 a	31.3a	14.0 a	15.9 a	12.04 a	0.04 b	0.44 a	89.13 a	0.05 c

Means with different letters in each column are significantly different at  $P < 0.05$  according to the LSD test.

**Table 3.** Main effect of different NO<sub>3</sub> concentration on some characteristics of cucumber.

Nitrogen level	Shoot fresh weight/plant (g)	Root fresh weight/plant (g)	Shoot dry weight/plant (g)	Root dry weight/plant (g)	Chlorophyll content (SPAD value)	FV/FM	Antioxidant activity % inhibition	Total phenol content (mg GAE g <sup>-1</sup> FW)	NR activity (mg g <sup>-1</sup> FW)
NO <sub>3</sub> -50	136.2 b	6.1 b	10.8 b	12.25 a	7.72 b	4.83 a	0.26 c	61.51 b	0.06 c
NO <sub>3</sub> -75	158.2 a	8.8 a	13.1 a	12.40 a	10.88 a	5.98 a	0.46 b	65.50 a	0.07 b
NO <sub>3</sub> -100	158.1 a	8.2 a	12.9 a	12.18 a	9.02 ab	4.34 a	0.75 a	49.65 c	0.09 a




Means with different letters in each column are significantly different at  $P < 0.05$  according to the LSD test.

**Table 4.** The effect of mycorrhiza/NO<sub>3</sub> levels on MD, ND, NE and TOL of root and shoot

Arbuscular ycorrhiza treatment	MDR	MDS	Nitrogen level	NDS	NDR	SNE	RNE	TOL
AM1	0.14	0.10	NO <sub>3</sub> -50	0.13	0.25	83.72	100.57	21.9
AM2	0.06	0.08	NO <sub>3</sub> -75	-0.0006	-0.07	101.55	101.80	-0.1
T-test	0.01	0.41		0.027	0.02	0.02	0.27	0.002

Mycorrhizal dependency for shoot (MDS) and root (MDR); Nitrogen dependency for shoot (NDS) and root (NDR); Shoot nutrient efficiency (SNE); Root nutrient efficiency (RNE); Stress Tolerance (TOL)


Fresh weight of shoots increased significantly in NO<sub>3</sub>-75×AM1 treatment but it did not show any significant difference between other treatments (Fig 1a). Root fresh weight increased significantly by AM1 and AM2 applications compared to non-mycorrhizal treatment (AM0) in all levels of NO<sub>3</sub> and the most root fresh weight was observed in NO<sub>3</sub>-75×AM1 treatment (Fig 1b). Dry weight of shoots increased significantly in NO<sub>3</sub>-75 and NO<sub>3</sub>-100 treatments compared to NO<sub>3</sub>-50 treatment in all mycorrhiza levels and the lowest dry weight of shoots was found in NO<sub>3</sub>-50 treatment with all mycorrhiza levels (Fig 1c). Root dry weight increased by increasing NO<sub>3</sub> from NO<sub>3</sub>-50 to NO<sub>3</sub>-75 and NO<sub>3</sub>-100 in all mycorrhiza levels. However, in NO<sub>3</sub>-50 treatment, by applying mycorrhiza (AM1 and AM2), root dry weight significantly increased compared to the AM0 treatment (Fig 1d).




**Fig. 1.** Interactive effect of mycorrhiza inoculation and nitrogen concentration on shoot fresh weight (a), root fresh weight (b), shoot dry weight (c) and root dry weight (d) per plant. Bars with different letters across treatments are significantly different at  $P < 0.05$  according to the LSD test.

Mycorrhiza application in NO<sub>3</sub>-50 treatment increased SPAD values significantly when AM2 application was used. The highest SPAD value was observed in NO<sub>3</sub>-100×AM1 treatment. However, no significant differences were found between NO<sub>3</sub>-100×AM0 and NO<sub>3</sub>-100×AM2 treatments and between NO<sub>3</sub>-75 treatments in none of three levels of mycorrhiza applications (Fig 2).

No significant differences were observed in FV/FM between the different treatments. However, the highest and lowest levels of FV/FM were found in NO<sub>3</sub>-100×AM0 treatment and NO<sub>3</sub>-50×AM1 treatment, respectively (Fig 3).




**Fig. 2.** Interactive effect of mycorrhiza inoculation and nitrogen concentration on SPAD value. Bars with different letters across treatments are significantly different at  $P < 0.05$  according to the LSD test.




**Fig. 3.** Interactive effect of mycorrhiza inoculation and nitrogen concentration on FV/FM. Bars with different letters across treatments are significantly different at  $P < 0.05$  according to the LSD test.

Phenol content of cucumber leaves increased by AM1 and AM2 applications compared to AM0 treatment in all levels of NO<sub>3</sub>. The highest phenol content was observed in NO<sub>3</sub>-75×AM1 treatment (Fig 4).

Antioxidant activity decreased in NO<sub>3</sub>-75 and NO<sub>3</sub>-100 treatments, especially with AM2 application and the highest antioxidant activity was found in NO<sub>3</sub>-50 treatments with all three levels of AM applications (Fig 5).




**Fig. 4.** Interactive effect of mycorrhiza inoculation and nitrogen concentration on total phenol content. Bars with different letters across treatments are significantly different at  $P < 0.05$  according to the LSD test.



**Fig. 5.** Interactive effect of mycorrhiza inoculation and nitrogen concentration on antioxidant activity. Bars with different letters across treatments are significantly different at  $P < 0.05$  according to the LSD test.

NR activity of the leaves increased in NO<sub>3</sub>-100 treatment in all levels of AM applications. However, there were not significant differences between these treatments and NO<sub>3</sub>-50/AM2 and NO<sub>3</sub>-75/AM2 treatments. NR activity was the lowest in NO<sub>3</sub>-50/AM0, NO<sub>3</sub>-50/AM1 and NO<sub>3</sub>-75/AM0 treatments (Fig 6).



**Fig. 6.** Interactive effect of mycorrhiza inoculation and nitrogen concentration on NR activity. Bars with different letters across treatments are significantly different at  $P < 0.05$  according to the LSD test.

Nitrogen fertilization at moderate concentration level increased the plant growth of some leafy vegetable crops, while high nitrogen level reduced plant growth by increasing N supply (Chen et al., 2004). In the present study, by increasing NO<sub>3</sub> level from NO<sub>3</sub>-50 to NO<sub>3</sub>-75 and NO<sub>3</sub>-100, shoot and root fresh weights and shoot dry weight of cucumber significantly increased. Also, shoot fresh weight and root dry weight increased by mycorrhiza inoculation in NO<sub>3</sub>-50 treatment. Similar results were reported by Vazquez et al. (2002) who found that shoot and root weights of alfalfa increased by mycorrhiza colonization under the lowest nitrogen concentration. However, Hays et al. (1982) reported that plant growth of blue grama was enhanced by increasing nutrient solution (N and P), no mycorrhizal infection occurred at high N and P levels and mycorrhiza inoculation caused a lower shoot and root weight at high N level (Hays et al., 1982). The amount and timing of N application can also alter plant morphology, nutrient availability, and net photosynthesis (Zapata & Zaharah, 2002). It seems that by increasing nutrient absorption, the photosynthesis traits improved and caused more assimilate production (Zapata & Zaharah, 2002). Nitrogen and phosphorous are critical determinants of plant growth and productivity, and plant growth and root morphology are important parameters for evaluating the effects of supplied nutrients (Zapata & Zaharah, 2002). N is required in the largest quantities, and its availability and internal concentration affect the partitioning of biomass between roots and shoots (Zhao et al., 2008). Although high N availability affects root and shoot biomass production (Barraclough et al., 1989), P plays an important role in lateral root morphology and

root branching (López-Bucio et al., 2003) and influences not only root development, but also the availability of nutrients (Jin et al., 2005).

In the present experiment, MDS did not show any significant effect on shoot fresh weight, which shows that shoot fresh weight did not depend on mycorrhiza application (Table 3), except in  $\text{NO}_3\text{-}75 \times \text{AM1}$  treatment and AM2 application (Fig 1a). Also, in the medium N deficiency, mycorrhiza improved the root fresh weight effectively in  $\text{NO}_3\text{-}75 \times \text{AM1}$  treatment (Fig 1b). These results supported the data of MDR which showed that generally AM1 application was more effective in increasing root fresh weight (Table 3), especially in the  $\text{NO}_3\text{-}75$  treatment. In agreement with the present study, the result of another study showed that plant height of chickpea and shoot fresh and dry weights of this plant significantly increased by increasing N concentration and mycorrhiza inoculation (Tufenkci et al., 2005). Johansen et al. (1994) showed that dry weights of shoots and roots increased by increasing N concentration, and mycorrhiza colonization (*G. intraradices*) increased cucumber cv. Aminex weight at all nitrogen levels (100, 200, and 400 mg N per kg soil). In the present study, it seems that in the medium and the optimum N levels of the nutrient solution, mycorrhiza had the same effect on root fresh weight and in the high rate of N application, the mycorrhiza application was more effective in increasing root dry weight than shoot dry weight of cucumber (Fig 1c and 1d).

The SPAD value indirectly quantifies chlorophyll content and plant nitrogen status (Wu et al., 2007). In the present study, higher SPAD value was observed in the mycorrhizal inoculated plants compared to the non-mycorrhizal inoculated ones at all nitrogen levels. Huang et al. (2011) stated that high chlorophyll content in the leaves of the mycorrhizal inoculated plants confirms that symbiosis plays a key role in modifying photosynthesis and metabolic activity. The SPAD values were higher in the mycorrhizal inoculated plants than in the non-mycorrhizal inoculated ones in the study of Busquet et al. (2010). Similar results were observed by Campanelli et al. (2013) who found that SPAD value of alfalfa increased by mycorrhiza inoculation under non-stress conditions, but was not influenced under salinity stress. Chlorophyll and carotenoid synthesis are dependent upon mineral nutrients. N is the most important elemental factor in chlorophyll biosynthesis. It might be due to the optimum availability of N, which plays a vital role in cell division and the formation of active photosynthetic pigments including chlorophyll. Green pigments in leaves also depend on P concentration (Daughtry et al., 2000).

Zhu et al. (2011) reported that Fv/Fm was stimulated by mycorrhiza inoculation under non-stress conditions and temperature stress condition. In the present study, mycorrhiza inoculation increased Fv/Fm in  $\text{NO}_3\text{-}75$  treatment, while mycorrhiza symbiosis reduced Fv/Fm in  $\text{NO}_3\text{-}100$ .

In the present study, nitrogen at a moderate level ( $\text{NO}_3\text{-}75$ ) increased total phenol content of cucumber. Giorgi et al. (2009) showed that by reducing N supplementation, total phenol content of yarrow increased. In our study, mycorrhiza inoculation with

2000 spore (AM2) significantly increased total phenol content of cucumber leaves. In agreement with the present study, the increase in total phenol content by mycorrhiza inoculation was also observed by Kapoor (2008), Al-Askar and Rashad (2010), and Mathur and Vyas (1996). Chen et al. (2013) reported that polyphenol content of cucumber increased by mycorrhiza inoculation under non-stress and low temperature stress conditions. Similar results were observed in the present experiment as both mycorrhiza inoculations increased total phenol content of cucumber under N deficit and normal N supply. The highest total phenol content was observed in the N deficit ( $\text{NO}_3\text{-}75$ ) treatment by mycorrhiza inoculation with 1000 spores.

The highest deleterious effect of stress on cucumber was found in the  $\text{NO}_3\text{-}50$  treatment, and Mycorrhiza application did not significantly improve it, because it caused the highest antioxidant activity to moderate the stress condition, but in  $\text{NO}_3\text{-}75$  and  $\text{NO}_3\text{-}100$  treatments, mycorrhiza application effectively decreased stresses on plants, due to not stimulating antioxidant activities. Giorgi et al. (2009) showed that nitrogen deficit reduced plant growth, total nitrogen, chlorophyll, and carotenoids contents. Nitrogen significantly increased total phenol compound and antioxidant contents of root and leaves of yarrow (*Achillea collina* Becker ex Rchb.) compared to the normal nitrogen supply condition (Giorgi et al., 2009). In agreement with the present study, increasing N supply significantly reduced antioxidant activity of chrysanthemum flower (Liu et al., 2010). Liu et al. (2010) reported that an excess N supply negatively affected the antioxidant activity and, thereby, reduced the quality of chrysanthemum. In agreement with the present study, Ordookhani and Zare (2011) reported that antioxidant activity of tomato increased by mycorrhiza inoculation as compared to non-mycorrhiza inoculated plants. Similar results were observed by Akhtar Maya and Matsubara (2013) who reported that mycorrhiza inoculation increased antioxidant activity of leaf, root, and tuber of cyclamen.

Nitrate reductase (NR) activity of cabbage, spinach and rape significantly increased with high N supplementation (Chen et al., 2004). Similarly, an increase in NR activity by mycorrhiza inoculation was observed in onion under water stress by Azcon and Tobar (1998) and in rice under low temperature stress by Liu et al. (2013). Caravaca et al. (2005) showed that drought stress caused a reduction in NR activity and mycorrhiza inoculation improved NR activity of the plants under drought stress. In the present study, it seems that by increasing N level of nutrient solution and  $\text{NO}_3$  availability for plants, the NR activity increased to convert N to an usable form for improving plant metabolism. The application of mycorrhiza in high level (AM2) along with low  $\text{NO}_3$  levels ( $\text{NO}_3\text{-}50$  and  $\text{NO}_3\text{-}75$ ) effectively increased the absorption of  $\text{NO}_3$  which resulted in the same NR activity as was observed in the high level of  $\text{NO}_3$  application; i.e.,  $\text{NO}_3\text{-}100$  (Fig 6). Evidence is on the rise that the AM symbiosis with host plants changes the nutrient uptakes of the plants (Li et al., 2006). Whether and how the plant uptake pathway for N is affected by the colonization with AM fungi is

currently unknown, but some pieces of evidence were presented that showed some plant N transporters were downregulated in colonized roots (Kobae et al., 2010). It was also shown that fungal N uptake system have a high affinity for the uptake of N from the soil (Pérez-Tienda et al., 2012), but how the expression of these transporters is regulated is largely unknown. The transcript levels of some of these transporters are substrate inducible, and regulated by the  $\text{NH}_4^+$  supply and/or fungal  $\text{NH}_4^+$  status (Lopez-Pedrosa et al., 2006). Other researches showed different results on the efficiency of mycorrhiza on N absorption in different plants (Mensah et al., 2015). Evidence was presented that there is a high intraspecific diversity in the efficiency with which fungi are able to improve the N nutrition of their host (Mensah et al., 2015).

## CONCLUSIONS

In general, high nitrogen stress which resulted from low N supply and was observed in the  $\text{NO}_3$ -50 treatment had a deleterious effect on plant growth (decreasing root fresh and dry weights compared to high N supply treatments). Both mycorrhiza inoculation levels enhanced root fresh and dry weights while these parameters were unaffected by mycorrhiza inoculation

in nitrogen stress conditions. The lowest total phenol content of tested plants was found in the  $\text{NO}_3$ -50 treatment. However, both mycorrhiza inoculation levels stimulated total phenol content of tested plants. Mycorrhiza inoculation with 2000 spores (AM2) was more effective than AM1 treatment. The highest NR activity was observed in the  $\text{NO}_3$ -100 treatment in all mycorrhiza inoculation levels while the lowest NR activity was achieved in the  $\text{NO}_3$ -50 and  $\text{NO}_3$ -75 treatments. Mycorrhiza inoculation increased NR activity. Based on the results of this study, it can be concluded that the low nitrogen supply caused high stress on cucumber production during the vegetative growth stage. Mycorrhiza inoculation can compensate for some parameters like root fresh and dry weights, phenol content and NR activity under high nitrogen deficit stress condition ( $\text{NO}_3$ -50), especially under high mycorrhiza inoculation condition (AM2). It seems that mycorrhiza inoculation could compensate for moderate N deficiency ( $\text{NO}_3$ -75) in most parameters and had the same results as  $\text{NO}_3$ -100 provided. Therefore, it can be concluded that by mycorrhiza inoculation, especially the 1000 spores treatment (AM1), a moderate level of  $\text{NO}_3$  ( $\text{NO}_3$ -75) can be used and these application levels can be very effective for greenhouse cucumber production.

## REFERENCES

- Al-Askar, A. A., & Rashad, Y. M. (2010). Arbuscular mycorrhizal fungi: a biocontrol agent against common Bean fusarium root rot disease. *Plant Pathology*, 9 (1), 31–38.
- Azcon, R., & Maria Tobar, R. (1998). Activity of nitrate reductase and glutamine synthetase in shoot and root of mycorrhizal *Allium cepa* effect of drought stress. *Plant Science*, 133, 1–8.
- Aktar Maya, M., & Matsubara, Y. (2013). Influence of arbuscular mycorrhiza on the growth and antioxidative activity in cyclamen under heat stress. *Mycorrhiza*, 23, 381–390.
- Azcon, R., Gomez, M., & Tobar, R. (1992). Effects of nitrogen source on growth, nutrition photosynthetic rate and nitrogen metabolism of mycorrhizal and phosphorous fertilized plants of (*Lactuca sativa* L.). *New Phytologist*, 121, 227–234.
- Barraclough, P. B., Kuhlmann, H., & Weir, A. H. (1989). The effects of prolonged drought and nitrogen fertilizer on root and shoot growth and water uptake by winter-wheat. *Journal of Agronomy and Crop Science*, 163(5), 352–360.
- Busquets, M., Calvet, C., Camprubi, A., & Estaun, V. (2010). Differential effects of two species of arbuscular mycorrhiza on the growth and water relations of (*Spartium junceum*) and (*Anthyllis cytisoides*). *Symbiosis*, 52, 95–101.
- Campanelli, A., Ruta, C., Mastro, G. D., & Morone-Fortunato, I. (2013). The role of arbuscular mycorrhizal fungi in alleviating salt stress in (*Medicago sativa* L.) var. icon. *Symbiosis*, 59 (2), 65–76.
- Caravaca, F., Alguacil, M. M., Hernandez, J. A., & Roldan, A. (2005). Involvement of antioxidant enzyme and nitrate reductase activities during water stress and recovery of mycorrhizal (*Myrtus communis*) and (*Phillyrea angustifolia*) Plants. *Plant Science*, 169, 191–197.
- Ceccarelli, N., Curadi, M., Martelloni, L., Sbrana, C., Picciarelli, P., & Giovannetti, M. (2010). Mycorrhizal colonization impacts on phenolic content and antioxidant properties of artichoke leaves and flower heads two years after field transplant. *Plant and Soil*, 335, 311–323.
- Chen, B. M., Wang, Z. H., Li, S. X., Wang, G. X., Song, H. X., & Wang, X. N. (2004). Effects of nitrate supply on plant growth, nitrate accumulation, metabolic nitrate concentration and nitrate reductase activity in three leafy vegetables. *Plant Science*, 16, 635–643.
- Chen, Sh., Jina, W., Liu, A., Zhang, Sh., Liu, D., Wang, F., Lin, X., & He, C. (2013). Arbuscular mycorrhizal fungi (AMF) increase growth and secondary metabolism in cucumber subjected to low temperature stress. *Scientia Horticulturae*, 160, 222–229.
- Dai, J., Liua, Sh., Zhanga, W., Xua, R., Luoa, W., Zhang, S. H., Yin, Z., Han, L., & Chen, W. (2011). Quantifying the effects of nitrogen on fruit growth and yield of cucumber crop in greenhouses. *Scientia Horticulturae*, 130, 551–561.
- Daughtry, C. S. T., Walthall, C. L., Kim, M. S., Brown, D. E., Colstounm E., & McMurtrey, J. E. (2000). Estimating corn leaf chlorophyll concentration from leaf and canopy reflectance. *Remote Sensing of Environment*, 74(2), 229–239.
- Eftekhari, M., Alizadeh, M., & Ebrahimi, P. (2012). Evaluation of the total phenolics and quercetin content of foliage in mycorrhizal grape (*Vitis vinifera* L.) varieties and effect of postharvest drying on quercetin yield. *Industrial Crops and Products*, 38, 160–165.
- Farshadfar, E., Poursiahbidi, M. M., & Safavi, S. (2013). Assessment of drought tolerance in land races of bread wheat based on resistance/ tolerance indices. *International Journal of Advanced Biological and Biomedical Research*, 12, 143–158.

- Giorgi, A., Mingozzi, M., Madeo, M., Speranza, G., & Cocucci, M. (2009). Effect of nitrogen starvation on the phenolic metabolism and antioxidant properties of yarrow (*Achillea collina* Becker ex Rechb.). *Food Chemistry*, 114, 204–211.
- Gollany, H. T., Molina, J. E., Clapp, C. E., Allmaras, R. R., Layese, M.F., Baker, J.M., & Cheng, H.H. (2004). Nitrogen leaching and denitrification in continuous corn as related to residue management and nitrogen fertilization. *The Journal of Environmental Management*, 33, S289–S298.
- Goss, M. J., & de Varennes, A. (2002). Soil disturbance reduces the efficacy of mycorrhizal associations for early soybean growth and N<sub>2</sub> fixation. *Soil Biology & Biochemistry*, 34, 1167–1173.
- Hays, R., Reid, C. P. P., St, John, T. V., & Coleman, D. C. (1982). Effects of Nitrogen and Phosphorus on Blue Grama Growth and Mycorrhizal Infection. *Oecologia*, 54, 260–265.
- He, Z., He, C., Zhang, Z., Zou, Z., & Wang, H. (2007). Changes of antioxidative enzymes and cell membrane osmosis in tomato colonized by arbuscular mycorrhizae under NaCl stress. *Colloids and Surfaces B: Biointerfaces*, 59, 128–133.
- Huang, Z., Zou, Z., He, C., He, Z., Zhang, Z., & Li, J. (2011). Physiological and photosynthetic responses of melon (*Cucumis melo* L.) seedlings to three *Glomus* species under water deficit. *Plant and Soil*, 339, 391–399.
- Jin, J., Wang, G. H., Liu, X., Pan, X., & Herbert, S.J. (2005). Phosphorus application affects the soybean root response to water deficit at the initial flowering and full pod stages. *Soil Science and Plant Nutrition*, 51(7), 953–960.
- Johansen, A., Jakobsen, I., Jensen, E., & Hyphal, N. (1994). transport by a vesicular-arbuscular mycorrhizal fungus associated with cucumber grown at three nitrogen levels. *Plant and Soil*, 160, 1–9.
- Jones, J. B. (1930). *Hydroponics: a practical guide for the soilless grower*. CRC Press, USA.
- Kapoor, R. (2008). Induced resistance in mycorrhizal tomato is correlated to concentration of jasmonic acid. *The Journal of Bioscience*, 8 (3), 49–56.
- Kobae, Y., Tamura, Y., Takai, S., Banba, M., & Hata, S. (2010). Localized expression of arbuscular mycorrhiza-inducible ammonium transporters in soybean. *Plant Cell Physiology*, 51, 1411–1415.
- Koleva, I. I., Van Beek, T. A., Linssen, J. P. H., de Groot, A., & Evstatieva, L. N. (2002). Screening of plant extracts for antioxidant activity: A comparative study on three testing methods. *Phytochemical Analysis*, 13, 8–17.
- Lawlor, D.W. (2008). Carbon & nitrogen assimilation in relation to yield: mechanisms are the key to understanding production systems. *Journal of Experimental Botany*, 53, 773–787.
- Li, H. Y., Smith, S. E., Holloway, R. E., Zhu, Y. G., Smith, F.A. (2006). Arbuscular mycorrhizal fungi contribute to phosphorus uptake by wheat grown in a phosphorus-fixing soil even in the absence of positive growth responses. *New Phytologist*, 172, 536–543.
- Liu, D., Liu, W., Zhu, D., Geng, M., Zhou, W., & Yang, T. (2010). Nitrogen effects on total flavonoids, chlorogenic acid, and antioxidant activity of the medicinal plant *Chrysanthemum morifolium*. *Journal of Soil Science and Plant Nutrition*, 173 (2), 268–274.
- Liu, Z. L., Li, Y. L., Hou, H.Y., Zhu, Z. C., Rai, V., He, X. Y., & Tian, C. J. (2013). Differences in the arbuscular mycorrhizal fungi-improved rice resistance to low temperature at two N levels: Aspects of N and C metabolism on the plant side. *Plant Physiology and Biochemistry*, 71, 87–95.
- López-Bucio, J., Cruz-Ramírez, A., & Herrera-Estrella L. (2003). The role of nutrient availability in regulating root architecture. *Current Opinion in Plant Biology*, 6(3), 280–287.
- Lopez-Pedrosa, A., González-Guerrero, M., Valderas, A., Azcón -Aguilar, C., & Ferrol, N. (2006). GintAmt1 encodes a functional high-affinity ammonium transporter that is expressed in the extraradical mycelium of *Glomus intraradices*. *fungus genetics and biology*, 43, 102–110.
- Makhziah, K., Rochiman, R., & Purnobasuki, H. (2013). Effect of nitrogen supply and genotypic variation for nitrogen use efficiency in maize. *American Journal of Experimental Agriculture*, 3(1), 182–199.
- Masoumeh, F., Wichmann, S., Vierheilg, H., & Kaul, H. P. (2009). The effects of arbuscular mycorrhiza and nitrogen nutrition on growth of chickpea and barley. *Pflanzenbauwissenschaften*, 13 (1), 15–22.
- Mathur, N., & Vyas, A. (1996). Relative efficiency of different VAM fungi on growth and nutrient uptake in *Ziziphus mauritiana*. *Indian Forester*, 19, 129–131.
- Mensah, J. A., Koch, A. M., Antunes, P. M., Hart, M. M., Kiers, E.T., & Bücking, H. (2015). High functional diversity within arbuscular mycorrhizal fungal species is associated with differences in phosphate and nitrogen uptake and fungal phosphate metabolism. *Mycorrhiza*, 25, 533–546.
- McDonald, S., Prenzler, P.D., Antolovich, M., & Robard, K. (2001). Phenolic content and antioxidant activity of olive extracts. *Food Chemistry*, 73, 73–84.
- Mortimer, P. E., Perez-Fernandez, M. A., & Valentine, A. J. (2009). Arbuscular mycorrhizae affect the N and C economy of nodulated (*Phaseolus vulgaris* L.) during NH<sub>3</sub> nutrition. *Soil Biology & Biochemistry*, 41, 2115–2121.
- Ordookhani, K., & Zare, M. (2011). Effect of Pseudomonas, Azotobacter and Arbuscular Mycorrhiza Fungi on Lycopene, Antioxidant Activity and Total Soluble Solid in Tomato (*Lycopersicon Esculentum* F1 Hybrid, Delba). *Advances in Environmental Biology*, 5(6), 1290–1294.
- Pérez-Tienda, J., Valderas, A., Camañes, G., García-Agustín, P., & Ferrol, N. (2012). Kinetics of NH<sub>4</sub><sup>+</sup> uptake by the arbuscular mycorrhizal fungus *Rhizophagus irregularis*. *Mycorrhiza*, 22, 485–491.
- Planchette, C., Fortin, J. A., & Furlan, V. (1983). Growth response of several plant species to mycorrhiza in soil of moderate fertility. I. Mycorrhizal dependency under field condition. *Plant and Soil*, 70, 199–209.
- Sagi, M., Savidov, N. A., Lvov, N. P., & Lips, S. H. (1997). Nitrate physiological reductase and molybdenum cofactor in annual ryegrass as affected by salinity and nitrogen source. *Plant Physiology*, 99, 546–53.
- Smith, S. E., & Read, D. J. (1997). *Mycorrhizal Symbiosis*, second ed. Academic Press Inc London, UK.
- Tufenkci, S., Sonmez, F., & Sensoy, R. (2005). Effect of arbuscular mycorrhiza fungi inoculation and phosphours and nitrogen fertilization on some plant growth parameter and nutrient content of chickpea. *Journal of Biological Sciences*, 5(6), 738–743.
- Vazquez, M. M., Barea, J. M., & Azcon, R. (2002). Influence of arbuscular mycorrhizae and a genetically modified strain of *Sinorhizobium* on growth, nitrate reductase activity and protein content in shoots and roots of *Medicago sativa* as affected by nitrogen concentrations. *Soil Biology & Biochemistry*, 34, 899–905.
- Wu, Q., Zou, Y., Xia, R., & Wang, M. (2007). Five *Glomus* species affect water relations of Citrus tangerine during drought stress. *Botany Study*, 48, 147–154.



- Zand-Parsa, S., Sepaskhah, A.R., & Ronaghi, A. (2006). Development and evaluation of integrated water and nitrogen model for maize. *Agricultural Water Management*, 81, 227–256.
- Zapata, F., & Zaharah, A. R. (2002). Phosphate availability from phosphate rock and sewage sludge as influenced by addition of water soluble phosphate fertilizers. *Nutrient Cycling in Agroecosystems*, 63,43–48.
- Zhao, D., Kane, M., Borders, B., & Harrison, M. (2008). Pine growth response to different site- preparation methods with or without post-plant herbaceous weed control on North Florida's Lower Coastal Plain. *Forest Ecology and Management*, 255(7), 2512–2523.
- Zhu, X. C., Song, F. B., Liu, S. Q., & Liu, T. D. (2011). Effects of arbuscular mycorrhizal fungus on photosynthesis and water status of maize under high temperature stress. *Plant and Soil*, 346, 189–199.



## کاهش مصرف کود نیتروژن در خیار با استفاده از مایه زنی گیاه با میکوریزا

مریم حقیقی\*، بهزاد عبداللهی پور

گروه باغبانی، دانشکده کشاورزی، دانشگاه صنعتی اصفهان، اصفهان، ج.ا. ایران

\*نویسنده مسئول

### اطلاعات مقاله

#### تاریخچه مقاله:

تاریخ دریافت: ۱۳۹۷/۲/۳۱

تاریخ پذیرش: ۱۳۹۷/۱۲/۹

تاریخ دسترسی: ۱۳۹۸ /۳/۱۸

#### واژه‌های کلیدی:

فعالیت آنتی اکسیدانی

خیار

مایه‌زنی

میکوریزا

نیتروژن

فعالیت آنزیم نیترات رداکتاز

**چکیده-** این مطالعه به منظور بررسی تأثیر کاربردهای مختلف نیتروژن بر رشد، میزان فنل، آنتی‌اکسیدان و فعالیت آنزیم نیترات رداکتاز همراه با مایه‌زنی میکوریزا در رقم Super-N3 خیار به صورت فاکتوریل در قالب یک طرح کاملاً تصادفی با ۶ تکرار با ۳ سطح نیتروژن NO<sub>3</sub>-50، NO<sub>3</sub>-75 و NO<sub>3</sub>-100 و ۳ سطح مایه‌زنی میکوریزا شامل مایه‌زنی ۱۰۰۰ اسپور (AM1)، مایه‌زنی ۲۰۰۰ اسپور (AM2) و بدون مایه‌زنی میکوریزا (AM0) مورد بررسی قرار گرفت. نتایج حاصل از ترکیبات نیتروژن با مایه‌زنی میکوریزا نشان داد که وزن تر ریشه با مایه‌زنی میکوریزا در تمام سطوح نیتروژن افزایش یافت، درحالی‌که وزن تر ساقه با سطوح نیتروژن و مایه‌زنی میکوریزا تحت تأثیر قرار نگرفت. FV / FM در گیاهان با سطح NO<sub>3</sub>-75 و مایه‌زنی میکوریزا در مقایسه با تیمار بدون مایه‌زنی میکوریزا بالاتر بود. همزیستی میکوریزا فعالیت آنتی‌اکسیدانی گیاه را در تیمار بدون نیتروژن افزایش داد، به طوری‌که AM1 و AM2 فعالیت آنتی‌اکسیدانی را به ترتیب در NO<sub>3</sub>-50 و NO<sub>3</sub>-75 نسبت به گیاهان بدون مایه‌زنی میکوریزا افزایش دادند. بالاترین فعالیت آنزیم نیترات رداکتاز در NO<sub>3</sub>-50 و مایه‌زنی میکوریزا مشاهده شد و فعالیت آنزیم نیترات رداکتاز گیاهان در تمام سطوح نیتروژن کاهش یافت. در کل، می‌توان نتیجه گرفت که با مایه‌زنی میکوریزا، به ویژه تیمار ۱۰۰۰ اسپور (AM1)، و سطح متوسط NO<sub>3</sub> (NO<sub>3</sub>-75) می‌تواند مورد استفاده قرار گیرد و این میزان مصرف می‌تواند برای تولید خیار گلخانه‌ای بسیار موثر باشد.