

Oxidation of Sulfides to Sulfoxides by NaBrO_3 - NH_4Cl in Aqueous Acetonitrile

Shaabani, Ahmad*, Safaei, Hamid Reza and Bazgir, Ayoob

Department of Chemistry, University of Shahid Beheshti, P.O. Box 19396-4716, Tehran, I. R. Iran

ABSTRACT: NaBrO_3 combined with NH_4Cl is found to be an efficient reagent for the transformation of sulfides into sulfoxides in aqueous acetonitrile and under mild conditions.

KEY WORDS: Sulfide, Sulfoxide, Oxidation, Sodium bromate, Ammonium chlorid

INTRODUCTION

Sulfoxide synthesis constitutes an important research area [1-7] as they are versatile building blocks in organic synthesis [8,9]. A great number of oxidizing agents can effect the conversion of sulfides into sulfoxides [10-15]. However, the susceptibility of sulfoxide to further oxidation narrows the choice of reagents for the oxidation of sulfides to sulfoxides. Introduction of new oxidants for the transformation of sulfides to sulfoxides under mild condition is of importance in synthetic organic chemistry. NaBrO_3 and AgBrO_3 have been used for the oxidation of varieties of organic compounds in the presence of Lewis acids in aprotic organic solvents [16]. Recently mixture of NaBrO_3 - NH_4Cl is used as efficient oxidant for the oxidation of alcohols [17]. In this paper oxidation of sulfides to sulfoxides using NaBrO_3 - NH_4Cl is reported.

This reagent is a cheap, safe and very convenient one for the oxidation reaction of sulfides, compared with other oxidizing agents.


Table 1, summarizes the results of various sulfide oxidation with NaBrO_3 - NH_4Cl to corresponding sulfoxides in good yields. Sulfones were not detected in the crude mixture, which indicates high selectivity of the present method.

Sodium bromate in the absence of NH_4Cl is also able to transform dialkyl sulfides to corresponding sulfoxides (entry 9-11), but this reagent alone is not capable of oxidizing aryl sulfide (entry 12-14).

To study the oxidation ability of NaBrO_3 - NH_4Cl , a few reaction is run by using $\text{Bu}_4\text{NIO}_4/\text{AlCl}_3$ [15], $\text{Bi}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}/\text{HOAc}$ [18], $\text{Ba}(\text{MnO}_4)_2$ [13] and $\text{H}_2\text{O}_2/\text{MeCN}/\text{K}_2\text{CO}_3/\text{CH}_3\text{OH}$ [19] (Table 2).

EXPERIMENTAL

All products are known compounds and were identified by comparison of their physical and spectral data with those of authentic samples. Melting points were determined in open capillaries using an


























* To whom correspondence should be addressed.





oil-bath and are uncorrected. IR spectra were recorded as neat films or as KBr pellets on a Shimadzu 470 spectrophotometer. ^1H NMR spectra were recorded at

90 MHz on a JEOL EX-90 instrument with CDCl_3 as solvent and Me_4Si as an internal standard. All sulfides were purchased from Fluka, Aldrich and Merck

Table 1: Oxidation of sulfide to sulfoxide compounds by NaBrO_3 - NH_4Cl in acetonitrile-water (7:3 v/v) at 40°C

Entry	Substrate	Product	Reaction time (t/h)	Yield ^a (%)	m.p.($^\circ\text{C}$) or b.p.($^\circ\text{C}$)	
					Found	Report
1			2.5	95	190-192	189 ^b
2			4	95	103-105	104 ^b
3			3.5	90	28-30	32.6 ^b
4			1.5	95	75-76	71-72 ^c
5			30	85	73-74	70.5 ^d
6			3	90	30-33	33-34 ^c
7			4	90	147-148.13	146.13 ^c
8			3.5	90	135-137	133-135 ^c
9 ^e			4	85	190-191	189 ^b
10 ^e			4	80	28-30.6	32.6 ^b
11 ^e			4	75	75-76	71-72 ^c
12 ^e		no reaction	6	—	—	—

Continued

Entry	Substrate	Product	Reaction time (t/h)	Yield ^a (%)	m.p.(°C) or b.p.(°C)	
					Found	Report
13 ^e		no reaction	4	—	—	—
14 ^e		no reaction	4	—	—	—
15 ^f			8	—	—	—

a) Yield refers to isolated product

b) From ref. 20

c) From ref. 21

d) From ref. 22

e) In the absence of NH_4Cl

f) Isolated sulfoxide without any transformation to sulfone

Table 2: Comparison of the results of NaBrO_3 - NH_4Cl with those obtained from $\text{Bu}_4\text{NIO}_4/\text{AlCl}_3$ (1) [15], $\text{Bi}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}/\text{HOAc}$ (2) [18], $\text{Ba}(\text{MnO}_4)_2$ (3) [13] and $\text{H}_2\text{O}_2/\text{MeCN}/\text{K}_2\text{CO}_3/\text{CH}_3\text{OH}$ (4) [19]

Entry	Substrate	Product	Yield % (h)	Yield % (h) reported by other methods			
			NaBrO_3 - NH_4Cl	1	2	3	4
1	$(\text{C}_4\text{H}_9)_2\text{S}$	$(\text{C}_4\text{H}_9)_2\text{SO}$	90(3.45)	—	68(2)	—	84(0.5)
2	$\text{C}_6\text{H}_5\text{SCH}_3$	$\text{C}_6\text{H}_5\text{SOCH}_3$	90(3)	70(4)	—	—	82(2)
3	$(\text{C}_6\text{H}_5\text{CH}_2)_2\text{S}$	$(\text{C}_6\text{H}_5\text{CH}_2)_2\text{SO}$	90(3.5)	75(4)	85(4)	77(4)	—

Chemical Companies.

Oxidation of Dibenzyl Sulfide (entry 8)

Typical Procedure

Dibenzyl sulfide (1.072 g, 5 mmol) was added to a mixture of NaBrO_3 (0.755 g, 5 mmol) and NH_4Cl (0.400 g, 7.5 mmol) in aqueous acetonitrile (CH_3CN - H_2O ; 7:3 v/v; 10 mL) and stirred at 40°C for 3.5 h. The resulting mixture was extracted with methylene chloride (20 mL \times 2). The combined organic layers were washed with a saturated aqueous solution of NaHCO_3 (15 mL) and dried over anhydrous MgSO_4 and filtered. Evaporation of the filtrate afforded dibenzyl sulfoxide as a colorless solid, which was purified by crystallization from EtOH to afford the desired product, yield, 1.037 g, (90%), m.p. 135-137°C (lit. 133-135°C), [22].

ACKNOWLEDGMENTS

Financial support by research council of Shahid Beheshti University is gratefully acknowledged.

Received: 13th January 1999; Accepted: 24th January 2000

REFERENCES

- [1] a) Patai, S. and Rappoport, z., "Chemistry of Sulfoxides and Sulfones", Academic Press, New-York (1998), b) Block, E., "Reaction of Organo Sulfur Compounds", Academic Press, New York (1978).
- [2] Madesclaire, M., *Tetrahedron*, **42**, 5459(1986).
- [3] Hudlicky, M., "Oxidations in Organic Chemistry", Am. Chem. Soc. Publication, Washington, DC, pp. 252-259(1990).
- [4] Orito, K., Hatakeyama, T., Takeo, M. and Sugi-

- nome, H., *Synthesis*, 1357(1995).
- [5] Firouzabadi, H., Iranpoor, N. and Zolfigol, M. A., *Synth. Commun.*, **28**, 377(1998).
- [6] Heshmat Ali, M., Leach, D. R. and Schmitz, C. E., *Synth. Commun.*, **28**, 2969(1998).
- [7] Ravikumar, K. S., Begue, J. P. and Delpon, D. B., *Tetrahedron Lett.*, **39**, 3141(1998).
- [8] Trost, B. M., *Acc. Chem. Res.*, **11**, 453(1978).
- [9] Durst, T., "In Comprehensive Organic Chemistry", Barton, D., Ollis, W. D., Eds., Vol. 3, Pergamon, Oxford (1979).
- [10] Shaabani, A., Teimoori, M. B. and Safaei, H. R., *Synth. Commun.*, In Press.
- [11] Bellesia, F., Gheefi, F., Panoni, U. M. and Pinetti, A., *Synth. Commun.*, **23**, 1759(1993).
- [12] Mata, E. G., *Phosphorus, Sulphur and Silicon*, **117**, 231(1996).
- [13] Firouzabadi, H. and Seddighi, M., *Synth. Commun.*, **21**, 211(1991).
- [14] Firouzabadi, H. and Mohammadpoor Baltork, I., *Bull. Chem. Soc. Jpn.*, **65**, 1131(1992).
- [15] Firouzabadi, H., Sardarian, A. and Badparva, H., *Bull. Chem. Soc. Jpn.*, **69**, 685(1996).
- [16] Firouzabadi, H. and Mohammadpoor Baltork, I., *Bull. Chem. Soc. Jpn.*, **68**, 2319(1995).
- [17] Shaabani, A. and Ameri, M., *J. Chem. Res.(S)*, 100(1998).
- [18] Mashraqui, S. H., Mudaliar, C. D. and Karnik, M. A., *Synth. Commun.*, **28**, 939(1998).
- [19] Bullman Page, P. C., Graham, A. E., Bethell, D. and Park, B. K., *Synth. Commun.*, **23**, 1507(1993).
- [20] "Handbook of Chemistry and Physics", (CRC Press), 70th ed., (1989-1990).
- [21] "Dictionary of Organic Compounds", 6th ed., Chapman & Hall, London (1966).
- [22] Orito, H., Hatakeyama, T., Takeo, M. and Sugimoto, H., *Synthesis*, 1357(1995).

Entry	Substrate	Product	Yield (%)	Yield (%)
1	(C ₆ H ₅) ₂ S	(C ₆ H ₅) ₂ S	90(4)	90(4)
2	C ₆ H ₅ SC ₂ H ₅	C ₆ H ₅ SC ₂ H ₅	90(4)	90(4)
3	(C ₆ H ₅ CH ₂) ₂ S	(C ₆ H ₅ CH ₂) ₂ S	90(4)	90(4)

ACKNOWLEDGMENTS

Financial support by research council of Shahrood University of Technology is gratefully acknowledged.

REFERENCES

[1] J. E. McGrath, *Chem. Rev.*, **80**, 1473 (1980).

[2] M. J. Fréchet, *Chem. Rev.*, **80**, 1473 (1980).

[3] J. E. McGrath, *Chem. Rev.*, **80**, 1473 (1980).

[4] Orito, H., Hatakeyama, T., Takeo, M. and Sugimoto, H., *Synthesis*, 1357(1995).

[5] Firouzabadi, H. and Mohammadpoor Baltork, I., *Bull. Chem. Soc. Jpn.*, **65**, 1131(1992).

[6] Firouzabadi, H., Sardarian, A. and Badparva, H., *Bull. Chem. Soc. Jpn.*, **69**, 685(1996).

[7] Firouzabadi, H. and Mohammadpoor Baltork, I., *Bull. Chem. Soc. Jpn.*, **68**, 2319(1995).

[8] Shaabani, A. and Ameri, M., *J. Chem. Res.(S)*, 100(1998).

[9] Mashraqui, S. H., Mudaliar, C. D. and Karnik, M. A., *Synth. Commun.*, **28**, 939(1998).

[10] Bullman Page, P. C., Graham, A. E., Bethell, D. and Park, B. K., *Synth. Commun.*, **23**, 1507(1993).

[11] "Handbook of Chemistry and Physics", (CRC Press), 70th ed., (1989-1990).

[12] "Dictionary of Organic Compounds", 6th ed., Chapman & Hall, London (1966).

[13] Orito, H., Hatakeyama, T., Takeo, M. and Sugimoto, H., *Synthesis*, 1357(1995).