# Cellulose Sulfuric Acid: As an Efficient Bio Polymer Based Catalyst for the Selective Oxidation of Sulfides and Thiols by Hydrogen Peroxide

Shaabani, Ahmad\*<sup>+</sup>; Ganji, Nasim; Seyyedhamzeh, Mozhdeh; Mofakham, Hamid

Faculty of Chemistry, Shahid Beheshti University, P.O. Box 19396-4716 Tehran, I.R. IRAN

**ABSTRACT:** Cellulose sulfuric acid as a bio-polymer based solid catalyst efficiently catalyzes the selectively oxidation of sulfides to sulfoxides and thiols to disulfides using hydrogen peroxide as a green oxidant with good yields at room temperature. The developed method offers a number of advantages such as high selectivity, mild reaction conditions, simple operation, cleaner reaction profiles, low cost and biodegradability of the catalyst.

KEY WORDS: Cellulose sulfuric acid, Oxidation, Sulfide, Thiol, Hydrogen peroxide.

### **INTRODUCTION**

The oxidation of organosulfurs is of a great importance in synthesis of drug, drug metabolism and the preparation of bioconjugates compounds [1,2]. In addition, the removal of sulfur from fuels, and industrial products and wastes is important in chemical industry. In this regard, several new methods were recently proposed for the removal of sulfur by oxidation [1-3].

Heterogeneous solid acid catalysts for the synthesis of fine chemicals have attracted considerable attention from both environmental and economic points [4]. Majority of the reported supports for catalytic applications is based on inorganic materials or functional synthetic polymers in which the synthesis of the supports involves high temperature calcinations or polymerization of petrochemical base feed stocks [6,7]. Biopolymers, especially cellulose and its derivatives [8] which have some unique properties are attractive and alternative supports for catalytic process in synthesis of conventional organic or inorganic compounds. Cellulose is one of

7/\$/2.70

the most abundant natural bio-polymers in the world and has been widely studied during the past decades because of it is a biodegradable material and a renewable resource.

In this context, Cellulose Sulfuric Acid (CSA) is one of the useful heterogeneous catalysts which has a good performance as an inexpensive biopolymer-based catalyst and can be easily separated without contaminating the products [9]. The aim of this study is to utilize cellulose supported sulfuric acid as a catalyst for the selective oxidation of sulfides to sulfoxides and thiols to disulfides using hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) as a green oxidant.Over other traditional oxidants, H<sub>2</sub>O<sub>2</sub> has many advantages. The major benefit of using H<sub>2</sub>O<sub>2</sub> is its environmental concern in view of effective-oxygen content (47%) and more importantly, entire process is very clean since water is the only side product[10-13]. Furthermore, H<sub>2</sub>O<sub>2</sub> has acceptable safety in storage and operation [14, 15]. Hydrogen peroxide is a mild oxidizing agent, and often activated by homogeneous [16-19] or heterogeneous [20-22] catalysts.

<sup>\*</sup> To whom correspondence should be addressed.

<sup>+</sup>E-mail: a-shaabani@sbu.ac.ir

<sup>1021-9986/13/3/1</sup> 



Scheme 1: Selective oxidation of thiols and sulfides by hydrogen peroxide in the presence of a catalytic amount of CSA.

In connection with our studies on the development of new routes for the oxidation of organic compounds [23-29] and exploring CSA as a solid acid catalyst [9,30-33], we wish to report herein the selectively oxidation of alkyl and aryl sulfides to sulfoxides and thiols to disulfides, in good yields with aqueous hydrogen peroxide in the presence of CSA as an efficient solid acid catalyst (Scheme 1).

### **EXPERIMENTAL SECTION**

#### Apparatus

Melting points were measured on an Electrothermal 9200 apparatus. The chemicals used in this work were purchased from Merck and Fluka chemical companies. Conversion of the sulfides and selectivity for the sulfoxides was determined by GC by using Agilent GC with FID detector.

# General experimental procedure for the oxidation of sulfides and thiols

In a round bottom flask, to a solution of sulfide or thiol (1 mmol) in CH<sub>3</sub>CN (5 ml), 30% (w/w) H<sub>2</sub>O<sub>2</sub> (1mmol) and the CSA catalyst (0.05 g for oxidation of sulfides and 0.1 g for oxidation of thiols) was added successively and the mixture was stirred magnetically at room temperature for the indicated time in Table 3. The progress of the reaction was monitored by TLC or GC. After completion of the reaction, the reaction mixture was filtered and the excess of solvent was removed under reduced pressure to give the corresponding pure sulfoxide or disulfide (Table 3, Entries 1-15). All the products are known and they were characterized by melting point which were compared with those obtained from authentic samples [34-41].

# **RESULTS AND DISCUSSION**

Cellulose sulfuric acid is readily prepared by the dropwise addition of chlorosulfonic acid to cellulose in CHCl<sub>3</sub> at  $0^{\circ}$ C, as our previously reported method [9]. It is important to note that, this reaction is easy and clean without any further work-up due to the release of HCl gas from the reaction vessel immediately. The white homogeneous and nonhygroscopic solid acid is very stable under reaction conditions.

To study the catalytic oxidation of sulfides mediated by CSA, thioanisole was chosen as a model compound and reaction conditions were optimized. The yield and rate of the reaction were also found to be dependent on the amount of the catalyst (Table 1). The optimum molar ratio of sulfide: $H_2O_2$  (1:1) in the presence of CSA (0.05 g) is found to be ideal for complete conversion of sulfides to sulfoxide with no observation of sulfone. The catalyst is very active, stable to air and moisture, nontoxic and inexpensive.

The choice of the organic solvent is of particular importance. The reaction was carried out in different solvents and the results are tabulated in Table 2. The reaction gave very good yields in acetonitrile. It is comparatively slower in non-polar solvents such as toluene. In water, it affords lower yield.

To indicate the scope of reaction, a series of sulfides were chosen and subjected to the oxidation reaction.

Entry	CSA (g)	t (h)	Isolated Yield (%)		
1	0	30	35		
2	0.015	24	73		
3	0.025	7.5	81		
4	0.35	6	89		
5	0.05	4.5	92		
6	0.075	4.5	93		

Table 1: Effect of amount of catalyst on the oxidation of thioanisole in CH<sub>3</sub>CN.

Table 2: Influence of different solvents on oxidation of thioanisole.

	0 0 00		
Entry	Solvent	t(h)	Isolated Yield (%)
1	Acetonitrile	4.5	92
2	Acetone	6	81
3	Methanol	5	84
4	Ethanol	9	73
5	Toluene	14	66
6	Chloroform	9	68
7	Water	15	45

The results summarized in Table 3 (Entries 1-9) and they clearly indicated the generality of the reaction with respect to variety of alkyl and aryl sulfides.

In an effort to expand CSA efficiency within the family of organosulfur species, we turned our attention toward exploring the oxidation of thiols to their respective disulfides. In the first step, we carried out a set of initial experiments on benzothiol as a model substrate using 30%  $H_2O_2$  in the presence of different amounts of catalyst (Table 4) in various solvents (Table 5) at room temperature. The best result was obtained by carrying out the reaction with molar ratios of thiol: $H_2O_2(1:1)$ in the presence of 0.1 g of CSA in acetonitrile as solvent for 30 min.

The oxidative couplings of other thiols were then examined using the optimized reaction conditions and oxidation performed almost in excellent yields (Table 3, entries 10-15).

## CONCLUSIONS

In summary, we have developed an efficient and simple procedure for the selective oxidation of sulfides and thiols to their corresponding sulfoxides and disulfides with using  $H_2O_2$  in the presence of CSA as an environmentally friendly bio-supported proton source catalyst. Inexpensive nature and availability of the reagents, easy and clean workup process and the ability to be operated at room temperature are features of this new procedure to selective oxidation of sulfides and thiols.

### Acknowledgement

We gratefully acknowledge financial support from the Iran National Elites Foundation (INEF) and Research Council of Shahid Beheshti University.

Entry	Substrate	Product	t (h)	Isolated Yield (%)	M.p.(°C)	
					Found	Reported[Ref.]
1	S CH3	O S CH <sub>3</sub>	4.5	92	28-30	30-30.5 [34]
2	CI	CI CH3	5.5	87	135-138	136-137[35]
3	S CH3	O CH <sub>3</sub>	6	85	136-138	138-140 [35]
4	C) <sup>s</sup> C)		8	79	70-73	72-73 [34]
5	Br S CH3	Br CH <sub>3</sub>	5	93	71-73	74-76 [36]
6	€, s, , , , , , , , , , , , , , , , , ,		9	74	145-147	146 [36]
7		S-C	4	87	41-44	43-45 [36]
8	~~~s~~~	∽∽s∽∽ ö	12	71	27-29	29-32 [34]
9	~s~	∧s∧ ð	18	78	100-103	103-106 [37]
10	CH <sub>2</sub> SH	CH <sub>2</sub> S-SCH <sub>2</sub> -	0.5	90	70-73	69-72[38]
11	ClSH	CISSCI	1.5	89	68-70	70-71[39]
12	——————————————————————————————————————	-<	1.25	92	43-46	45-46[40]
13	K→SH	NS-S-S-KS	0.75	82	175-177	177-179[41]
14	N N H	$\underset{H}{\overset{N}{\underset{H}{\longrightarrow}}} S-S-\overset{N}{\underset{H}{\bigvee}} \overset{N}{\underset{H}{\longrightarrow}}$	0.5	79	200-203	202-204[41]
15	SH SH	CCC <sup>S-S</sup> CCC	1	88	137-139	139[40]

Table 3: Oxidation of sulfides to sulfoxides and thiols to disulfides with  $H_2O_2$  in the presence of CSAat room temperature in  $CH_3CN$ .

4

	00 0 0 0	U	
Entry	CSA (g)	t (h)	Isolated Yield (%)
1	0	8	77
2	0.025	5	81
3	0.05	2.5	85
4	0.075	2.5	87
5	0.1	0.5	90
7	0.12	0.5	91

Table 4: Effect of amount of catalyst on the oxidation of benzothiol in CH<sub>3</sub>CN.

Entry	Solvent	t (h)	Isolated Yield (%)		
1	Acetonitrile	0.5	90		
2	Acetone	3	86		
3	Methanol	2	79		
4	Ethanol	7	66		
5	Toluene	10	62		
6	Chloroform	5	78		
7	Water	10	55		

Table 5: Influence of different solvents on oxidation of benzothiol.

Received : Sep. 22, 2013 ; Accepted : Mar. 9, 2014

## REFERENCES

- Patai S., Rappoport Z., Stirling C.J.M., "The Chemistry of Sulphones and Sulphoxides", Chichester: John Wiley & Sons, (1988).
- [2] Bäckvall J.E., "Modern Oxidation Methods", Hoboken, NJ: Wiley-VCH, (2011).
- [3] Dreyer D. R., Jia H. P., Todd A. D., Geng J., Bielawski C. W., Graphite Oxide: a Selective and Highly Efficient Oxidant of Thiols and Sulfides, Org. Biomol. Chem., 9: 7292-7295 (2011).
- [4] Dhakshinamoorthy A., Alvaro M., Garcia H., Aerobic Oxidation of Thiols to Disulfides Using Iron Metal-Organic Frameworks as Solid Redox Catalysts, Chem. Commun., 46: 6476-6478 (2010).
- [5] Sheldon R.A., Bekkum H., "Fine Chemicals Through Heterogeneous Catalysis", Weinheim: Wiley-CH, (2001).
- [6] Clarck J. H., Macquarrie D. J., "Green Chemistry and Technology", Abingdon: Blackwell, (2002).
- [7] Guibal E., Heterogeneous Catalysis on Chitosan-Based Materials: a Review, *Prog. Polym. Sci.*, 30: 71-109 (2005).

- [8] Klemm D.,Heublein B., Fink H. P.,Bohn A.,Cellulose: Fascinating Biopolymer and Sustainable Raw Material, Angew. Chem. Int. Ed., 44: 3358- (2005).
- [9] Shaabani A., Maleki A., Moghimi-Rad J., Soleimani E., Cellulose Sulfuric Acid Catalyzed One Pot Three-Component Synthesis of Imidazoazines, *Chem. Pharm. Bull.*, 55: 957-958 (2007).
- [10] GrigoropoulouG., Clark J., Elings J., Recent Developments on the Epoxidation of Alkenes Using Hydrogen Peroxide as an Oxidant, *Green Chem.*, 5: 1-7 (2003).
- [11] Noyori R., Aoki M., SatoK., Green Oxidation with Aqueous Hydrogen Peroxide, Chem. Commun., 1977-1986 (2003).
- [12] Strukul G., "Catalytic Oxidations with Hydrogen Peroxide as Oxidant", London: Kluwar Academic Publishers, (1992).
- [13] Trost B. M., Masuyama Y., Chemoselectivity in Molybdenum Catalyzed Alcohol and Aldehyde Oxidations, *Tetrahedron Lett.*, 25: 173-176 (1984).

- [14] Arends I., Sheldon R., Recent Developments in Selective Catalytic Epoxidations with H<sub>2</sub>O<sub>2</sub>, *Top. Catal.*, **19**: 133-141 (2002).
- [15] Ramirez-Verduzco L., Torres-Garcia E.,Gomez-Quintana R., Gonzalez-Pena V., Murrieta-Guevara F., Desulfurization of Diesel by Oxidation/Extraction Scheme: Influence of the Extraction Solvent, *Catal. Today*, **98**: 289- 294 (2004).
- [16] Kuhn F. E., Santos A. M., Herrmann W. A., Organorhenium(VII) and Organomolybdenum(VI) Oxides: Syntheses and Application in Olefin Epoxidation, *Dalton Trans.*, 2483-2491 (2005).
- [17] Kuhn F. E., Scherbaum A., Herrmann W. A., Methyltrioxorhenium and Its Applications in Olefin Oxidation, Metathesis and Aldehyde Olefination, *J. Organomet. Chem.*, 689: 4149-4164 (2004).
- [18] Al-Ajlouni A. M., Espenson J. H., Epoxidation of Styrenes by Hydrogen Peroxide As Catalyzed by Methylrhenium Trioxide, J. Am. Chem. Soc., 117: 9243-9250 (1995).
- [19] Al-Ajlouni A. M., Espenson J. H., Kinetics and Mechanism of the Epoxidation of Alkyl-Substituted Alkenes by Hydrogen Peroxide, Catalyzed by Methylrhenium Trioxide., J. Org. Chem., 61: 3969-3976 (1996).
- [20] Ziolek M., Catalytic Liquid-Phase Oxidation in Heterogeneous System as Green Chemistry Goal-Advantages and Disadvantages of MCM-41 Used as Catalyst, Catal. Today,90: 145-150 (2004).
- [21] Sheldon R.A., Dakka J., Heterogeneous Catalytic Oxidations in the Manufacture of Fine Chemicals, *Catal. Today*, **19**: 215-245 (1994).
- [22] Sheldon R.A., Arends I., Lempers H., Liquid Phase Oxidation at Metal Ions and Complexes in Constrained Environments, *Catal. Today*, **41**: 387-407 (1998).
- [23] Shaabani A., Behnam M., Rezayan A.H., Tungstophosphoric Acid (H<sub>3</sub>PW<sub>12</sub>O<sub>40</sub>) Catalyzed Oxidation of Organic Compounds with NaBrO<sub>3</sub>, Catal. Commun., 10: 1074-1078 (2009).
- [24] Shaabani A., Rezayan A. H., Silica Sulfuric Acid Promoted Selective Oxidation of Sulfides to Sulfoxides or Sulfones in the Presence of Aqueous H<sub>2</sub>O<sub>2</sub>, *Catal. Commun.*, 8: 1112-1116 (2007).

- [25] Shaabani A., Naderi S., Rahmati A., Badri Z., Darvishi M., Lee D.G., Cleavage of Oximes, Semicarbazones, and Phenylhydrazones with Supported Potassium Permanganate, Synthesis, 3023-3025 (2005).
- [26] Shaabani A., Mirzaei P., Naderi S., Lee D.G., Green Oxidations. The Use of Potassium Permanganate Supported on Manganese Dioxide, *Tetrahedron*, 60: 11415-11420 (2004).
- [27] Shaabani A., Bazgir A., Teimouri F., Lee, D. G., Selective Oxidation of Alkylarenes in Dry Media with Potassium Permanganate Supported on Montmorillonite K10, *Tetrahedron Lett.*, 43: 5165-5167 (2002).
- [28] Shaabani A., Lee D.G., Solvent Free Permanganate Oxidations, Tetrahedron Lett., 42: 5833-5836 (2001).
- [29] Shaabani A., Mofakham H., Rahmati A., Farhangi E., 1-Butyl-3-Methylimidazolium Bromide Promoted Selectively Oxidation of Sulfur Compounds by NaBrO<sub>3</sub>, *Iran. J. Chem. Chem. Eng. (IJCCE)*, **31**: 1-8 (2012).
- [30] Shaabani A., Maleki A., Cellulose Sulfuric Acid as a Bio-Supported and Recyclable Solid Acid Catalyst for the One-Pot Three-Component Synthesis of α-Amino Nitriles, *Appl. Catal. A: Gen.*, **331**:. 149-151 (2007).
- [31] Shaabani A., Rahmati A., Badri Z., Sulfonated Cellulose and Starch: New Biodegradable and Renewable Solid Acid Catalysts for Efficient Synthesis of Quinolines, Catal. Commun., 9: 13-16 (2008).
- [32] Shaabani A., Seyyedhamzeh M., Maleki A., Rezazadeh F., Cellulose Sulfuric Aacid: An Efficient Biopolymer-Based Catalyst for the Synthesis of Oxazolines, Imidazolines and Thiazolines Under Solvent-Free Cconditions, Appl. Catal. A: Gen., 358: 146-149 (2009).
- [33] Mofakham H., Hezarkhani Z., Shaabani A., Cellulose-SO<sub>3</sub>H as a biodegradable solid acid catalyzed one-Pot three-Component Ugi reaction: Synthesis of-Amino amide,3,4-Dihydroquinoxalin-2-Amine,4*H*-Benzo[*b*][1,4]Thiazin-2-Amine and 1,6-Dihydropyrazine-2,3-Dicarbonitrile Derivatives, *J. Mol. Catal. A: Chem.*, **360**: 26-34 (2012).

- [34] Sato K., Hyodo M., Aoki M., Zheng X. Q., Noyori R., Oxidation of Sulfides to Sulfoxides and Sulfones with 30% Hydrogen Peroxide under Organic Solvent- and Halogen-Free Conditions, Tetrahedron, 57: 2469-2476 (2001).
- [35] "The Sadtler Standard Spectra, NMR", Sadtler Research Laboratories, 12327M, (1972).
- [36] "Dictionary of Organic Compounds", Sixth ed., London, Chapman and Hall,(1996).
- [37] "Handbook of Chemistry and Physics", 93th ed., CRC Press, (2012).
- [38] Field L., Lawson J. E., Organic Disulfides and Related Substances. I. Oxidation of Thiols to Disulfides with Lead Tetraacetate, J. Am. Chem. Soc., 80: 838-841 (1958).
- [39] Khodaei M., Mohammadpoor-Baltork I., Nikoofar K., Bismuth(III) Nitrate Pentahydrate Bi(NO<sub>3</sub>)<sub>3</sub>·5H<sub>2</sub>O: An Inexpensive and Mild Reagent for the Efficient and Clean Oxidation of Thiols to Disulfides, Bull. Korean Chem.Soc.,24: 885-886 (2003).
- [40] Wallace T.J., Reactions of Thiols with Sulfoxides. I. Scope of the Reaction and Synthetic Applications, J. Am. Chem. Soc., 86: 2018-2021 (1964).
- [41] Karami B., Montazerozohori M., Moghadam M., Habibi M.H., Niknam K., Selective Oxidation of Thiols to Disulfides Catalyzed by Iron(III) Tetra Phenyl Porphyrin Using Urea-Hydrogen Peroxide as Oxidizing Reagent, Turk. J. Chem., 29: 539-546 (2005).