

The Effects of Water Deficit on Crop Yield and the Physiological Characteristics of Barley (*Hordeum vulgare L.*) Varieties

E. Mamnouie¹, R. Fotouhi Ghazvini¹, M. Esfahany^{1*} and B. Nakhoda²

ABSTRACT

The effects of water deficit on grain yield, yield components and the physiological characteristics of barley (*Hordeum vulgare L.*) varieties were studied in a split plot design during the 1999-2000 growing season in Karaj, Iran. Five irrigation levels [0, 25, 50, 75, and 100% crop water requirements (CWR)] and six barley varieties Karoon × Kavir, Reihani (drought tolerant), Torkman, C-74-9 (intermediate), Kavir × Badia and Gorgan-4 (2 rowed type, drought sensitive) were arranged in the main plots and sub plots respectively. Results showed that water deficit induced stress and this decreased the grain yield and yield components. Water deficit increased the canopy temperature and those varieties showed a higher canopy temperature under non-stress conditions, performed better under drought conditions. Severe stress reduced the chlorophyll content (SPAD values) considerably, but the differences were not significant between the 50, 75 and 100% CWR treatments. In addition, no significant differences were observed in the chlorophyll content of barley varieties under drought conditions. The effect of irrigation on the photochemical efficiency of photosystem II (F_v/F_m) and the other fluorescence parameters for all varieties were significant. Although the F_v/F_m values were not significant in barley varieties at any level of irrigation, in general Karoon × Kavir and Reihani varieties showed a better performance under water deficit conditions. Proline content was significantly different in various irrigation treatments, but its accumulation at any level of irrigation did not differ significantly in barley varieties. It was concluded that the higher canopy temperature (less negative ΔT) under well irrigated conditions and higher grain yield, 1000-grain weight, F_v/F_m values under water stress conditions could possibly be the proper criteria for screening the drought tolerant barley genotypes under field or laboratory conditions.

Keywords: Barley (*Hordeum vulgare L.*), Canopy temperature, Drought tolerance, Photosystem II, SPAD, Water deficit.

INTRODUCTION

Water deficit (drought) at different stages of barley's life cycle, (from seed germination to maturity) may adversely affect the final yield of the crop. The effect of drought on the yield of cereals depends on the duration and the severity of the stress [9]. Long-term drought under high temperature conditions, coinciding with the reproductive stage of the plant, may cause a decrease in the size

and number of the grains, and eventually its yield.

Drought may cause abortion of the embryonic sac, dehydration of the style and pollen and, hence, interference in pollination [8, 9]. Singh, *et al.* [10] found that drought reduces the number of the grains/spike and thereby yield [11]. Water deficit at the tillering stage had no significant effect on the number of grains on spikes but, at the flowering stage, it reduced grain weight significantly [9].

1. Faculty of Agriculture, University of Guilan, Rasht, Islamic Republic of Iran.

2. Seed and Plant Improvement Institute, Karaj, Islamic Republic of Iran.

* Corresponding author.



Since drought may affect osmotic adjustment and canopy temperature, these two can be used as an index for screening varieties of barley to drought stress. Varieties that showed warmer canopies under irrigated conditions had a lower stomatal conductance and higher yield under drought conditions [2]. Liu and Zhang [6] reported that, under water deficit, the difference between air and canopy temperatures (ΔT) in wheat varieties was significantly higher at spike exertion stage but it had no effect on the yield. Under irrigated conditions, the reduction in the yield was significantly related to ΔT . Chlorophyll content is positively correlated with the photosynthesis rate and soluble proteins [3, 13]. The chlorophyll content which was measured by SPAD. 502 correlated to the photosynthesis of leaves ($R^2=0.77^{**}$) [7]. A significant correlation between the SPAD reading and spectrophotometric method of chlorophyll determination ($R^2 = 0.95^{**}$) was also reported [13]. In addition, chlorophyll fluorescence has been accepted as a reliable index for drought tolerance screening of barley varieties. This phenomenon is a criterion for thylacoid membrane integrity and electron transfer efficiency from photosystem II to photosystem I [7]. The photochemical efficiency of photosystem II is determined by the F_v/F_m ratio, which is reduced during periods of drought stress.

Free proline accumulation can also be used as another index for screening the tolerance of varieties [5]. Water stress may cause a disorder in the mitochondrial membrane and proline oxidation to glutamate might therefore be decreased. Proline accumulation by way of the glutamic pathway may help osmotic adjustment, energy production and further absorption of the released ammonia from protein breakdown [11].

The objectives of this experiment were to evaluate the effects of water deficit on the physiological characteristics of barley varieties in order to exploit them for drought tolerance screening.

MATERIALS AND METHODS

Field experiments were conducted at the Seed and Plant Improvement Institute (SPII), Karaj (Iran) (35°N Latitude) during the growing season of 1999-2000 in a split plot design with three replications. Five levels of irrigation: 0, 25, 50, 75 and 100%, of crop water requirements and six barley varieties, namely: Karoon \times Kavir, Reihani (drought tolerant), Torkman, C-74-9 (intermediate), Kavir \times Badia and Gorgan-4 (2-rowed type) (sensitive)-all obtained from Cereals Department of Seed and Plant Improvement Institute (SPII)- were laid out in the main plots and sub plots respectively. Seeds were sown in 16 rows in 10×2.5 m plots (400 plants. m^{-2}) in Autumn. Plants were irrigated twice after sowing equally ($1100 \text{ m}^3/\text{ha}$) and irrigation treatments were applied after the plants established and after their further growth in Spring (since April, 4). Their evapotranspiration potential (ETP) was determined according to:

$$\text{ETP} = E_{\text{pan}} \times K_p \quad (1)$$

E_{pan} = Evaporation from Class A Evaporation pan

K_p = Evaporation pan coefficient (0.85 in karaj) (obtained from Meteorology Station of Karaj).

Evapotranspiration of the crop (ETC = Crop water requirement) was calculated according to the following equation:

$$\text{ETC} = \text{ETP} \times K_c \quad (2)$$

K_c = Crop coefficient (0.96, 1.12, 1.15, 0.95 and 0.65 for growth stages of all barley varieties; stem elongation, booting, heading, anthesis and grain filling, in karaj). (Data obtained from Cereals Dept. - Seed & Plant Development Inst. (SPII))

The amount of irrigation water for each treatment (T in mm) was calculated as follows:

$$T = \text{ETC} \times \text{Percent of crop water requirements for each treatment (mm)} \quad (3)$$

Irrigation water was applied according to 80% irrigation efficiency and effective rainfall (twice during growing season; 3.5 and 2.5 mm in April and May, respectively) at

10-day intervals and its volume was measured using a WSC¹ Phloem type III fixed in the irrigation ditches. The total amount of irrigation water was 0, 2107.2, 3739, 5786 and 6450 m³/ha applied 0, 4, 5, 5 and 5 times for 0, 25, 75 and 100% of crop water requirements, respectively.

Grain yield was determined by harvesting the eight middle rows of each plot at the maturity stage and the yield components (number of grains/spike and number of spikes. m²) were determined from the samples taken randomly (10 spikes from the remaining parts of each plot) at the dough stage. Canopy temperature was measured between 12:00-2 pm. using an infrared thermometer [Kane - May Co. UK] at anthesis for determination of the ΔT (temperature difference between canopy and ambient air). The apparatus was targeted at the canopy with a 10-20° angle from four directions and at least 1 m distance after initial calibration. Ambient air temperature was measured using a mercurial thermometer and ΔT calculated as following:

$$\Delta T = T_{\text{canopy}} - T_{\text{air}} \quad (4)$$

The effect of water deficit on the chlorophyll content of flag leaves was determined using a portable chlorophyll meter (SPAD 502-Minolta Co. Japan) at the heading stage. The measurements were performed twice with an interval of one week between each

and 30 flag leaves were chosen randomly and the SPAD value recorded. Chlorophyll fluorescence and the parameters of F_0 , F_v , F_m and F_v / F_m (Fluorescence Initial, Fluorescence Maximum and Fluorescence Variable, respectively) were measured on the flag leaves of barley varieties by means of a plant stress meter (Biomonitor 1989, UK) at the heading stage. Measurements were carried out on 30 flag leaves from each plot five days after every irrigation, subsequent to the canopies approximate closure. The broadest part of the flag leaves was clipped in the measuring chamber of the apparatus and exposed to 650 nm wavelength radiation and the F_0 , F_m and F_v values were measured after initial calibration. All measurements were carried out at 10 am in order to avoid the effects of dew and air humidity. The free proline content was determined according to Bates *et al.* [1] at different irrigation levels in 15 flag leaves from each plot at the heading stage. 0.5 gr of leaf samples were crushed in a precooled mortar and pestle with a small amount of sand and 10 ml 3% (W/V) solfocalicylic acid. Leaf extracts, Ninhydrin and glacial acetic acid (equal amounts) were incubated in a water bath for one hour and the proline isolated by adding 4 ml Toluene to each and measured in 520 nm in a spectrophotometer (M 330- 230 Jenway – UK). The proline content was determined against its standard curve and converted to μg proline in 1gr fresh weight. All

Table 1. Summery analysis of variance for barley varieties parameters recorded at different irrigation levels.

S.O.V.	Mean Squares (MS)							
	1000 Grain Wt.	No. Spikes/m ²	No. Grains/Spike	Grain yield	ΔT	SPAD Values	F_v/F_m	Proline Content
Irrigation regime	348.75**	121803.74**	246.19**	0.43**	256.71**	229.6**	0.349**	36412.7**
E _a	4.599	1868.86	10.72	0.013	1.47	7.517	0.001	19.36
Variety	35.823**	93934.95**	577.72**	0.06**	16.29**	30.5**	0.31**	33.11*
Irrigation × variety	9.206 ^{n.s}	8748.98**	11.83 ^{n.s}	0.008*	4.19**	3.247 ^{n.s}	0.003**	7.79 ^{n.s}
E _b	7.543	1648.37	8.15	0.007	0.96	7.67	0.001	14.38

*, ** Significant at 0.05 and 0.01, respectively

n.s Non significant

**Table 2.** Irrigation regime and variety effects on the performance of barley varieties.

	1000 Grain Wt. (gr)	No. Spikes/m ²	No. Grains/ Spike	Grain yield (Kg/m ²)	ΔT (°C)	SPAD Values	F _v /F _m	Proline Content μg.gr ⁻¹ fw	
Irrigation	100%	40.56 a	499.8 a	31.32 a	0.635 a	-2.89 d	52.76 a	0.73 a	43.57 e
	75%	39.92 a	432.9 b	31.13 a	0.540 ab	-0.28 c	53.74 a	0.67 b	66.83 d
	50%	38.17 ab	399.1 b	30.81 ab	0.458 bc	0.89 c	53.19 ab	0.59 c	84.49 c
	25%	35.83 b	350 c	27 b	0.365 c	3.53 b	50.14 b	0.50 d	131 b
	0%	29.67 c	252 d	22.67 c	0.186 d	7.11 a	45.96 c	0.30 e	152 a
Variety	Torkman	35.8 bc	330.5 cd	27.93 b	0.36 b	3.07 a	52.99 a	0.57 b	94.98 ab
	Karoon × Kavir	37.2 ab	370 c	32.3 a	0.48 a	1.27 cd	51.32 ab	0.62 a	97.16 a
	Reihani	38.43 ab	324.6 d	33.68 a	0.46 a	0.47 d	53.48 a	0.58 b	96.62 a
	C-74-9	38.8 a	351.7 cd	33.63 a	0.49 a	0.6 d	51.13 ab	0.53 c	95.83 ab
	Gorgan-4	35.43 c	494.4 a	17.27 c	0.33 b	2.67 ab	50.74 ab	0.57 b	92.95 b
	Kavir × Badia	35.3 c	449.49 b	26.7 b	0.49 a	1.97 bc	49.69 b	0.52 c	95.95 ab
Irrigation × Variety	n.s	**	n.s	*	**	n.s	**	n.s	

*, ** Significant at 0.05 and 0.01, respectively

n.S Non significant

data were analysed by MSTATc, SAS and SPSS softwares and the Excel software was used for graphical descriptions.

RESULTS AND DISCUSSION

Grain Yield and Yield Components

An analysis of variance showed that water deficit significantly reduced the 1000-grain weight, number of spikes. m⁻², number of grains/spike and grain yield (Tables 1 and 2). The 1000-grain weight and its size reduction is possibly due to a decrease in the assimilation rate and lower photoassimilate translocation to physiological sinks. Water deficit is known to reduce the 1000-grain weight by shortening the grain-filling period [3, 9]. These findings are in agreement with Mohammad *et al.* [9] who found that water stress at the flowering stage reduced 1000-grain weight significantly. There were significant differences in the number of grains/spike between barley varieties (Table 2). However, differences were not significant between 50, 75 and 100% crop water requirements, suggesting that moderate wa-

ter deficit had no effect on the abortion of young florets and did not reduce the number of grains/spike. The number of grains/spike of the Gorgan-4 variety was the lowest due to its 2-rowed type (distichum) spike feature. Water deficit reduced the number of spikes m⁻² in barley varieties (Table 2). The average number of spikes m⁻² was 494 in Gorgan-4 which was related to its higher tillering capacity as 2-rowed barley and the lowest spikes m⁻², with an average of 324, was observed in the Reihani variety. The grain yield of barley varieties at different irrigation levels is shown in Tables 2 and 3. Grain yields in the Karoon × Kavir and Kavir × Badia varieties were the highest under well-irrigated condition, while the Karoon × Kavir and Reihani varieties had the highest grain yield under stress conditions (Table 3). Moffat *et al* [8] also showed that water stress at the reproductive growth stage reduced the grain yield in wheat significantly.

Canopy Temperature

Water deficit had a significant effect ($P \leq 0.01$) on the canopy temperature difference between canopy and ambient air (ΔT)

Table 3. Interaction between irrigation regime and variety effects on the performance of barley varieties.

Irrigation Regime	Variety	No. Spikes/m ² **	Grain yield (Kg/m ²)*	ΔT(°c) **	F _v /F _m **	
100%	Torkman	340	ghij	0.479 cdefgh	-3 mn	0.72 abc
	Karoon × Kavir	472	cdef	0.685 a	-1 klm	0.72 abc
	Reihani	374	fghi	0.578 abcde	-1 klm	0.69 abcd
	C-74-9	328	ghij	0.597 abcdef	-3 mn	0.72 abc
	Gorgan-4	680	a	0.507 bcdefg	-5 n	0.77 a
	Kavir × Badia	577	b	0.665 ab	-4.3 n	0.77 a
75%	Torkman	397	efgh	0.422 efghijk	3 efg	0.64 cde
	Karoon × Kavir	378	fgh	0.537 abcdef	0 hijk	0.68 bcd
	Reihani	369	ghi	0.603 abcd	-1 klm	0.62 def
	C-74-9	339	jghi	0.570 abcdef	-2.7 lmn	0.63 de
	Gorgan-4	580	b	0.441 defghij	-0.33ijkl	0.71 abc
	Kavir × Badia	534.7	bc	0.607 abc	-0.66 jklm	0.73 ab
50%	Torkman	357	ghi	0.370 ghijkl	2.3 fgh	0.57 efg
	Karoon × Kavir	359	ghi	0.478 cdefghi	1.6 ghij	0.62 de
	Reihani	343	ghij	0.512 bcdefg	-0.33 ijkl	0.54 gh
	C-74-9	307	hijkl	0.470 cdefghi	0 hijk	0.54 gh
	Gorgan-4	493	bcde	0.367 ghijkl	2 ghi	0.63 de
	Kavir × Badia	520	bcd	0.549 abcdef	-0.3 ijkl	0.62 def
25%	Torkman	342	ghij	0.331 hijklm	3 efg	0.49 hi
	Karoon × Kavir	341	ghij	0.408 ghijk	2.7 efg	0.57efg
	Reihani	318	ijk	0.362 ghijkl	3 efg	0.54 fgh
	C-74-9	267	ijkl	0.321 ijklmn	3 efg	0.44 ij
	Gorgan-4	430	defg	0.277 klmno	2 cde	0.53 gh
	Kavir × Badia	402	efgh	0.293 jklmn	4.5 def	0.41 j
0%	Torkman	250	jkl	0.188 mnop	8 a	0.41 j
	Karoon × Kavir	300	hijkl	0.301 jklmn	7 abcd	0.47 ij
	Reihani	219	kl	0.323 jklm	5 cde	0.44 j
	C-74-9	223	kl	0.170 nop	5.7 abcde	0.33 k
	Gorgan-4	307	hijkl	0.104 p	7.7 ab	0.29 k
	Kavir × Badia	213	l	0.132 op	7 abcd	0.31 k

*, ** Significant at 0.05 and 0.01, respectively

(Tables 2 and 3). Negative and positive ΔTs under well irrigated and water stress conditions are shown in Figure 1. Negative ΔTs indicated a cooler canopy and positive ΔTs indicated a warmer canopy. With regard to the heat absorbing nature of transpiration, the cooler canopy (negative ΔT) is equivalent to a higher transpiration rate and higher stomatal conductance, while the warmer canopy (positive ΔT) indicated a lower transpiration rate and higher stomatal resistance. A higher water pressure deficit induces higher

stomatal resistance and, so, a higher leaf surface temperature. The ΔT increment due to water stress has been reported in wheat [4].

ΔT values for Karoon × Kavir and Reihani varieties were both -1°C under well irrigated conditions (Table 3) which had the highest grain yield under drought conditions (0.3 and 0.32 Kg.m⁻² respectively). It seems that a considerable relationship could be observed between higher canopy temperature (less negative ΔT) under well irrigated conditions and higher grain yield under stress




Figure 1. Temperature difference between canopy and ambient air in barley varieties under different irrigation levels, on the basis of crop water requirements.

conditions. Golestani and Assad [4] also reported similar results. They proposed that the mean comparison of ΔT s could not be considered in drought tolerance screening of wheat varieties by itself, while the higher canopy temperature under well irrigated conditions could be exploited as a screening criteria for drought tolerance selection of different genotypes under non-stress conditions.

Chlorophyll Content

The irrigation regime and varieties of barley both had significant effects on chlorophyll content (SPAD value) (Table 1). Water deficit decreased the chlorophyll content (SPAD value) in barley varieties and its reduction was significant at lower irrigation rates (50.14 and 45.96 at 25% and lower CWR respectively). (Table 2) There were not any significant differences between varieties. Similar results has been reported by Yavad [13].

A positive and significant correlation between the photosynthesis rate and SPAD values has been reported [3]. A lower photosynthesis rate under moderate water stress has been attributed to lower stomatal con-

ductance although, under severe stress, stomatal resistance may accompanied by non stomatal (mesophyll) resistance [7].

It seems that the SPAD value measurement at the heading stage may not be a reliable indicator for screening the drought tolerant varieties, because the pre-anthesis photosynthesis and carbohydrate remobilization ability might have compensated for the current photosynthesis reduction at the grain filling stage. Plant height is an important morphological trait related to carbohydrate remobilization under stress conditions [2, 3]. The higher grain yield of the Karoon \times Kavir and Reihani varieties under stress conditions may be related to their taller architecture (58 and 59 cm respectively) and better remobilization performance rather than to current photosynthesis. Sink activity and the grain filling period are two major important factors in carbohydrate remobilization capacity too. Sink activity is the sink's ability to absorb and utilize carbohydrates on the basis of the sink's tissue weight. The grain filling period, along with the grain filling rate, determine final grain weight and the former is greatly shortened due to environmental stresses. However, the grain filling period was relatively longer in the Karoon \times Kavir

and Kavir × Badia varieties (45 and 44.5 days respectively) under drought stress which may have enhanced their grain yield.

Photochemical Efficiency of Photosystem II

Irrigation regimes, the varieties of barley and their interactions all had significant effects on photochemical efficiency (F_v/F_m). (Table 1) Irrigation levels had significant effects on all chlorophyll fluorescence parameters especially on the F_v and F_m values which were reduced and increased, respectively, with an increase in stress severity (data were not shown). Such an effect of water stress had been expressed in a significant reduction in the F_v/F_m values (Table 2). The main effects of variety on the F_v/F_m values were different at all irrigation levels and distinctly higher in Karoon × Kavir (0.62) (Table 2). The F_v/F_m values for Karoon × Kavir and Reihani varieties were highest under severe stress condition (0% CWR) (0.47 and 0.44, respectively) which had better yield performance under drought conditions (Table 3). It can be concluded that the adverse effect of water stress on F_v/F_m could be adjusted by the varietal difference between plant types.

It seems that higher photochemical efficiency of Karoon × Kavir played a major role in its drought tolerance. Reports had shown that wheat plants could acclimatise to high temperatures by using higher F_v/F_m retention during reproductive growth. Such an acclimatisation could be related to the protection capability of the Q_A complex or its protein regeneration capacity in chlorophyll-protein complexes [8] which had led to higher grain yield under stress condition [8].

Proline Content

An analysis of variance showed that irrigation levels had significant effects on the free proline content of barley varieties and a

lower significant difference ($P \leq 0.05$) was observed between the barley varieties (Table 1). Water deficit significantly increased the proline content of all barley varieties to a maximum $152 \mu\text{g}\cdot\text{gr}^{-1}$ Fw, but was at its minimum ($43.5 \mu\text{g}\cdot\text{gr}^{-1}$ Fw) under well-irrigated treatment (Table 2).

Irrigation regimes and varieties of barley both had significant effects on proline content and the irrigation effect was more distinct than that of the variety (Table 1). As regards to proline accumulation under environmental stresses and its role in osmotic adjustment (1, 4, 8 and 12), it seems that its accumulation procedure were more or less similar in all varieties. In comparison with the results of Lingose and Bergman [5], it may be suggested that the Karoon × Kavir and Reihani varieties were much more tolerant to drought stress with a higher grain yield at 0% CWR (0.31 and $0.323 \text{ Kg}\cdot\text{m}^{-2}$ respectively).

CONCLUSION

The present study showed that the higher canopy temperature (less negative ΔT) under well irrigated conditions and higher grain yield, 1000-grain weight and F_v/F_m under water stress conditions were the proper criteria (morphological or physiological markers) for screening the drought tolerant barley genotypes under field or laboratory conditions. Further experiments should be focused on screening for drought tolerance at the early stages of plant growth using such criteria in order to determine the relationship with field experiments and shortening the screening duration.

REFERENCES

1. Bates, I. S., Waldren, R. P., and Teare, I. D. 1973. Rapid Determination of Free Proline for Water Stress Studies. *Plant Soil*, **39**:205-207.
2. Blum, A., Shpiler, L., Golan, A. and Mayer, J. 1989. Yield Stability and Canopy Temperature of Wheat Genotypes under Drought



- Stress. *Field Crop Res.*, **22(41)**:289-286.
- Earl, H. and Tollenaar, M. 1977. Maize Leaf Absorption of Photosynthetically Active Radiation and its Estimation using Chlorophyll Meter. *Crop Sci.*, **37**:36-440.
 - Gloestani, S. and Assad, M. T. 1998. Evaluation of Four Screening Techniques for Drought Resistance and their Relationship to Yield Reduction in Wheat. *Euphytica*, **103**:293-299.
 - Lingose, V. and Bergman, H. 1995. Changes in the Yield Lignin Content and Protein Pattern of Barley Induced by Drought Stress. *Angewante- Botanik.*, **69(6)**: 206-210.
 - Liu, X. Z. and Zhang, L. G. 1994. Differences in Diurnal Changes in Canopy Temperature of Winter Wheat under Water Stress Condition. *Acta Agr.*, **20**: 229-232
 - Ma, B. L., Morison, M. J. and Videng, H. D. 1995. Leaf Greenness and Photosynthetic Rates in Soybean. *Crop Sci.*, **35**: 1411- 1414.
 - Moffatt, J., Sears, M. R. G. and Paulsen, G. M. 1990. Wheat High Temperature Tolerance During Reproductive Growth. I. Evaluation by Chlorophyll Fluorescence. *Crop Sci.*, **30**: 881- 885.
 - Mohammad, J., Aaziri, M., Nazir, A., Shah, D. and Jamal, H. 1996. Wheat Yield Components as Affected by Low Water Stress at Different Growth Stages. *Sarhad J. Agr.*, **12**:19- 26
 - Singh, T., Paleg, L. and Aspinall, D. 1973. Stress Metabolism Variations in Response to Water Deficit in the Barley Plants. *Cab. Abstrct.*, 1972-1975.
 - Turner, B. 1990. The Extent and Pattern of Osmotic Adjustment in Winter Clover During the Development of Water Stress. *Ann. Bot.*, **66**:721- 727.
 - Umedi, L. V. 1989. A Rapid and Nondestructive Method to Determine Response to Water Deficit in the Barley Plant. *Crop Sci.*, **35**:655- 650.
 - Yavad, U. 1989. A Rapid and Nondestructive Method to Determine Chlorophyll in Intact Leaves. *Hortscience*, **21(6)**: 1449- 1450.

اثرات کمبود آب بر عملکرد و خصوصیات فیزیولوژیکی ارقام جو

۱. ممنوعی، ر. فتوحی قزوینی، م. اصفهانی و ب. ناخدا

چکیده

اثرات کمبود آب بر عملکرد، اجزای عملکرد و خصوصیات فیزیولوژیکی ارقام جو طی آزمایشی در قالب طرح بلوک‌های خرد شده در سال ۷۸-۱۳۷۷ در مؤسسه تحقیقات اصلاح و تهیه نهال و بذر کرج مورد بررسی قرار گرفت. در این آزمایش ۵ سطح آبیاری (صفر، ۲۵، ۵۰، ۷۵ و ۱۰۰ درصد نیاز آبی گیاه) در کرت‌های اصلی و ۶ رقم جو، کارون × کویر، ریحانی (مقاوم به خشکی)، ترکمن، ۹-۷۴- C (نیمه مقاوم)، کویر × بادپاو گرگان ۴ (جو دو ردیفه حساس به خشکی) در کرت‌های فرعی مشخص شدند. نتایج نشان دادند که کمبود آب باعث القای تنش به گیاه و کاهش عملکرد و اجزای عملکرد آن گردید. کمبود آب باعث افزایش دمای پوشش گیاهی گردید و ارقامی که در شرایط بدون تنش دارای پوشش گیاهی گرمتری بودند در شرایط کمبود آب مقاومت بهتری نسبت به خشکی از خود نشان دادند. تنش شدید باعث کاهش چشمگیر میزان کلروفیل (مقادیر SPAD) در ارقام جو گردید، لیکن میزان این

کاهش در تیمارهای ۵۰، ۷۵ و ۱۰۰ درصد آبیاری معنی‌دار نبود. بعلاوه در شرایط تنش آب تغییرات میزان کلروفیل در ارقام مختلف جو معنی‌دار نبود. اثرات آبیاری بر راندمان فتوشیمیایی فتوسیستم ۲ (F_v/F_m) و سایر اجزای فلئورسانس کلیه ارقام جو معنی‌دار بود. اگر چه مقادیر F_v/F_m ارقام جو در هر سطح آبیاری معنی‌دار نبود، لیکن در مجموع میزان فلئورسانس در ارقام کارون \times کویر و ریحانی در شرایط تنش آبی کمتر از سایر ارقام بود. سطوح آبیاری باعث تغییرات معنی‌داری در میزان اسید آمینه پرولین انباشته شده در ارقام جو گردید، لیکن میزان پرولین انباشته شده در ارقام مختلف جو در سطوح مختلف آبیاری تفاوت معنی‌داری نداشت. در مجموع به نظر می‌رسد که در شرایط آبیاری کامل، بالاتر بودن دمای پوشش گیاهی ($\Delta T =$ تفاوت دمای بین پوشش گیاهی و اتمسفر، کمتر منفی) و در شرایط تنش آبی، بالاتر بودن عملکرد دانه، وزن هزار دانه و F_v/F_m از مؤثرترین شاخص‌های شناسایی ژنوتیپ‌های مقاوم در جو در شرایط آزمایشگاهی و مزرعه‌ای محسوب می‌شوند.