

## Rice Bran Phytosterols of Three Widespread Iranian Cultivars

M. Derakhshan-Honarparvar<sup>1</sup>, M. M. Hamed<sup>2\*</sup>, and M. Kh. Pirouzifard<sup>1</sup>

### ABSTRACT

Rice bran sterols have antioxidant activity as well as physiological and biological effects. Because of these benefits, the aim of this study is evaluating the quantity and quality of Iranian rice bran sterols. Three widespread Iranian rice cultivars (Khazar, Hashemi, and Alikazemi) were used for determination of their sterol contents. Two methods of sample preparation were compared and the better one was used in this work. Rice bran samples were saponified directly after acid hydrolysis. Unsaponified materials were extracted, purified by solid phase extraction, silylated, and their sterol fractions determined by GC-MS. The sterol composition (in mg kg<sup>-1</sup> bran) of three cultivars (Khazar, Alikazemi, and Hashemi) were 1,330.69, 1,279.95, 1,313.17  $\beta$ -sitosterol; 747.52, 696.05, 756.8 campesterol; 112.8, 115.36, 114.24  $\Delta$ -5-avenasterol, 38.912, 33.08, 38.24  $\Delta$ -7-avenasterol; 8.05, 7.07, 7.56c cholesterol; 4.20, 3.99, 4.23 brassicasterol; and 2,722.016, 2,706.176, 2,717.68 total sterols, respectively. The results showed that there were no significant differences between the individual and total sterol contents of these three cultivars ( $P > 0.05$ ), except campesterol,  $\beta$ -sitosterol, and  $\Delta$ -7-avenasterol. The highest and lowest sterols were  $\beta$ -sitosterol and campesterol.

**Keywords:** Phytosterols, Plant sterols, Rice bran, Sterol content determination.

### INTRODUCTION

Phytosterols (Ps) are biosynthetically derived from squalen and belong to the group of triterpenes that includes more than 4,000 different compounds. Phytosterols are made up of a tetracyclic cyclopenta ( $\alpha$ ) phenanthrene ring and a flexible side chain at C17. The most common Ps -sitosterol, campesterol and stigmasterol- are structurally closely related to cholesterol. In plants, phytosterols exist as free and conjugated (Loharson, 2004).

Sterol in rice-bran oil contain considerable quantities of  $\Delta$ -5-avenasterol and related sterols. As with common vegetable oils,  $\beta$ -sitosterol is the major sterol in rice oil. More than 75% of the sterols of rice bran-oil are esterified and are collectively called

orizanol. These sterol esters of ferulic acid show antioxidant activity as well as physiological and biological effects (Gunstone, 2002). Vegetables and fruits as well as cereals are suitable sources of plant sterols in diet. However, the quantity of phytosterols in a regular diet is not sufficient for these health effects (Gunstone, 2002). Therefore, they are extracted and added to some foodstuffs for enrichment.

Rice bran is a byproduct in Iran which is used primarily as domestic animal foodstuff. So, in this work we used the bran of three widespread cultivars as a cheap source for the evaluation of their sterol content. We used the solid phase extraction (SPE) method which is more rapid and convenient than TLC (Thin Layer Chromatography) for cleaning up unsaponifiable matters

<sup>1</sup> Department of Food Science and Technology, College of Agriculture, University of Urmia, Urmia, Islamic Republic of Iran.

<sup>2</sup> Department of Food Science and Engineering, Faculty of Biosystem, University of Tehran, Karaj, Islamic Republic of Iran.

\* Corresponding author, e-mail: mmhamed@ut.ac.ir



(Azadmard-Damirchi, 2006, Pironen and Toivo, 1999; Toivo, 2000)

## MATERIALS AND METHODS

The bran of three widespread cultivars of Iranian rice (*Oryza sativa*) named Khazar, Hashemi, and Alikazemi were taken from the Rice Research Center in Rasht, Iran. The samples after milling (Cyclotec 1093 sample mill, Tekator, Sweden) and sieving (20 meshe) were packaged in nylon bags and kept frozen until analysis. The average oil content of the bran was 16%.

### Saponification

Two methods were used for saponification of plant materials for phytosterol extraction which had been developed by Toivo (1999) and Pironen (2001). Table 1 shows the fatty acid profile of extracted unsaponifiable matters by two methods. So, we used the Piironen method which showed lower residual fatty acids in unsaponifiable fraction. Acid hydrolysis can hydrolyze glycoside sterol esters in cereals (Toivo, 2000), so we used it as a pretreatment of saponification in cereals.

One gram of each sample was weighed in a 50 ml flask and betulin (Sigma-Aldrich) was added as an internal standard. For acid hydrolysis, 5 ml of 6M HCl was added to the flask and the contents were mixed for 10 seconds then refluxed for 60 minutes on a water bath. The content of the flask was

mixed manually every 10 minutes, then the flask was cooled to the room temperature, and its contents were transferred to a screw capped test tube. Lipids were extracted with 20 ml of hexane and diethyl ether (1:1) solvent mixture by shaking for 10 minutes. The organic layer was separated by letting the tube stand for 15 minutes (and centrifuging if necessary), and was transferred to a round-bottom flask and evaporated to dryness in a rotary evaporator at 50°C. For saponification, 8 ml of absolute ethanol and 0.5 ml of saturated aqueous KOH solution were added. The flask contents were mixed (10 seconds) and refluxed at 80-85°C for 30 minutes, then cooled to room temperature, and transferred into a screw capped test tube; 12 ml of water and 20 ml of cyclohexane were added, followed by shaking for 10 minutes. An aliquot of 15 ml of the organic layer was transferred to a round-bottom flask and evaporated to dryness in a rotary evaporator at 50°C. The residue was dissolved in 1 ml of hexane (Pironen and Toivo, 1999).

A SiOH cartridge (chromo bond SiOH, 6 ml, 1 gr) was activated by 5 ml of hexane. The sample solution was first filtered through a nylon acrodisc (13 Pall Perman Laboratory, Ann Arbor, and MI). The solution was eluted by gravity flow for several minutes and then by vacuum (5-10 Hg) for 5 minutes. The sterol fraction was eluted with 5ml of hexane and diethyl ether (90:10) solvent and evaporated to dryness at 50°C. The sterol fraction residue was dissolved in 500 µl of dichloromethane.

**Table 1.** Fatty acid residual of two methods of saponification.

Fatty acid residue	Toivo Method (%)	Piironen Method (%)
Lauric acid	0.15	1.89
Myristic acid	1.97	1.97
Palmitic acid	16.8	2.8
Margaric acid	0.6	1.6
Oleic acid	18.17	4.87
Linoleic acid	24.59	7.99
Stearic acid	6.53	5.53
Total fatty acid	44.22	23.76

### Identification and Quantification

Sterols were analyzed on GC by their trimethylsilyl (TMS) ether derivatives. In a presanitized sample vial, the solvent was evaporated under nitrogen (50°C) and then 100 µl of anhydrous pyridine (E. Merck, Darmstadt, Germany) and 100 µl of derivatization reagent containing 99% of BSTFA (E. Merck, Darmstadt, Germany) and 1% of TMCS (Flucka Chemie AG, Switzerland) were added. To complete the silylation, solutions were allowed to stand overnight at room temperature before GC analysis (Pironen and Toivo 1999).

GC-MS analysis was used to identify the major sterols in sample matrix. MS analysis was also used to evaluate peak purities of sterols.

A Hewlett-Packard (HP, Palo Alto, USA) HP 6890 series GC equipped with a split/splitless injector and a HP 5973 mass-selective detector system were used. The MS was operated in the EI mode (70 eV). Helium (99.999%) was employed as carrier gas and its flow rate was adjusted to 1 ml min<sup>-1</sup>. The separation was performed on a 60 m×0.25 mm I.D. fused silica capillary column of HP-1. The GC column temperature was programmed at 100°C and raised to 270°C at 30°C min<sup>-1</sup>, 90 minutes hold at this temperature. The injector temperature was set at 260°C, and all the injection was carried out on the splitless mode. The ion source and quadrupole

temperature were set at 230°C and 150°C, respectively.

The area of each peak was measured relative to the internal standard peak area and, then, the concentration of each sterols was calculated.

### Analysis of Results

All the analysis was performed in triplicate. SPSS was used for calculation of the statistical analysis.

## RESULTS AND DISCUSSION

The sterol contents of the three cultivars are given in Table 2. Table 3 shows there is not significant difference ( $P > 0.05$ ) between their sterol and individuals except campesterol,  $\Delta$ -7 avenasterol and  $\beta$ -sitosterol. The decreasing order of sterols was sitosterol, campesterol,  $\Delta$ -5 avenasterol,  $\Delta$ -7 avenasterol, cholesterol, and brassicasterol. Because the rice bran sterols reported by others (19, 20) were in mgkg<sup>-1</sup> oil, for comparison we calculated our results in mgkg<sup>-1</sup> oil. Table 4 demonstrates that the total contents of Iranian cultivars are higher than the sterols reported by Firestone (1999) and lower than those of Rossel, (2001).

**Table 2.** Sterol content of rice bran and calculated rice bran oil.

Sterol	Khazar		ALIKAZEMI		HASHEMI	
	mg/kg oil*	Sterol Fraction Percentage	mg/kg oil*	Sterol Fraction Percentage	mg/kg oil*	Sterol Fraction Percentage
Cholesterol	50.3±2.1	0.3	44.16±3.1	0.26	47.23±0.5	0.28
Campesterol	4672±11.0	27.8	4350.3±47.5	25.72	4730±19.9	27.84
Sitosterol	8129.3±706.3	47.8	7999.7±108.1	47.30	8207.33±7.4	48.3
Brassicasterol	26.3±1.0	0.15	24.93±0.9	0.15	26.45±0.7	0.16
$\Delta$ -5-Avenasterol	705±6.0	4.14	720.97±5.9	4.26	714±2.0	4.20
$\Delta$ -7-Avenasterol	243.2±2.1	1.43	206.77±1.7	1.22	239±5.3	1.41
Others	3185±1.9	18.72	3566.77±101.1	21.11	3021.43±2.1	17.88
TOTAL	17012.6±11.3		16913.6±118.0		16985.5±30.2	

**Table 3.** Statistics assessment of sterols content.

Source*	SS	d f	MS	F
1 SS <sub>M</sub>	57.0422	2	28.5211	6.21
SS <sub>P</sub>	27.9600	6	4.6600	
2 SS <sub>M</sub>	251,322.6667	2	125,666.3333	135.7251**
SS <sub>P</sub>	5,555.3333	6	925.8889	
3 SS <sub>M</sub>	66022.89	2	33011.44	7.96**
SS <sub>P</sub>	24882.00	6	4,147.000	
4 SS <sub>M</sub>	4.2654	2	2.1327	2.9925
SS <sub>P</sub>	4.2760	6	0.7127	
5 SS <sub>M</sub>	384.4689	2	192.2344	7.6159
SS <sub>P</sub>	151.4467	6	25.2411	
6 SS <sub>M</sub>	12,386.7267	2	1,193.3633	101.0564**
SS <sub>P</sub>	70.8533	6	11.8089	
7 SS <sub>M</sub>	15,681.7267	2	7,840.8633	1.5715
SS <sub>P</sub>	29,935.7533	6	4,989.2922	

\* 1=Cholesterol, 2= campesterol, 3=  $\beta$ -Sitosterol, 4=Brassicasterol, 5= $\Delta$ -5-avenasterol, 6=  $\Delta$ -7-avenasterol, and 7=Total sterol.

\*\* Significant difference ( $P < 0.05$ ).

**Table 4.** Rice bran oil Sterol composition % reported by others.

	Rossel(2001)	Firestone(1999)
Cholesterol	2.2	—
Brassicasterol	0.3	—
Campesterol	20.4	24
Stigmasterol	10.8	11.5
Sitosterol	34.8	51.5
$\Delta$ -5-avenasterol	3.2	8
$\Delta$ -7-Stigmasterol	11.6	1.5
$\Delta$ -7-avenasterol	2.4	3
Total(mg/kg)	36758	10550

## CONCLUSIONS

The average of total sterol contents of Khazar, Alikazemi and Hashemi cultivars were in the range of 2,706.176-2,722.016 mg kg<sup>-1</sup> bran or 16,913-1,712.06 mg kg<sup>-1</sup> oil,  $\beta$ - sitosterol was in the highest and brassica in the lowest proportions.

The result showed that there was not any significant differences ( $P > 0.05$ ) between the total sterol content of the three cultivars, and between individual sterols except campesterol, brassicaesterol, and  $\Delta$ -7 avenasterol.

## ACKNOWLEDGEMENTS

We thank Department of Food Science and Engineering, Faculty of Biosystem, University of Tehran for friendly corpora ting helps to do this study.

## REFERENCES

1. ISO, Animal and Vegetable Fats and Oil Determination of Individual and Total Sterols Content-gas Chromatographic Method. International standard, ISO 12228.

2. Azadmard-Damirchi, A. and Dutta, P. C. 2006. Novel Solid-phase Extraction Method to Separate 4-desmethyl-4-monomethyl-, and 4,4'-dimethylsterols In Vegetable Oils. *J. Chromatography*, **1108(A)**: 183-187
3. Bailey, A. E. 1996. Rice Bran Oil. *Oil and Fat Products*, **2**: 393-409
4. Burn, P. and Kishor, G. M. 2000. Food as a Source of Health Enhancing Compounds. *J. AgBIOFORM*, **3(1)**: 3-9
5. Firestone, D. 1999. *of Oils, Fats and Waxes*. AOCS Technical Services Champaign, IL.
6. Gordon, D. T. 1982. Sterols in Mollusks and Crustaceans of the Pacific Northwest. *J. AOCS*, **59(12)**
7. Gunstone, F. 2002. *Vegetable Oil in Food Technology*. CRC Press.
8. *Identity and Purity*. Standard 1.3.4, Australia New Zealand Food Standards, PP. 7.
9. Lagarda, M. J., Garcia-Llatas, G. and Farre, R. 2006. A Review of Analysis of Phytosterols in Foods. *J. Pharmaceutical and Biomedical Analysis*, **41**:1486-1496
10. Lampi, A., Moreau, R., Piironen, V. and Hicks, K. 2004. Pearling Barley and Rye to Produce Phytosterol-rich Fraction. *Lipids, AOCS Press*, **39(8)**: 783-787
11. Loharson, J. 2004. Phytosterol Oxidation Products. Doctoral thesis, Department of Food Science, Swedish University of Agricultural Sciences, Upsala.
12. Nagao, T., Kobayashi, T., Hirota, Y., Kitano, M., Kishimoto, T., Fujita, T., Watanabe, Y. and Shimada, Y. 2005. Improvement of a Process for Purification Soybean Oil Deodorizer Distillate. *Molecular Catalysis: Enzymatic*, **1(6)**: 56-62
13. European Union, 2000; 2004. Phytosterols and Their Esters for Use as Ingredient in a Range of Food Applications. European Commission, 500/EC and 335/EC.
14. European Union, 2003. *Opinion of the Scientific Committee on Food on an Application from Multibene for Approval of Plant Sterol Enriched Foods*. European Commission, 15 April 2003, Scientific Committee on Food, SCF/ CS/ NF/ DOS/24 ADD 2 Final.
15. European Union, 2000. Opinion of the Scientific Committee On Food on a Request for the Safety Assessment of the Use of Phytosterol Esters in Yellow Fat Spreads. European Commission, 6 April 2000, SCF/ CS/ NF/ DOS/1 Final.
16. VERIS, 2003. Plant Sterols Esters/Sterols May Reduce the Risk of Heart. Disease. VERIS Research Information Service, November 2003.
17. Piironen, V., Toivo, J. and Lampi, A. 1999. Plant Sterols in Cereals Products. *J. Cereal Chemistry*, **79(1)**: 148-158
18. Phytosterol Esters Derived From, (August 2005). First Review Report, Application A433, A434, A508, Australia/New Zealand Food Standards, 5-05, 3.
19. *Plant Sterols and Stanols*. A Document Adopted from [www.heartfoundation.com.au](http://www.heartfoundation.com.au)
20. Rossell, J. B. 2001. *Frying Improving Quality*. CRC PRESS.
21. Toivo, J., Lampi, A., Aalto, S. and Piironen, V. 2000. Factors Affecting Sample Preparation in the Gas Chromatographic Determination of Plant Sterols in Whole Wheat Flour. *Food Chemistry*, **68**: 239-245
22. Taguchi, G. 1986. *Introduction to Quality Engineering*. Dearborn, MI: American Supplier Institute.
23. Verleyen, T., Verhe, R., Garcia, L., Dewettinck, K., Huyghebaert, A. and Greyt, D. 2001. Gas Chromatographic Characterization of Vegetable Oil Deodorization Distillate, *J. Chromatography*, **921(A)**: 277-285
24. Weber, N., Weitkamp, P. and Mukherjee, K. D. 2002. Cholesterol Lowering Food Additives: Lipase-catalysed Preparation of Phytosterol and Phytostanol Esters. *Food Research International*, **35**: 177-181



## تعیین محتوی فیتوسترول سه واریته برنج ایرانی

م. درخشان هنرپرور، م. م. حامدی و م. خ. پیروزی فرد

### چکیده

محتوی استرول سه واریته پرمصرف برنج ایرانی (خزر، هاشمی و علی کاظمی) اندازه گیری گردید. سبوس ها بعد از هیدرولیز اسیدی و قلیایی، صابونی شدند. مواد غیر قابل صابونی استخراج گردیدند، به روش SPE خالص سازی شدند، سیلیله شدند و قسمت استرولی آنها توسط دستگاه GC-MS اندازه گیری شد. محتوی استرولی سه واریته به ترتیب نزولی عبارت بودند از: بتا سیوسترول (۱۳۳۰/۶۹، ۱۲۷۹/۹۵، ۱۳۱۳/۱۷ به ترتیب در مورد واریته های خزر، علی کاظمی، و هاشمی) برحسب میلی گرم در کیلوگرم سبوس، کمپسترول (۷۴۷/۵۲، ۶۹۶/۰۵، ۷۵۶/۸۰ به ترتیب در مورد واریته های خزر، علی کاظمی، و هاشمی) برحسب میلی گرم در کیلوگرم سبوس،  $\Delta$ -۵-آوناسترول (۱۱۲/۸، ۱۱۵/۳۶، ۱۱۴/۲۴ به ترتیب در مورد واریته های خزر، علی کاظمی، و هاشمی) برحسب میلی گرم در کیلوگرم سبوس،  $\Delta$ -۷-آوناسترول (۳۸/۹۱۲، ۳۳/۰۸، ۳۸/۲۴ به ترتیب در مورد واریته های خزر، علی کاظمی، و هاشمی) برحسب میلی گرم در کیلوگرم سبوس، کلسترول (۸/۰۵، ۷/۰۷، ۷/۵۶ به ترتیب در مورد واریته های خزر، علی کاظمی، و هاشمی) برحسب میلی گرم در کیلوگرم سبوس، براسیکاسترول (۴/۲۰، ۳/۹۹، ۴/۲۳ به ترتیب در مورد واریته های خزر، علی کاظمی، و هاشمی) برحسب میلی گرم در کیلوگرم سبوس، و استرول تام (۲۷۲۲/۰۱۶، ۲۷۰۶/۱۷۶، ۲۷۱۷/۶۸ به ترتیب در مورد واریته های خزر، علی کاظمی، و هاشمی) برحسب میلی گرم در کیلوگرم سبوس. نتایج نشان داد که تفاوت قابل ملاحظه آماری بین مقادیر استرول تام در سه واریته وجود نداشت و در بین اجزاء استرولی به جز کمپسترول و  $\Delta$ -۷-آوناسترول تفاوت قابل ملاحظه آماری وجود نداشت.