

Functional Surface Display of Laccase in a Phenol-Inducible Bacterial Circuit for Bioremediation Purposes

Mehrnoosh Fathi-Roudsari^{*1}, Mehrdad Behmanesh^{*2}, Ali-Hatef Salmanian¹, Majid Sadeghizadeh² and Khosro Khajeh³

¹National Institute of Genetic Engineering and Biotechnology, Tehran, Iran; ²Department of Genetics, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran; ³Department of Biochemistry, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran

Received 11 July 2017; revised 1 August 2017; accepted 6 August 2017

ABSTRACT

Background: Phenolic compounds, which are produced routinely by industrial and urban activities, possess dangers to live organisms and environment. Laccases are oxidoreductase enzymes with the ability of remediating a wide variety of phenolic compounds to more benign molecules. The purpose of the present research is surface display of a laccase enzyme with adhesin involved in diffuse adhesion (AIDA-I) autotransporter system on the surface of *Escherichia coli* cells for bioremediation of phenolic compounds. **Methods:** The expression of laccase was regulated by a phenol-responsive promoter (a σ^{54} promoter). The constitutively-expressed CapR transcription activator was able to induce laccase expression in the presence of phenolic compounds. **Results:** Western blot analysis showed the expression and correct transfer of the enzyme to the outer membrane of *E. coli* cells in the presence of phenol. Activity assay confirmed the correct folding of the enzyme after translocation through the autotransporter system. HPLC analysis of residual phenol in culture medium showed a significant reduction of phenol concentration in the presence of cells displaying laccase on the surface. **Conclusion:** Our findings confirm that autodisplay enables functional surface display of laccase for direct substrate-enzyme availability by overcoming membrane hindrance. **DOI: 10.22034/ibj.22.3.202**

Keywords: Laccase, Phenol, Bioremediation, AIDA-I

Corresponding Authors: Mehrnoosh Fathi-Roudsari and Mehrdad Behmanesh

National Institute of Genetic Engineering and Biotechnology, Pajoohesh BLVD, 17 km Karaj Highway, Tehran, Iran; Tel.: (+98-21) 44787401; Fax: (+98-21) 44787399; E-mail: mfathi@nigeb.ac.ir

Department of Genetics, Faculty of Biological Sciences, Tarbiat Modares University, P.O. Box 14115-154, Tehran, Iran; Tel.: (+98-21) 82884451; Fax: (+98-21) 82884717; E-mail: behmanesh@modares.ac.ir

INTRODUCTION

Human activities have resulted in the release of a plethora of aromatic and organic chemicals into the environment^[1]. Phenols and their halogenated derivatives, poly-aromatic hydrocarbons and pesticides, are classes of toxic environmental pollutants that are suspected for carcinogenicity, teratogenicity, and genotoxicity effects^[2,3]. Compared to traditional physico-chemical methods, bioremediation is generally the safest, least disruptive and most cost-effective treatment for the removal of these

toxic compounds. Biological tools such as whole microorganisms or isolated enzymes are able to degrade persistent contaminants into non- or less-toxic compounds^[4-7].

Laccases are a diverse family of cuproprotein enzymes that oxidize a broad range of aromatic compounds, including phenols, orthodiphenols, p-dihydroxybenzenes, aminophenols, polycyclic aromatic hydrocarbons, and aromatic polyamines^[8]. The active site of these enzymes contain a total of four copper ions arranged as a type 1 blue copper site and type 2 and type 3 copper ions that forms a trinuclear

cluster for O₂ binding and activation^[9]. Laccases catalyze oxidation of the phenolic compounds with concomitant reduction of oxygen to water^[10]. Due to wide reaction capabilities and broad substrate specificity, these enzymes have great biotechnological potentials, including bioremediation, xenobiotic degradation, polymer synthesis, textile dye bleaching, pulp, and paper bleaching as well as biosensor applications^[11,12].

Previously, we have prepared a genetically-modified *Escherichia coli* capable of expressing a laccase enzyme in the presence of common phenolic pollutions^[13]. A transcription activator (CapR) was constitutively expressed inside the cells but triggered the expression of the laccase gene when exposed to phenolic pollutions^[13,14]. It has been shown that upon the presence of phenolic contamination, laccase is able to be expressed and accumulated inside the cells^[13]. However, a general problem in using whole microorganisms for bioremediation purposes is bio-availability. Intracellular presence of the enzymes restricts the accessibility towards substrates, therefore decreases the efficiency of bioremediation.

Transfer of the biocatalyst to the cell surface will result in direct contact of the enzyme and substrates without the need for the contaminants to cross membranes^[5,15-20]. Among different ways of transferring enzymes to the outer membrane of Gram-negative bacteria, an *E. coli* virulence factor, adhesin involved in diffuse adhesion (AIDA-I), has been widely used. Naturally, the N-terminal signal sequence of AIDA-I is followed by a passenger domain that is transported across the cell envelope through a translocator domain in the C-terminal^[21-24]. Here, we have fused the translocator domain of AIDA-I to the C-terminal of the locally-isolated thermostable laccase in our CapR-regulated inducible system. Therefore in the environments with phenolic contaminations, the aromatic molecules that pass through the membrane will result in the expression of chimeric laccase-AIDA gene through activation of CapR. Laccase translocation to the outer cell membrane can increase the exposure of enzyme-substrate and facilitates the detoxification procedure.

Successful display of laccase on the surface of *Bacillus subtilis* spores, *Pseudomonas putida*, and *Saccharomyces cerevisiae* has previously been reported, but there is no report on the surface display of laccase in *E. coli*^[18,9,25-27]. Here, we have shown that autodisplay can be used for surface expression of the laccase enzyme, leading to development of a whole cell biocatalyst. Laccase displayed on the surface of the cells showed activity towards its synthetic and natural substrates such as syringaldazine (SGZ) and phenol.

MATERIALS AND METHODS

Chemicals and reagents

Taq DNA polymerase, high fidelity PCR enzyme mix, restriction endonucleases, and Ins/TA clone PCR cloning kit were obtained from Fermentas (Germany). SGZ and 3,3'-diaminobenzidine (DAB) were purchased from Sigma (St. Louis, MO, USA). All other chemicals were from Merck (Darmstadt, Germany) with reagent grade. Antibodies were prepared from Biovision (USA).

DNA manipulation

The previously reported pCap-lac plasmid was purified and digested with *Bam*HI and *Sma*I enzymes^[13]. In this construct, a laccase gene, which had formerly been isolated from a local *Bacillus* species (GenBank accession no. FJ663050), was placed under the control of a phenol-inducible Po promoter^[13]. Subsequently, AIDA-I transfer unit (TU) region (linker and β -barrel) was amplified using pAng-AIDA plasmid (a generous gift from Dr. Michael Mourez) with forward and reverse primers containing *Bg*III and *Sma*I sites, respectively. The sequences of the primers are presented below:

AIDAF: AGATCTCTTAATCCTACAAAAGAAAGT

AIDAR: CCC GGTTAG AAGCTGTATTTTATCCC

The amplified fragment was cloned in pTZ57R/T and digested with the mentioned enzymes. Then the TU was joined in frame with laccase gene in the pCap-lac plasmid. The final construct was named pCap-lac-TU, and its authenticity was confirmed by sequencing.

Growth condition and expression of laccase in the presence of inducer

Both pCap-lac and pCap-lac-TU were transformed into chemically prepared *E. coli* BL21 competent cells, and the recombinant clones were selected on Lauria-Bertani (LB) medium containing 100 μ g/mL of ampicillin. Five mL of LB medium was inoculated with single colonies of selected transformants and incubated at 37 °C overnight. Subsequently, 50 mL fresh medium was inoculated with the overnight preculture and incubated at 37 °C until an OD₆₀₀ of 0.5 was reached. The induction was carried out with the addition of 100 nm phenol at 37 °C for 4 hours^[13].

Cell fractionation and outer membrane preparation

First, 1 mL of the overnight culture was inoculated into 20 mL fresh medium and grown until reached an OD₆₀₀ of 0.5-0.6. Then phenol was added, and expression was carried out for four hours. After harvesting and washing with 0.2 M Tris-HCl, pH 8,

cells were sonicated for total protein analysis, or differential cell fractionation was performed^[28,29]. The protein accessibility test was also carried out in order to find whether laccase is exposed to the surface side of the outer membrane or trapped to the periplasmic side. After expression of laccase, cells were harvested and resuspended in PBS containing proteinase K (50 µg/mL). Treatment was carried out at 37 °C for 1 hour. The reaction was stopped after three times washing of the cells with PBS containing 5% FBS. The fractionation was carried out to prepare the outer membrane of the treated cells^[30-33].

SDS-PAGE and Western blot

Total cell lysate, outer membrane proteins, and the periplasmic/cytoplasmic fractions of the cells were diluted twofold with the SDS-PAGE sample buffer, loaded on 12.5% SDS-PAGE and blotted to PVDF membrane. A monoclonal anti-His-tag was applied in 1% BSA- Tris-buffered saline, 0.1% Tween 20 at 4 °C overnight. A horseradish peroxidase-conjugated goat anti-mouse polyclonal antibody was applied to the membrane with 1:2500 dilution for one hour to develop the signals.

Whole cell activity assay

Whole cell activity experiment was carried out through the incubation of the whole cells containing pCap-lac or pCap-lac-TU with SGZ, as a spectrophotometric substrate. After the induction of expression with phenol for four hours, cells were harvested and resuspended in PBS at 4 °C overnight. The cells were then harvested and incubated with 0.05 mM SGZ at 37 °C. After 1 hour, cells were sedimented by centrifugation, and the absorbance of the supernatant was measured at 525 nM.

Effect of laccase on phenol conversion and HPLC analysis

Fresh medium (20 mL) was inoculated with pre-cultured cells, and phenol (100 nM) was added as the inducer after reaching the suitable turbidity of 0.7-0.8. Cells were incubated with phenol for an additional period of 24 hours. Medium without any cells was used as the control in order to find spontaneous conversion of phenol during 24 hours. After this period, cells were harvested, and 800 µl of each medium was used for isolation of organic compounds. Afterwards, 200 µl trichloroacetic acid was mixed with 800 µl medium and stored on ice for 10 minutes. Then the tubes were centrifuged at 12,000 ×g at 4 °C for 15 min. Supernatant was transferred to new tubes and mixed three times with the same volume of trichloromethane. Lower phase containing trichloromethane and all

organic extracts of the medium was transferred to new tubes for evaporation of the solvent. The pellet was applied for the HPLC experiment using C18 reverse phase column. Methanol-water with the percentage of 50:50 was used as the mobile phase with the speed of 1 mL/min.

RESULTS

Construction of an inducible laccase-displaying circuit in *E. coli*

For preparing the pCap-lac-TU construct, the TU of AIDA-I (linker-β-barre) was joined to the C terminal of laccase through compatible ends of *Bam*HI and *Bg*II. Figure 1 shows the schematic view of the pCap-lac and the pCap-lac-TU final constructs. The recombinant plasmids were transformed into competent *E. coli* BL21 cells and screened on solid media containing 100 µg/ml ampicillin. The reason for choosing *E. coli* BL21, as the host of expression, was the absence of outer membrane protease T (*ompT*) on the external surface of the bacteria. Since the linker region of the AIDA-I contains a cleavage site for *ompT*, here, we have used *E. coli* BL21 as the host.

Expression and surface display of laccase

Fusing laccase enzyme (with the molecular mass of 65 kDa) to the TU of AIDA-I results in the formation of a fusion protein with the molecular mass of 106 kDa after the cleavage of signal sequence. Following induction of expression with 100 nM phenol, a band with the molecular mass of 106 kDa was observed in total cell lysate. After cell fractionation, the same band was obvious in the outer membrane fraction (Fig. 2A). Following the induction of expression with 100 nM phenol and cell fractionation, Western-blot analysis showed a band with the correct size, just in the outer membrane fraction of the cells. The band was absent in cytoplasmic/periplasmic fraction, which shows the efficient transfer of the fusion protein to the outer membrane (Fig. 2). Proteinase K treatment was also performed to find the direction of laccase attachment to the outer membrane. Since the proteinase K molecules are unable to pass the outer membrane, degradation of the fusion protein supports the correct transfer of the enzyme to the outer side of the cells. Trapping the laccase in the periplasmic side can protect the fusion protein from degradation. As shown in Figure 2, the band corresponding to the fusion protein was disappeared after proteinase treatment, which proves the surface display of laccase.

It has previously been shown that the formation of disulfide bonds and complete folding of the passenger




Fig. 1. Schematic view of the constructs expressing laccase and laccase-TU fusion protein. (A) The map of the pCap-lac construct. CapR transcription activator is expressed constitutively by the Pr promoter and ends in a T7 terminator. Divergently, Po inducible promoter can be occupied by CapR and results in transcription of the periplasmic signal peptide (pelB), His-tag, and laccase gene in the presence of phenols. The laccase expression is terminated by the T7 terminator. (B) The map of the pCap-lac-TU construct. The TU region corresponding to the linker and β -barrel domain of AIDA-I is joined to the C-terminal of the laccase gene. Upon activation of the Po promoter by CapR, laccase can be expressed in fusion with TU and transferred through the β -barrel to the surface of the outer membrane.

proteins may result in unsuccessful transfer of the protein through β -barrel to outer membrane^[34]. Laccase enzymes contain a disulfide bond between cysteine residues 229 and 322 in crystallographic studies^[35]. Although Western-blot analysis of outer membrane fraction with or without proteinase treatment confirmed correct transfer of laccase to the surface, we also prepared the outer membrane fraction from cells, which were grown in highly reducing medium containing 10 mM of 2-mercaptoethanol. Western blot analysis showed no difference in the transfer of fusion protein in normal or reducing conditions (data not shown). It seems that the presence of disulfide bond in laccase structure cannot hinder the transfer of the enzyme through outer membrane.

Cell surface activity assay of laccase

To investigate the biological activity of the laccase molecules, which were transferred to the outer membrane, whole cells were assayed in the presence of SGZ. After the expression of laccase in the presence of phenol as inducer and dialysis against PBS, cells were incubated with SGZ substrate for one hour. The absorbance of the supernatant at 525 nm showed significantly higher oxidative ability of cells, which harbor laccase-AIDA fusion. Cells containing pCap-lac express laccase inside the cells and show the activity of 0.2 μ M/hour in comparison to 2 μ M/hour for cells containing pCap-lac-TU. These results prove almost 10 times higher oxidative ability for cells, which display laccase on the surface.

The biological activity of the displayed laccase was additionally assayed in the presence of phenol as a natural substrate. In the present expression system, phenol not only is an inducer of the laccase expression but also acts as a substrate. After four-hour induction

of laccase expression with phenol (100 nM), cells were kept at 37 °C for an additional 24 hours. Medium containing 100 nm phenol was kept at 37 °C for 24 hours, as the control. After harvesting the cells, the organic fractions of the medium were extracted. Figure 3A shows the organic fraction of medium containing cells, which express laccase inside the cells (pCap-lac) or display laccase on the surface (pCap-lac-TU). Obviously, the sample prepared from surface-displayed laccase contained remarkable higher amounts of brown precipitates. These precipitates are known as phenolic polymers formed as the result of oxidation reaction. The organic fraction obtained from pCap-lac containing cells showed much lower amounts of brown precipitates. After centrifugation, the supernatant was analyzed by reverse-phase HPLC to find any residual soluble phenol in the medium. The obtained peaks are shown in Figure 3B. The medium containing 100 nm phenol for 24 hours, without any cells, was used as the control. A single peak with retention time of approximately two minutes was observed. Phenol extracts from medium containing BL21 cells harboring pCap-lac show a reduction in the height of the mentioned peak. Looking at the graph obtained from cells displaying laccase, a remarkable reduction of phenol molecules can be seen. The height and surface area of the peak corresponding to the phenol shows almost four times reduction in comparison to the control.

DISCUSSION

Laccases (E.C. 1.10.3.2, *p*-benzenediol:oxygen oxidoreductase) are copper-containing oxidoreductase enzymes able to oxidize a variety of aromatic




Fig. 2. (A) Protein separation using SDS-PAGE analysis. Lane 1, the molecular weight marker; lanes 2 and 3, *E. coli* cell lysate containing pCap-lac-TU before and after induction with phenol, respectively; lane 4, outer membrane fraction after phenol induction. The chimeric protein with the mass of 106 kDa is obvious after phenol induction. (B) Western blot on cells harboring pCap-lac-TU. The cells were fractioned before and after proteinase K treatment. Lane 1, cytoplasmic/periplasmic soluble proteins; lane 2, outer membrane fraction; lane 3, protein weight marker; lane 4, outer membrane fraction after proteinase K treatment. The band of the fusion protein is present in the outer membrane fraction. After proteinase treatment, the band is disappeared, confirming the correct translocation of the laccase through TU to the outer surface of the cells.

compounds, including mono-, di-, and poly-phenols^[36,37]. These enzymes can oxidize, polymerize, degrade, or transform phenolic compounds to less toxic derivatives^[38,39]. An interesting characteristic of laccases is their low specificity towards substrates. A diverse and broad range of chemicals, including phenols, dyes, pesticides, endocrine disrupters, and polycyclic aromatic hydrocarbons can be converted by the laccase active site to more benign compounds^[40,41]. Therefore, a wide range of applications from food stabilization, dye bleaching, and textile preparation to bioremediation of wastewaters can be catalyzed using these enzymes^[4,42].

Although laccases are considered as relatively stable enzymes, prevention of inactivation under industrial conditions still remains a priority. Tolerance of the

enzyme towards temperature, pH, organic solvents, and free radicals, which might be produced as the result of laccase function, is a highly important criterion in industrial use of this enzyme. Immobilization by various chemicals or physical methods can increase the stability and tolerance of enzymes to denaturation in harsh industrial situations and extend its functional life-time^[5,38,39]. Enzymes can also be immobilized on the surface of live organisms through linking to one of the cell wall or outer membrane molecules. Surface display has other benefits such as direct contact of enzyme with substrates and elimination of the expensive steps of enzyme purification^[43].

From different strategies of surface display in Gram-negative *E. coli*, we have used the autotransporter system. Functional transport of several enzymes to the surface of *E. coli* has been reported frequently^[43-45]. Here, we have substituted the passenger domain of




Fig. 3. Conversion of phenol and HPLC analysis of residual phenol. (A) The organic fraction of culture medium after 24 hours of phenol (100 nM) treatment. Brown precipitates are the oxidized polymeric forms of phenol remarkably higher in medium of laccase-displaying cells. (B) The supernatant of organic fractions were injected to HPLC. Chromatograms show the amount of residual phenols. (a) Medium without any cells shows a single peak with retention time of 2 minutes. (b) Medium which contained cells with pCap-lac plasmid. (c) Medium contained surface-display laccase cells due to the presence of pCap-lac-TU plasmid. The peak shows an obvious reduction in phenol after treatment with cells displaying laccase.

AIDA-I with laccase (laccase-TU) and prepared a pelB signal sequence at the beginning of the coding region. We have also used the inducible pathway of phenol degradation in *Pseudomonas putida* KCTC1452; therefore, laccase expression is controlled by the CapR transcription activator and σ^{54} -RNA polymerase that recognizes the -12 (GC)/24 (GG) promoter^[11].

The present study indicated that in the presence of phenol, laccase was expressed from pCap-lac-TU and successfully transferred to the cell surface. Western blot analyses proved the absence of the chimeric protein in the cytoplasmic fraction of the phenol-treated cells and showed a band with the correct size of chimeric protein in the outer membrane fraction. Proteinase K treatment of whole cells and subsequent disappearing of the laccase-TU band confirmed the correct direction and exposure of the chimeric protein to the outer membrane.

Similar to other immobilization methods, chimerization may also interfere with catalytic properties of the displayed enzymes. Although correct transfer of laccase to the cell surface was confirmed, the functionality of the enzyme was investigated through activity assays. Treatment of the whole cell biocatalysts with SGZ confirmed the remarkable higher oxidative ability of laccase-displaying cells for conversion of this substrate. Additionally, long treatment of cells harboring pCap-lac and pCap-lac-TU with phenol as a natural substrate showed the high oxidative ability of laccase, which was transferred to the surface. Brown polymers, which are the result of phenol polymerization, were remarkably higher in medium containing cells with surface-displayed laccase. The polymerized precipitates were also observed in medium with intracellular-expressing laccase. The negligible amount of phenol polymerization was also observed in medium without cells, as the negative control, showing the spontaneous polymerization of phenol in the condition of experiment. In agreement to these data, the residual phenol of the culture media was also analyzed using reverse phase HPLC. HPLC analysis of phenol moieties from medium containing BL21 cells with pCap-lac plasmid showed a smaller peak in comparison to the control (medium without cells). It is assumed that phenols can be partially degraded even when laccase is expressed inside the cells. Laccase release, as the result of cell lysis or transfer of a fraction of phenol molecules through the membrane, might be the reason of phenol degradation in this experiment. Presence of the laccase on the surface of *E. coli* cells dramatically reduced the amount of residual phenol in the medium, which indicates the functional transfer of the enzyme and increased

bioavailability of substrates.

We conclude that laccase, as the remediating enzyme, is expressed and successfully transferred to the outer surface of *E. coli* in our phenol inducible expression system. The enzyme is capable of passing through the β -barrel of AIDA-I TU. The present study confirms that the displayed enzyme is functional and active on the surface of bacteria and shows the ability to polymerize phenol as the substrate. The cells displaying laccase on the surface can be used as whole-cell biocatalysts.

ACKNOWLEDGMENTS

Our great thanks goes to Dr. Michael Mourez for the kind gift of plasmid containing AIDA-I autotransporter gene. The authors also thank Tarbiat Modares University (Tehran, Iran) for the financial support of the work.

CONFLICT OF INTEREST. None declared.

REFERENCES

1. Tropel D, Van der Meer JR. Bacterial transcriptional regulators for degradation pathways of aromatic compounds. *Microbiology and molecular biology reviews* 2004; **68**(3): 474-500.
2. Juhasz AL, Naidu R. Bioremediation of high molecular weight polycyclic aromatic hydrocarbons: a review of the microbial degradation of benzo(a)pyrene. *International biodeterioration and biodegradation* 2000; **45**(1-2): 57-88.
3. Leedjarv A, Ivask A, Virta M, Kahru A. Analysis of bioavailable phenols from natural samples by recombinant luminescent bacterial sensors. *Chemosphere* 2006; **64**(11): 1910-1919.
4. Ang EL, Zhao H, Obbard JP. Recent advances in the bioremediation of persistent organic pollutants via biomolecular engineering. *Enzyme and microbial technology* 2005; **37**(5): 487-496.
5. Alcalde M, Ferrer M, Francisco JP, Ballestros A. Environmental biocatalysis: from remediation with enzymes to novel green processes. *Trends biotechnology* 2006; **24**(6): 281-287.
6. Paul D, Pandey G, Pandey J, Jain RK. Accessing microbial diversity for bioremediation and environmental restoration. *Trends biotechnology* 2005; **23**(3): 135-142.
7. Sutherland TD, Home I, Weir KM, Coppin CW, Williams MR, Selleck M, Russell RJ, Oakeshott JG. Enzymatic bioremediation: from enzyme discovery to applications. *Clinical and experimental pharmacology and physiology* 2004; **31**(11): 817-821.
8. Wang W, Zhang Z, Ni H, Yang H, Li Q, Li L.

- Decolorization of industrial synthetic dyes using engineered *Pseudomonas putida* cells with surface-immobilized bacterial laccase. *Microbial cell factories* 2012; **11**: 1-14.
9. Cho EA, Seo J, Lee DW, Pan JG. Decolorization of indigo carmine by laccase displayed on *Bacillus subtilis* spores. *Enzyme and microbial technology* 2011; **49**(1): 100-104.
 10. Durán N, Rosa MA, D'annibale A, Gianfreda L. Applications of laccases and tyrosinases (phenoloxidases) immobilized on different supports: a review. *Enzyme and microbial technology* 2002; **31**(7): 907-931.
 11. Sharma P, Goel R, Capalash N. Bacterial laccases. *World journal of microbiology and biotechnology* 2007; **23**(6): 823-832.
 12. Viswanath B, Rajesh B, Janardhan A, Kumar AP, Narasimha G. Fungal laccases and their applications in bioremediation. *Enzyme research* 2014; 163242.
 13. Fathi-Roudsari M, Behmanesh M, Salmanian AH, Sadeghizadeh M, Khajeh K. Substrate-dependent expression of laccase in genetically modified *Escherichia coli*: design and construction of an inducible phenol-degrading system. *Preparative biochemistry and biotechnology* 2013; **43**(5): 456-467.
 14. Park SM, Park HH, Lim WK, Shin HJ. A new variant activator involved in the degradation of phenolic compounds from a strain of *Pseudomonas putida*. *Journal of biotechnology* 2003; **103**(3): 227-236.
 15. Saleem M, Brim H, Hussain S, Arshad M, Leigh MB, Hassan Z. Perspectives on microbial cell surface display in bioremediation. *Biotechnology advances* 2008; **26**(2): 151-161.
 16. Samuelson P, Gunneriusson E, Nygren PA, Stahl S. Display of proteins on bacteria. *Journal of biotechnology* 2002; **96**(2): 129-154.
 17. Chen W, Georgiou G. Cell-surface display of heterologous proteins: from high-throughput screening to environmental applications. *Biotechnology and bioengineering* 2002; **79**(5): 496-503.
 18. Van Bloois E, Winter RT, Kolmar H, Fraaije MW. Decorating microbes: surface display of proteins on *Escherichia coli*. *Trends in biotechnology* 2011; **29**(2): 79-86.
 19. Lee SY, Choi JH, Xu Z. Microbial cell-surface display. *Trends biotechnology* 2003; **21**(1): 45-52.
 20. Yang Z, Liu Q, Wang Q, Zhang Y. Novel bacterial surface display systems based on outer membrane anchoring elements from the marine bacterium *Vibrio anguillarum*. *Applied and environmental microbiology* 2008; **74**(14): 4359-4365.
 21. Rutherford N, Mourez M. Surface display of proteins by Gram-negative bacterial autotransporters. *Microbial cell factories* 2006; **5**: 22-37.
 22. Maurer M, Jose J, Meyer TF. Autodisplay: one-component system for efficient surface display and release of soluble recombinant proteins from *Escherichia coli*. *Journal of bacteriology* 1997; **179**(3): 794-804.
 23. Jose J. Autodisplay: efficient bacterial surface display of recombinant proteins. *Applied microbiology and biotechnology* 2006; **69**(6): 607-614.
 24. Fang Y, Ngeleka M, Middelton DM, Simko E. Characterization and immuno-detection of AIDA-I adhesin isolated from porcine *Escherichia coli*. *Veterinary microbiology* 2005; **109**(1-2): 65-73.
 25. Gupta N, Farinas ET. Directed evolution of CotA laccase for increased substrate specificity using *Bacillus subtilis* spores. *Protein engineering, design and selection* 2010; **23**(8): 679-682.
 26. Lu J, Guo Q, Cui M, Yang L, Du S, Ruan H, He G. Construction of a yeast cell-surface display system and expression of *trametes* sp. laccase. *Advanced materials research* 2012; **347**: 3635-3640.
 27. Bleve G, Lezzi C, Spagnolo S, Rampino, P, Perrotta C, Mita G, Grieco F. Construction of a laccase chimerical gene: recombinant protein characterization and gene expression via yeast surface display. *Applied biochemistry and biotechnology* 2014; **172**(6): 2916-2931.
 28. Konieczny MP, Benz H, Hollinderbaumer B, Beinke C, Niederweis M, Schmidt MA. Modular organization of the AIDA autotransporter translocator: The N-terminal β 1-domain is surface-exposed and stabilizes the transmembrane β 2-domain. *Antonie van leeuwenhoek* 2001; **80**: 19-34.
 29. Hantke K. Regulation of ferric iron transport in *Escherichia coli* K12: isolation of a constitutive mutant. *Molecular and general genetics* 1981; **182**(2): 288-292.
 30. Jose J, Bernhardt R, Hannemann F. Cellular surface display of dimeric Adx and whole cell P450-mediated steroid synthesis on *E. coli*. *Journal of biotechnology* 2002; **95**(3): 257-268.
 31. Klauser T, Pohlner J, Meyer TF. Extracellular transport of cholera toxin B subunit using *Neisseria* IgA protease beta-domain: conformation-dependent outer membrane translocation. *EMBO journal* 1990; **9**(6): 1991-1999.
 32. Schultheiss E, Weiss S, Winterer E, Maas R, Heinzle E, Jose J. Esterase autodisplay: enzyme engineering and whole-cell activity determination in microplates with pH sensors. *Applied and environmental microbiology* 2008; **74**(15): 4782-4791.
 33. Schultheiss E, Paar C, Schwab H, Jose J. Functional esterase surface display by the autotransporter pathway in *Escherichia coli*. *Journal of molecular catalysis B: enzymatic* 2002; **18**(1-3): 89-97.
 34. Jose J, Zangen D. Autodisplay of the protease inhibitor aprotinin in *Escherichia coli*. *Biochemical and biophysical research communication* 2005; **333**(4): 1218-1226.
 35. Enguita FJ, Martins LO, Henriques AO, Carrondo MA. Crystal structure of a bacterial endospore coat component: A laccase with enhanced thermostability properties. *The journal of biological chemistry* 2003; **278**(21): 19416-19425.
 36. Karam J, Nicell JA. Potential applications of enzymes in waste treatment. *Journal of chemical technology and biotechnology* 1999; **69**(2): 141-153.
 37. Thurston CF. The structure and function of fungal laccases. *Microbiology* 1994; **140**: 19-26.

- [DOI: 10.22034/ibj.22.3.202]
38. Madhavi V, Lele SS. Laccase: Properties and applications. *Bioresources* 2009; **4**(4): 1694-1717.
 39. Majeau JA, Brar SK, Tyagi RD. Laccases for removal of recalcitrant and emerging pollutants. *Bioresource technology* 2010; **101**(7): 2331-50.
 40. Morozova OV, Shumakovich GP, Gorbacheva MA, Shleev SV, Yaropolov AI. Blue laccases. *Biochemistry* 2007; **72**(10): 1136-1150.
 41. Rodgers CJ, Blanford CF, Giddens SR, Skamnioti P, Armstrong FA, Gurr SJ. Designer laccases: a vogue for high-potential fungal enzymes. *Trends biotechnology* 2010; **28**(2): 63-72.
 42. Durán N, Esposito E. Potential applications of oxidative enzymes and phenoxidase-like compounds in wastewater and soil treatment: a review. *Applied catalysis B: environmental* 2008; **28**(2): 83-99.
 43. Schumacher SD, Hannemann F, Teese MG, Bernhardt R. Autodisplay of functional CYP106A2 in *Escherichia coli*. *Journal of biotechnology* 2012; **161**(2): 104-112.
 44. Schumacher SD, Jose J. Expression of active human P450 3A4 on the cell surface of *Escherichia coli* by autodisplay. *Journal of biotechnology* 2012; **161**(2): 113-20.
 45. Jose J, Bernhardt R, Hannemann F. Functional display of active bovine adrenodoxin on the surface of *E. coli* by chemical incorporation of the (2Fe-2S) cluster. *Chembiochem* 2001; **2**(9): 695-701.