Brief Report

Protective Effect of Safranal against Gentamicin-Induced Nephrotoxicity in Rat

Mohammad Taher Boroushaki, Hamid Reza Sadeghnia

Abstract

Gentamicin is an important aminoglycoside antibiotic. However its use is limited to serious and life threatening gram negative infections, because of its high nephrotoxicity potential in patients. There are reports that safranal, the active ingredient of saffron with antioxidant properties, exerts protective effect against ischemic injuries occurred by certain nephrotoxins including gentamicin. Therefore, in the present study, we examined the protective effect of safranal against gentamicin-induced nephrotoxicity in rat. After acclimatization, animals were randomly divided to three groups (8 rats /each group). On day one, each animal was placed separately in a metabolic cage for collecting 24-hour urine samples. On day two, after collecting urine samples for measuring glucose and protein, the rats in group 1 received saline 1 ml/kg for 6 days, those in group 2 received gentamicin 80 mg/kg/day for 6 days, and the remaining rats in group 3 received safranal 0.5 ml/kg followed by gentamicin 80 mg/kg/day for 6 days. Injections were intraperitoneally. All the animals were euthanized 24 hours after the last dose. Blood samples were collected by cardiac puncture, and concentration of blood urea, creatinine, and urinary glucose and protein, as the indicators of nephrotoxicity were measured. Our results showed that in group 2, concentration of blood urea nitrogen p<0.01, creatinine p<0.05, urinary glucose p<0.001, and protein p<0.01 were significantly increased compared with the control and safranal-treated groups. There was no significant difference between the control and safranal-treated groups. Safranal exerts protective effects against gentamicin-induced nephrotoxicity in rat.

Iran J Med Sci 2009; 34(4): 285-288.

Keywords • Safranal • gentamicin • nephrotoxicity

Introduction

entamicin is an important aminoglycoside antibiotic. Its use is limited to serious and life threatening gram negative infections because of its high nephrotoxicity potential by inhibiting the protein synthesis in renal cells in 13-30% of treated patients. This mechanism specifically causes necrosis of cells in the proximal tubule, resulting in acute tubular necrosis which can lead to acute renal failure.¹

Crocus sativus L. (Iridaceae), commonly known as saffron is used in folk medicine for various purposes such as an aphrodisiac, antispasmodic, and expectorant.² Modern pharmacological studies have demonstrated that saffron extracts exert antitumor,³⁵ free radical scavenger, hypolipemic,⁶ and anticonvulsant activities,⁷ as well as improving learning and memory,⁶⁸

Department of Pharmacology and Pharmacological, Research Center of Medicinal Plants, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

Correspondence:

Mohammad Taher Boroushaki PhD, Department of Pharmacology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran. **Tel:** +98 511 8828566 **Fax:** +98 511 8828567 **Email:** <u>boroushakimt@mums.ac.ir</u> Received: 28 December 2008 Revised: 8 March 2009 Accepted: 23 April 2009

Archive of SID

M.T. Boroushaki, H.R. Sadeghnia

Phytochemical studies on *Crocus sativus* have shown the presence of constituents such as crocin, crocetin, safranal, and picrocrocin.⁹⁻¹¹ Among constituents of saffron extract, crocetin is mainly responsible for these pharmacological activities.⁶

In traditional medicine, the stigma of this plant is used as an antiedematogenic remedy.² In addition some studies indicated that saffron has antioxidant activity.¹²⁻¹⁹ An investigation showed that aqueous extract of saffron could inhibit cisplatin, cyclophosphamide, mitomycin-C, and urethane induced alterations in lipid peroxidation.¹⁶ Also our laboratory findings showed that safranal was able to protect kidney against hexachlorobutadiene-induced nephrotoxicity and ischemia/reperfusion injury in rat.^{17,18} Therefore, it was decided that validation of the protective effect of safranal against gentamicininduced nephrotoxicity also be investigated.

Materials and Methods

Gentamicin was purchased from (Daru-Pakhsh Pharmaceutical Company, Tehran, Iran) and safranal was obtained from (Fluka Chemie, Switzerland).

In the present study, Wistar albino rats weighing 150-200g of either sex, provided by Animal Breeding Unit, Department of Pharmacology, School of Medicine, Mashhad, Iran, were used. The animals were kept under light/dark cycle with 20 °C and 50% humidity for 12 hours. After acclimatization, the animals were divided randomly into three groups (8 rats / each group), and placed in metabolic cages separately for collecting 24-hour urine samples.

After collecting the first urine samples, the rats in group 1 received saline 1 ml/kg for 6 days, those in group 2 received gentamicin 80mg/kg/day for 6 days,¹⁹ and the remaining rats in group 3 received safranal 0.5ml/kg¹⁷ plus gentamicin 80 mg/kg/day after 1 hour for 6 days. Urine samples were collected daily. All injections were intraperitoneal.

Twenty-four hours after the last injection, urine samples were collected for measuring glucose and protein, using enzymatic method glucose oxidase that catalyses the oxidation of glucose to gluconic acid, and turbidimetry (trichloroacetic acid),²⁰ respectively. Then the animals were euthanized under ether anesthesia. Blood samples were collected by cardiac puncture for measuring urea and creatinine as an indicator of kidney damage, using urease (a nickel-metallo enzyme that catalyzes the degradation of urea to ammonia and carbon dioxide),²¹ and Jaffé (the combined use of creatinine, amidohydrolase, and alkaline sodium picrate) methods, respectively.²²

Statistical Analysis

Data were expressed as mean ± standard error of mean (SEM). The differences among treated groups were analyzed by one-way ANOVA followed by Tukey test. P<0.05 was considered statistically significant.

Results

Concentrations of urea, creatinine, glucose, and protein, as indicators of kidney damage, are shown in table-1. Concentration of urea in gentamycin-treated group was significantly higher than the control and safranal-treated groups (P<0.001). Concentration of creatinine, in gentamycin- treated group was also higher (P<0.05) than the control and safranal-treated groups. Concentration of urinary glucose in gentamycin- treated group was significantly higher than the control (P<0.001) and safranaltreated groups (P<0.01). And concentration of urinary protein in gentamycin- treated group was significantly higher than the control (P<0.01) and safranal-treated groups (P<0.05). There was no significant difference between the control and safranal-treated groups.

Discussion

Our findings indicated that safranal, the active constituent of crocus sativus L. is able to protect kidneys against gentamycin-induced nephrotoxicity in rats. Concentration of blood urea, creatinine, and urinary glucose and protein were used as indicators of damage to kidney.

It has been shown that aqueous extract of saffron inhibits oxidative stress caused by cisplatin, cyclophosphamide, and mitomycin-C.¹⁶ Saffron also increases the reduced intracellular glutathione and enzymes including glutathione reductase and glutathione-S-transferase.²³

Table 1: Concentration of blood urea, creatinine, and urinary glucose and protein in different animal groups in the last day of study

Animal group	Blood urea (mg/dl)	Blood creatinine (mg/dl)	Urinary glucose (mg/dl)	Urinary protein (mg/dl)
Control	33.75 ± 1.6***	0.53 ± 0.02*	7 ± 0.71***	1.68 ± 0.09**
Gentamicin	85 ± 7.6	1.95 ± 0.49	15.75 ± 0.85	3.83 ± 0.17
Safranal + gentamicin	30.95 ± 3.7***	0.59 ± 0.08*	9.88 ± 0.89**	2.45 ± 0.35*

* P<0.05 **P<0.01 ***P<0.001 compare with gentamicin-treated group

There are other investigations on pharmacological effect of safranal including the effect on extracellular hippocampal levels of glutamate and aspartate,²⁴ inhibition of pentylenetetrazolinduced seizures,²⁵ and radical scavenging activity.²⁶ These studies indicated that safranal was a potent antioxidant and able to protect body organs against certain toxic materials.

In conclusion, findings of the present study show that safranal is a protective agent against gentamicin-induced nephrotoxicity in rat. However, the exact protective mechanism(s) of safranal is unknown and need to be more investigated.

Acknowledgement

This study was financially supported by the Vice-Chancellor for Research, Mashhad University of Medical Sciences, Mashhad, Iran.

Conflict of Interest: None declared

References

- 1 Sundin DP, Sandoval R, Molitoris BA: Gentamicin inhibits renal protein and phospholipid metabolism in rats: implications involving intracellular trafficking. *J Am Soc Nephrol* 2001; 12: 114-23.
- Zargari A. Medicinal Plants. 4th ed. Tehran, Iran: Tehran University Press, 1990. p. 574-8.
- 3 Nair SC, Pannikar B, Panikkar KR. Antitumour activity of saffron (Crocus sativus). *Cancer Lett* 1991; 57: 109-14.
- 4 Nair SC, Kurumboor SK, Hasegawa JH. Saffron chemoprevention in biology and medicine: a review. *Cancer Biother* 1995; 10: 257-64.
- 5 Salomi MJ, Nair SC, Panikkar KR. Inhibitory effects of Nigella sativa and saffron (Crocus sativus) on chemical carcinogenesis in mice. *Nutr Cancer* 1991; 16: 67-72.
- 6 Abe K, Saito H. Effects of saffron extract and its constituent crocin on learning behaviour and long-term potentiation. *Phytother Res* 2000; 14: 149-52.
- 7 Hosseinzadeh H, Khosravan V. Anticonvulsant Effects of Aqueous Andethanolic Extracts of *Crocus Sativus* L. Stigmas in Mice. *Arch Irn Med* 2002; 5: 44-47.
- 8 Zhang Y, Shoyama Y, Sugiura M, Saito H. Effects of Crocus sativus L. on the ethanolinduced impairment of passive avoidance performances in mice. *Biol. Pharm. Bull* 1994; 17: 217-21.
- 9 Tarantilis PA, Tsoupras G, Polissiou M. Determination of saffron (Crocus sativus

L.) components in crude plant extract using high-performance liquid chromatography-UV-visible photodiode-array detectionmass spectrometry. *J Chromatogr A* 1995; 699: 107-18.

- 10 Escribano J, Alonso GL, Coca-Prados M, Fernandez JA. Crocin, safranal and picrocrocin from saffron (Crocus sativus L.) inhibit the growth of human cancer cells in vitro. *Cancer Lett* 1996; 100: 23-30.
- 11 Lozano P, Delgado D, Gómez D, et al. A non-destructive method to determine the safranal content of saffron (Crocus sativus L.) by supercritical carbon dioxide extraction combined with high-performance liquid chromatography and gas chromatography. *J Biochem Biophys Methods* 2000; 43: 367-78.
- 12 Ochiai T, Ohno S, Soeda S, et al. Crocin prevents the death of rat pheochromyctoma (PC-12) cells by its antioxidant effects stronger than those of alphatocopherol. *Neurosci Lett* 2004; 1362: 61-4.
- 13 Martínez-Tomé M, Jiménez AM, Ruggieri S, et al. Antioxidant properties of Mediterranean spices compared with common food additives. *J Food Prot* 2001; 64: 1412-9.
- 14 Pham TQ, Cormier F, Farnworth E, et al. Antioxidant properties of crocin from Gardenia jasminoides Ellis and study of the reactions of crocin with linoleic acid and crocin with oxygen. J Agric Food Chem 2000; 48: 1455-61.
- 15 Verma SK, Bordia A. Antioxidant property of Saffron in man. *Indian J Med Sci* 1998; 52: 205-7.
- 16 Premkumar K, Abraham SK, Santhiya ST, Ramesh A. Protective effects of saffron (Crocus sativus Linn.) on genotoxinsinduced oxidative stress in Swiss albino mice. *Phytother Res* 2003; 17: 614-7.
- 17 Boroushaki M.T, Mofidpour H, Sadeghnia H.R. Protective Effect of Safranal against Hexachlorobutadiene-Induced Nephrotoxicity in Rat *Irn J of Med Sci* 2007: 32: 173-6.
- 18 Sadeghnia HR, Boroushaki MT, Mofidpour H. Effect of Safranal, A Constituent of Saffron (Crocus Sativus L.), On Lipid Peroxidation Level during Renal Ischemia-Reperfusion Injury in Rats. *Iranian J. of basic Medical Sciences* 2005; 8: 179-85.
- 19 Ali BH, Al-Qarawi AA, Mousa HM. The effect of calcium load and the calcium channel blocker verapamil on gentamicin nephrotoxicity in rats. *Food Chem Toxicol* 2002: 40, 1843-7.

Iran J Med Sci December 2009; Vol 34 No 4 287

Archive of SID M.T. Boroushaki, H.R. Sadeghnia

- 20 Burtis CA, Ashwood ER, Bruns DE. Textbook of clinical chemistry and molecular diagnostics. 4th ed. Washington, Sanders, 2006. p. 215.
- 21 Guilbault GG, Montalvo JG Jr. A ureaspecific enzyme electrode. J Am Chem Soc 1969: 91: 2164-5
- 22 Masson P, Ohlsson P, Björkhem I. Combined enzymic-Jaffé method for determination of creatinine in serum. Clin Chem 1981; 27: 18-21.
- 23 Nair SC, Salomi MJ, Varghese CD, et al. Effect of saffron on thymocyte proliferation, intracellular glutathione levels and its antitumor activity. Biofactors 1992; 4: 51-4.
- 24 Hosseinzadeh H, Sadeghnia HR, Rahimi A. Effect of safranal on extracellular hippocampal levels of glutamate and aspartate during kainic Acid treatment in anesthetized rats. Planta Med. 2008, 74: 1441-5.
- 25 Hosseinzadeh H, Sadeghnia HR. Protective effect of safranal on pentylenetetrazolinduced seizures in the rat: involvement of GABAergic and opioids systems. Phytomedicine 2007; 14: 256-62.
- 26 Assimopoulou AN, Sinakos Z, Papageorgiou VP. Radical scavenging activity of Crocus sativus L. extract and its bioactive constituents. Phytother Res 2005; 19: 997-1000.