A SIMPLE TRANSFORMATION METHOD IN SKEWNESS REDUCTION

B. Abbasi

Department of Mathematical and Geospatial Sciences, RMIT University Victoria, Australia b.abbasi@gmail.com

S.T. Akhavan Niaki* and S.E. Seyedan

Department of Industrial Engineering, Sharif University of Technology P.O. Box 11155-9414, Tehran, Iran niaki@sharif.edu - amirsdn@gmail.com

*Corresponding Author

(Received: May 16, 2009 – Accepted in Revised Form: June 21, 2011)

Abstract Statistical analysis of non-normal data is usually more complicated than that for normal distribution. In this paper, a simple root/power transformation technique developed by Niaki, et al [1] is extended to transform right and left skewed distributions to nearly normal. The value of the root/power is explored such that the skewness of the transformed data becomes almost zero with an acceptable error. The proposed method is then compared to the well-known and complicated Box, et al [2] transformation method for different left and right skewed distributions using Monte Carlo simulation. While the proposed procedure is easy to understand and to implement, the results of the simulation study show that it works as good as the Box-Cox method.

Keywords Root/Power Transformation, Box-Cox Transformation, Non-Normal Data Analysis

چکیده روشهای تحلیل آماری داده های غیر نرمال معمولا" پیچیده تر از روشهای تحلیل داده های نرمال است. در این مقاله، یک روش سادهٔ تبدیل توانی برای توزیعهای احتمالی که به سمت راست و چپ چولگی دارند توسعه داده می شود تا آنها را به توزیع نرمال نزدیک کند. مقدار عددی توان با استفاده از روش عددی برش نیم طوری جستجو و به دست می آید که چولگی موجود در داده ها را تقریبا" از بین ببرد. عملکرد روش پیشنهادی آنگاه با عملکرد روش معروف و پیچیدهٔ باکس -کاکس برای توزیعهای غیر نرمال متفاوت با چولگیهای چپ و راست با استفاده از شبیه سازی مقایسه می شود. نتایج مقایسه نشان می دهد گه بر خلاف سادگی، روش پیشنهادی دست کم به خوبی روش باکس و کاکس عمل می کند.

1. INTRODUCTION

The statisticians George Box, et al [2] developed a procedure to identify an appropriate exponent (λ) to use to transform data (Y) into a "normal shape." The λ value indicates the power to which all data should be raised. In order to do this, the Box-Cox power transformation searches for λ until the best value is found. Note that for $\lambda = 0$, the transformation is not Y⁰ (because this would be 1 for every value), but instead the logarithm of *Y*.

The Box-Cox power transformation is not a guarantee for normality. This is because it actually

does not really check for normality. The assumption is that amongst all transformations with λ values between -5 and +5, transformed data has the highest likelihood-but not a guarantee-to be normally distributed when standard deviation is the smallest. Therefore, it is absolutely necessary to always check for the normality of the transformed data using a probability plot for example.

The original form of the Box-Cox transformation takes the following form:

$$Y(1) = \begin{cases} \frac{Y^{-1} - 1}{l} ; & \text{if } l \neq 0\\ Log(Y) ; & \text{if } l = 0 \end{cases}$$
(1)

Vol. 24, No. 2, June 2011 - 169

IJE Transactions A: Basics

Additionally, the Box-Cox power transformation only works for positive-valued data. This, however, can usually be achieved easily by adding a constant (C) to all negative-valued data such that they all become positive before transformation. The transformation equation is then:

$$Y'(1) = \begin{cases} \frac{(Y+C)^{l}-1}{l} ; & \text{if } l \neq 0\\ Log(Y+C) ; & \text{if } l = 0 \end{cases}$$
(2)

As an extension to the Box-Cox power transformation technique, Manly [3] proposed the following exponential transformation:

$$Y(1) = \begin{cases} \frac{e^{IY} - 1}{I} ; & \text{if } I \neq 0 \\ Y ; & \text{if } I = 0 \end{cases}$$
(3)

In which Y can take negative values. This transformation was reported to be successful in transforming uni-modal skewed distribution into normal distribution. However, its performance is not good for bimodal or U-shaped distributions [4].

John, et al [5] proposed the "modulus transformation" as a modification to the Box-Cox power transformation method in which Y can take negative values. The modulus transformation takes the form

$$Y(I) = \begin{cases} Sign(Y) \frac{(|Y|+1)^{l} - 1}{l}; & \text{if } I \neq 0\\ Sign(Y) Log((|Y|+1)+1); & \text{if } I = 0 \end{cases}$$
(4)

Where

$$Sign(Y) = \begin{cases} 1 & ; if \ Y \ge 0 \\ -1 & ; if \ Y < 0 \end{cases}$$
(5)

This transformation works best for distributions that are somewhat symmetric. We note that a power transformation on a symmetric distribution is likely going to introduce some degree of skewness.

Bickel, et al [6] gave the following slight modification in their examination of the asymptotic performance of the parameters in the Box-Cox transformations model:

$$Y(1) = \frac{|Y|^{l} Sign(Y) - 1}{l} ; for l > 0$$
 (6)

Yeo, et al [7] made a case for the following transformation:

$$Y(1) = \begin{cases} \frac{(Y+1)^{l}-1}{l} ; & \text{if } l \neq 0, Y \ge 0\\ Log(Y+1) ; & \text{if } l = 0, Y \ge 0\\ \frac{(1-Y)^{2-l}-1}{l-2} ; & \text{if } l \neq 2, Y < 0\\ -Log(1-Y) ; & \text{if } l = 0, Y < 0 \end{cases}$$
(7)

When estimating the transformation parameter, they found the value of λ that minimizes the Kullback-Leibler distance between the normal distribution and the transformed distribution.

1.1. The Estimation Process of λ in Box-Cox Transformation Method The main objective in the analysis of Box-Cox transformation model is to make inference on the transformation parameter λ and Box, et al [2] considered two approaches. In the first approach that employs the maximum likelihood estimator (MLE), one first assumes that transformed responses Y(l) follow a the multivariate normal distribution. i.e.. $Y(I) \approx N(X b, s^2 I_n)$. Then, in order to estimate the model parameters $(1, b, s^2)$, the design matrix X and the raw data Y are observed. Using the density for Y(1) as f(Y(1)) =

$$\frac{exp\left(\frac{-1}{2s^{2}}(Y(1)-Xb)'(Y(1)-Xb)\right)}{(2ps^{2})^{n/2}}$$
(8)

and letting J(I,Y) to be the Jacobian of the transformation from Y to Y(I), the density for Y (which is also the likelihood for the whole model) is

$$L(\boldsymbol{l},\boldsymbol{b},\boldsymbol{s}^{2}) = f(\boldsymbol{Y}) =$$
(9)

$$\frac{exp\left(\frac{-1}{2s^{2}}(Y(1)-Xb)'(Y(1)-Xb)\right)}{(2ps^{2})^{n/2}}J(1,Y)$$

To obtain the MLE from the likelihood Equation 9, we observe that for each fixed l, the likelihood

IJE Transactions A: Basics

170 - Vol. 24, No. 2, June 2011

equation is proportional to the likelihood equation for estimating (b, s^2) for observed Y(I). Thus the MLE for (b, s^2) are

$$b(1) = (X'X)^{-1}XY(1)$$
 (10)

$$s^{2}(1) = \frac{Y(1)'(I_{n} - X(X'X)^{-1}X')Y(1)}{n}$$
(11)

Substituting b(1) and $s^2(1)$ into likelihood equation, the likelihood function (9) will be maximized for 1 [4].

This method is a commonly used procedure since it is conceptually easy and the profile likelihood function is easy to compute. Also, it is easy to obtain an approximate confidence interval (CI) for l because of the asymptotic property of MLE.

For the second approach that is based on the Bayesian method, one needs to first ensure that the model is fully identifiable (see [8-10]).

Other families of transformations such as the folded power family can be found in literature (see [11]). However, they are rarely used because the resulting transformations have poor properties.

In this paper, instead of using the above two commonly used approaches of Box-Cox transformation procedures to find the required power transformation, a simple procedure based on the bisection method is proposed to make the inherent skewness of the existing skewed distribution zero. The method is developed for both positive and negative data.

The rest of the text is organized as follows. In Section 2, the proposed transformation method is developed. In order to investigate the performance of the proposed procedure and to compare it with the ones of the Box-Cox transformation technique, a simulation experiment is performed in Section 3 using differentknownskewedprobability distributions. Finally, Section 4 presents the conclusion.

2. ROOT TRANSFORMATION TECHNIQUE IN SKEWNESS REDUCTION

The most serious issue in the statistical analysis of non-normal data is the existing skewness in their probability distributions such that the usual standard statistical analysis is not applicable. Originally, Niaki, et al [1] used a root transformation technique to design multi-attribute control charts that are applicable to non-normal attribute data. Since the discrete probability distributions of the attribute data are right-skewed, their technique is only applicable to right-skewed distributions. In this research, after describing this technique, we generalize it to be applicable to almost any kind of non-normal data.

In the proposed power/root transformation technique of Niaki, et al [1], one searches for the best power (r) within (0, 1) such that if the data is raised to the power of r (i.e. Y^{r}), the transformed data will have almost zero skewness. The search is based on the bisection method to find the desired value of r.

The bisection method is a root-finding algorithm that repeatedly divides an interval in half and then selects the subinterval in which a root exists. Suppose one needs to solve the equation f(r) = 0 in the interval (a,b). The bisection method starts with two points a_0 and b_0 in (a,b)such that $f(a_0)$ and $f(b_0)$ have opposite signs. The method then divides the interval in two by computing $c_0 = (a_0 + b_0)/2$. There are now two possibilities; either $f(a_0)$ and $f(b_0)$ have opposite signs, or $f(c_0)$ and $f(b_0)$ have opposite signs. The location of the root is determined as lying within the subinterval with opposite signs. The bisection algorithm is then applied recursively to the subinterval where the sign-change occurs and eventually finds the root.

In order to find a root of f(r) = 0 in the interval (a,b), we start with the subinterval (a_0,b_0) such that $f(a_0)f(b_0) < 0$, pick a tolerance e and then apply the following algorithm:

$$k = 0$$

While $|f(r_{k+1}| > e$
 $r_{k+1} = \frac{a_k + b_k}{2}$
If $f(r_{k+1})f(a_k) < 0$ then
 $a_{k+1} = a_k$ and $b_{k+1} = r_k$

IJE Transactions A: Basics

Vol. 24, No. 2, June 2011 - 171

Else $b_{k+1} = b_k$ and $a_{k+1} = r_k$ End If k = k + 1End while $r^* = r_k$

In the root transformation technique, if we define f(r) to be the amount of skewness on the *r*th power-transformed *Y*, i.e. Y^r , we want to find *r* such that f(r) becomes zero. Therefore, applying the bisection method, we try to find a root (r) for f(r) = 0 in the interval (0,1).

mentioned previously, the procedure As described above is only applicable to right-skewed or slightly skewed (i.e. skewness very close to zero, either negative or positive) distributions. In cases where the distribution is heavily left-skewed (i.e. its skewness is a significant negative number), the data should be raised to a power bigger than one to dampen non-normality. To search for this power, we modify the power transformation technique for left-skewed data. This modification is quite simple; if data is heavily left-skewed, instead of raising Y to the power of r, it is raised to the power of 1/r, where r is still within interval (0,1). Therefore, data is actually raised to a power bigger than one. Furthermore, the results of a simulation experiment for negative-skewed data showed that if we initially subtract the data by its minimum value, the resulted skewnesses would become significantly less. Hence. before employing the modified power transformation technique for negative-skewed distributions, the data is first subtracted from its minimum value.

The second modification to the original power transformation technique applies to lognormal data. The power transformation technique of Niaki, et al [1]) is not able to make the skewness of lognormal data close to zero and leaves them almost unchanged (i.e. r > 0.99.) Therefore, we modify the proposed algorithm as follows; if the power transformation technique leaves the data

unchanged and skewness is still high, it means that the data follows a logarithmic-type distribution, and hence it is transformed by taking natural logarithm.

Finally, it should be mentioned that similar to the original Box-Cox transformation method, the original data of the proposed power transformation technique need to be positive. In this case, the absolute value of the minimum-valued data should be added to all to make them non-negatives. In summary the flowchart of the proposed algorithm is shown in Figure 1.

3. SIMULATION EXPERIMENTS

In order to investigate the performance of the proposed procedure and to compare it with the one of the complicated well-known Box-Cox transformation, simulation experiments have been conducted. In the simulation study, four different probability distributions of gamma, beta, weibull, and lognormal were used to produce all major types of non-normal skewed data. The simulation experiments are categorized into four classes of right-skewed, slightly-skewed, left-skewed, and logarithmic.

The results of the simulation experiments based on 10,000 samples of sizes n = 30, n = 100, and n=1,000, are presented in Tables 1 and 2. In both of these tables, there are smaller 4-by-4 tables for each distribution. The first column of each smaller table represents the basic statistics of the original data, the second and the third columns exhibit information on the transformed data using the proposed power transformation technique (namely treatment 1: where the first modification described in section two is employed, and treatment 2: in which the second modification is also used), and the fourth column represents information on the Box-Cox transformation of the data. The transformed data of treatment 2 has been subjected to bigger sample sizes than those of treatment 1 (i.e., more than 30, 100, and 1000). That is why treatment 2 outperforms treatment 1 in most cases.

Archive of SID



Figure 1. The flowchart of the proposed method.

The first and the second rows of the smaller tables show the skewness and the kurtosis of the data, respectively. The third and the fourth rows of these tables show the results of JB-test (Jarque Bera) for each column. The third row shows the average value of the test statistics (h is zero if the normality hypothesis is accepted, and one if rejected) for 10000 runs, and the fourth row is the average of the *p*-values. The less the third-row value, and the more the fourth-row value are, the better is the performance of the corresponding

treatment. Hence, we can use these two values as measures of performances to describe and compare the effectiveness of the treatments.

The results of both Tables 1 and 2 show that the proposed power transformation method performs quite well in all right-skewed, left-skewed and logarithmic distributions and transforms non-normal data to normal data efficiently in almost all cases (in most cases treatment 2 provides better results than those of treatment 1). The Box-Cox method also performs well and does slightly better

IJE Transactions A: Basics

Vol. 24, No. 2, June 2011 - 173

			Sample Size = 30				Sa	mple Siz	e = 100		Sample Size = 1000				
			Original Data	1	2	Box- Cox	Original Data	1	2	Box- Cox	Original Data	1	2	Box- Cox	
gamma (2, 2)	Skewness		1.0884	0.1500	0.1491	-0.0403	1.2854	0.0100	0.0097	-0.0231	1.3980	0.0008	-0.0004	-0.0135	
	Kurtosis		4.1470	2.8407	2.8401	2.5405	5.1440	2.7700	2.7692	2.7491	5.8738	2.8515	2.8508	2.8535	
	Jarque- Bera test	h	0.5783	0.0868	0.0866	0.0013	0.9950	0.0076	0.0077	0.0020	1.0000	0.0279	0.0254	0.0245	
		р	0.0756	0.4221	0.4222	0.4621	0.0044	0.4508	0.4515	0.4546	0.0010	0.3866	0.3926	0.3939	
gamma (1, 2)	Skewness		1.4788	0.0775	0.0765	-0.0682	1.7868	0.0020	0.0004	-0.0489	1.9725	0.0004	-0.0001	-0.0392	
	Kurtosis		5.2355	2.6568	2.6562	2.4833	7.1596	2.6488	2.6397	2.6504	8.7169	2.7105	2.7093	2.7199	
	Jarque- Bera test	h	0.8203	0.0373	0.0373	0.0005	1.0000	0.0011	0.0009	0.0008	1.0000	0.2179	0.2100	0.2074	
		р	0.0269	0.4412	0.4414	0.4576	0.0013	0.4346	0.4353	0.4356	0.0010	0.2013	0.2081	0.2069	
Weibull (1.5,3)	Skewness		0.1359	-0.0515	-0.0530	-0.0683	0.1546	-0.0292	-0.0251	-0.0494	0.1674	-0.0014	-0.0002	-0.0392	
	Kurtosis		2.6184	2.5437	2.5435	2.4814	2.6981	2.6508	2.6486	2.6510	2.7259	2.7100	2.7088	2.7193	
	Jarque- Bera test	h	0.0311	0.0140	0.0140	0.0005	0.0357	0.0012	0.0009	0.0006	0.8291	0.2192	0.2113	0.2072	
		р	0.3755	0.4316	0.4317	0.4581	0.3319	0.4310	0.4313	0.4358	0.0318	0.2009	0.2067	0.2055	
beta (2,2) (slightly right skewed)	Skewness		-0.0026	-0.1074	-0.1087	-0.1560	0.0012	-0.0679	-0.0602	-0.1506	-0.0002	-0.0146	-0.0205	-0.1493	
	Kurtosis		2.1793	2.1756	2.1764	2.1816	2.1562	2.1697	2.1667	2.2144	2.1442	2.1484	2.1499	2.2205	
	Jarque- Bera test	h	0.0016	0.0013	0.0013	0.0000	0.0434	0.0312	0.0319	0.0244	1.0000	1.0000	1.0000	1.0000	
		р	0.3359	0.3680	0.3680	0.3846	0.1465	0.1703	0.1692	0.1787	0.0010	0.0010	0.0010	0.0010	

 TABLE 1. Results of Simulation in 10,000 Replications of Right-Skewed Distributions.

than the proposed procedure for small sample sizes. In medium and large sample sizes, even treatment 1 performs better than the Box-Cox method. However, the difference is not significant. Combining this finding with the ease of understanding and conducting of the proposed method, one may conclude that the proposed procedure provides a better approach for practical applications.

4. CONCLUSIONS

In this paper, the power transformation technique that was developed by Niaki, et al [1] was extended to transform right and left skewed distributions to distributions that are closer to normal. The value of the power was searched such that the skewness of transformed data becomes almost zero with an acceptable error. The performance of the proposed method was then compared to the ones of the wellknown and complicated Box-Cox transformation method for different left-skewed, right-skewed and logarithmic distributions using Monte Carlo simulation. Besides simplicity in understanding and implementing, the results of the simulation study showed that the proposed method works well, and is quite compatible to the complicated method of Box, et al [2] in all of the simulated experiments. Although the *p*-values of the normality tests of the Box-Cox method are slightly more than the ones obtained by the proposed procedure, the simplicity of the latter method makes it preferable especially when large data are treated.

5. REFERENCES

- Niaki, S.T.A. and Abbasi, B., "Skewness Reduction Approach in Multi-Attribute Process Monitoring", *Communications in Statistics-Theory and Methods*, Vol. 36, (2007), 2313-2325.
- Box, G. and Cox, D.R., "An analysis of Transformation", Journal of Royal Statistical Society B, Vol. 26, (1964), 211–243.
- 3. Manly, B.F.J., "Exponential Data Transformations", *Statistician*, Vol. 25, (1976), 37-42.
- 4. Li, P., "Box-Cox Transformations: An Overview", Technical Report (2005), on the web: http://www.stat. uconn.edu/~studentjournal/index_files/pengfi_s05.pdf.
- John, J.A. and Draper, N.R., "An Alternative Family of Transformations", *Applied Statistics*, (1980), Vol. 29, (1980), 190-197.

174 - Vol. 24, No. 2, June 2011

IJE Transactions A: Basics

			Sa	ample Siz	ze = 30		Sa	mple Siz	e = 100		Sample Size = 1000				
			Original Data	1	2	Box- Cox	Original Data	1	2	Box- Cox	Original Data	1	2	Box- Cox	
beta (5,1)	Skewness		-1.0243	-0.0009	-0.0005	-0.2814	-1.1367	-0.0008	0.0000	-0.2891	-1.1782	-0.0014	0.0000	-0.2911	
	Kurtosis		3.5865	1.7770	1.7764	2.0055	4.0139	1.7726	1.7715	2.0005	4.1802	1.7832	1.7818	1.9965	
	Jarque- Bera test	h	0.5615	0.0001	0.0001	0.0000	0.9994	0.7808	0.7809	0.5886	1.0000	1.0000	1.0000	1.0000	
		р	0.0627	0.2174	0.2176	0.2498	0.0046	0.0419	0.0419	0.0523	0.0010	0.0010	0.0010	0.0010	
weibull (3,10)	Skewness		-0.5173	0.0185	0.0194	-0.0680	-0.5963	0.0003	0.0005	-0.0490	-0.6322	-0.0006	0.0000	-0.0392	
	Kurtosis		3.1165	2.3784	2.3776	2.4842	3.4046	2.5217	2.5207	2.6486	3.5495	2.6223	2.6210	2.7189	
	Jarque-	h	0.1955	0.0006	0.0006	0.0008	0.5531	0.0008	0.0010	0.0002	1.0000	0.5351	0.5267	0.2063	
	test	р	0.2557	0.4393	0.4394	0.4586	0.0955	0.3948	0.3955	0.4370	0.0010	0.0927	0.0957	0.2045	
5)	Skewness		-0.6222	0.0105	0.0115	-0.0679	-0.7245	-0.0008	0.0001	-0.0490	-0.7813	-0.0008	0.0000	-0.0394	
weibull (2,1	Kurtosis		3.3074	2.3536	2.3526	2.4830	3.7141	2.4932	2.4919	2.6470	3.9690	2.5980	2.5968	2.7185	
	Jarque- Bera test	h	0.2643	0.0004	0.0004	0.0004	0.7056	0.0014	0.0013	0.0008	1.0000	0.6252	0.6171	0.2132	
		р	0.2199	0.4368	0.4369	0.4576	0.0578	0.3824	0.3829	0.4362	0.0010	0.0719	0.0741	0.2055	
•	Skewness		1.1825	0.1625	0.1605	0.0008	1.4875	0.0969	0.0935	0.0001	1.7085	0.0391	0.0322	0.0000	
logn (0, 0.5	Kurtosis		4.6450	2.6979	2.6978	2.5847	6.4724	2.8838	2.8833	2.8347	8.4256	2.9878	2.9875	2.9814	
	Jarque- Bera test	h	0.6222	0.0244	0.0244	0.0018	0.9942	0.0269	0.0268	0.0072	1.0000	0.0332	0.0326	0.0123	
		р	0.0760	0.4190	0.4191	0.4657	0.0035	0.4175	0.4177	0.4589	0.0010	0.4112	0.4136	0.4501	
logn (0, 5)	Skewness		4.4544	0.1581	0.1557	0.0003	8.2382	0.0930	0.0900	0.0001	24.7876	0.0240	0.0211	0.0000	
	Kurtosis		22.2253	2.6885	2.6881	2.5800	74.0223	2.8892	2.8890	2.8397	678.2497	2.9901	2.9901	2.9824	
	Jarque- Bera test	h	1.0000	0.0248	0.0248	0.0020	1.0000	0.0290	0.0289	0.0062	1.0000	0.0355	0.0353	0.0136	
		р	0.0010	0.4198	0.4199	0.4654	0.0010	0.4152	0.4154	0.4575	0.0010	0.4061	0.4072	0.4492	

TABLE 2. Results of Simulation in 10,000 Replications of Left-Skewed and Logarithmic Distributions.

- 6. Bickel, P.J. and Doksum, K.A., "An Analysis of Transformations Revisited", *Journal of the American Statistical Association*, Vol. 76, (1981), 296-311.
- 7. Yeo, I.K. and Johnson, R., "A New Family of Power Transformations to Improve Normality or Symmetry", *Biometrika*, Vol. 87, (2000), 954-959.
- 8. Perrichi, L.R., "A Bayesian Approach to Transformations to Normality", *Biometrika*, Vol. 68, (1981), 35-43.
- Sweeting, T.J., "On the Choice of Prior Distribution for the Box-Cox Transformed Linear Model", *Biometrika*, Vol. 71, (1984), 127–134.
- Hinkley, D.V. and Runger, G., "The Analysis of Transformed Data", *Journal of the American Statistical Association*, Vol. 79, (1984), 302–309.
- Cook, R.D. and Weisberg, S., "Applied Regression Including Computing and Graphics", Wiley, New York, U.S.A., (1999).

IJE Transactions A: Basics