



# Chemical Composition of the Essential Oil from *Marrubium persicum* C. A. Mey. (Lamiaceae)

Sanaz Hamedeyazdan<sup>1</sup>, Fatemeh Fathiazad<sup>2\*</sup>, Solmaz Asnaashari<sup>3</sup>

<sup>1</sup> Student's Research Committee, Faculty of Pharmacy, Tabriz University of Medical Sciences, Iran.

<sup>2</sup> Department of Pharmacognosy, Faculty of Pharmacy, Tabriz University of Medical Sciences, Iran.

<sup>3</sup> Drug Applied Research Centre, Tabriz University of Medical Sciences, Iran.

## ARTICLE INFO

### Article Type:

Research Article

### Article History:

Received: 25 February 2013

Accepted: 19 September 2013

### Keywords:

*Marrubium persicum*

Lamiaceae

Essential oil composition

GC-MS

Non-terpenoids

Tolualdehyde

## ABSTRACT

**Background:** The essential oil of *Marrubium persicum* C. A. Mey (Lamiaceae) was obtained by hydrodistillation from aerial parts of the plant during flowering stage. **Methods:** The chemical analyses of the essential oil by GC/MS and GC/FID, allowed us to identify thirty three compounds, representing 94.4% of the total oil. **Results:** The essential oil of *M. persicum* was typically a complex mixture of mainly oxygenated non-terpenoids (51.5%), sesquiterpene hydrocarbons (27.9%), oxygenated sesquiterpenes (4.8%), monoterpene hydrocarbons (9%), and oxygenated monoterpenes (1.3%). The major components of the essential oil were m-tolualdehyde (19.2%) followed by acetophenone (14.6%), germacrene D (10.5%),  $\beta$ -caryophyllene (7.4%),  $\beta$ -farnesene (6.2%), and  $\alpha$ -pinene (4.6%). **Conclusion:** In spite of the fact that *Marrubium* sp. contain quite lower amounts of aliphatic and non-terpenoid fractions, *M. persicum* revealed rather higher proportions of non-terpenoid compounds, namely acetophenone with different isomers of tolualdehyde which are considered to be reported for the first time in the genus *Marrubium*.

## Introduction

Not surprisingly, the wide range of herbal medicines available today has been brought about by one of the most significant ancient heritages, the sophisticated experience of people who have tried over millennia to find useful plants for health improvement, with each generation adding its own experience to this tradition. Following on from scientific advancements, in a time of increased considerations to natural products as a safe herbal remedies, it is as important as ever to gather detailed documentations of the latter in every aspect.<sup>1</sup>

Lamiaceae family has been holding a place of value for hundreds of years due to the infusions and tinctures of numerous aromatic species used as components of herbal treatments for a variety of ailments.<sup>2</sup> Considering the endemism and diversity of the species used in traditional and folk medicine in Iran, *Marrubium* a genus of about 40 species of flowering plants in this family has a widespread usage by different cultures and also a high reputation in herbal medicine by several known healing attributes.<sup>3</sup>

*Marrubium* sp. are often perennial tomentose herbs, with wrinkled leaves, and small flowers in dense axillary clusters that are native to temperate regions of Europe and Asia.

Regarding the phytochemical analysis of the plants of this genus, they mainly produce diterpenes,

polyphenols, steroids, phenylpropanoids and flavonoids, some of which have important biological properties.<sup>4-8</sup> Generally, species of this genus are characterized through having potential therapeutic activities, which have been supported by various studies demonstrating cytotoxicity, immunomodulating, vasorelaxant, antispasmodic, hypolipidemic, hypoglycemic, and analgesic properties of this genus.<sup>9-15</sup> Furthermore, studies dealing with the composition, antimicrobial and antioxidant activities of essential oils extracted from different species of the genus *Marrubium* have been previously reported.<sup>16-25</sup> Based on Flora Iranica, *M. persicum*, is endemic to the countries; Armenia, Azerbaijan, Turkey and Iran which is considered to be an unknown source of chemical constituents, since there has been no study on *M. persicum*.<sup>26</sup> Regarding the number of reports on the chemical diversity and biological properties of the essential oils within a genus demonstrating both quantitative and qualitative diversity in the composition of the essential oils, assessing the volatile constituents naturally occurring in the essential oil of a plant seems to be of value. Hence, the present study was conducted to provide the very first and detailed GC-MS analysis of the volatile constituents from the aerial parts of *M. persicum* grown in Azarbaijan province in Iran. This

\*Corresponding Author: Fatemeh Fathiazad, Department of Pharmacognosy, Faculty of Pharmacy, Tabriz University of Medical Sciences, Iran.

Tel: +98 (411) 6692405, Fax: +98 (411) 3344798, Email: fathiazad@tbzmed.ac.ir

Copyright © 2013 by Tabriz University of Medical Sciences

report is a leading point in our study on the species of genus *Marrubium* of Iranian flora.

## Materials and Methods

### Plant Material

Aerial parts of *Marrubium persicum* C. A. Mey. were collected during the flowering stage from Varzeghan in East Azarbaijan province, Iran (38° 30' 33.9" 'N latitude, 46° 30' 41" E longitude, and 1940 m above sea level), in June 2011. A voucher specimen of the plant representing this collection has been deposited at the Herbarium of the Faculty of Pharmacy, Tabriz University of Medical science, Iran.

### Essential oil extraction

Air-dried plant material of the aerial parts of *M. persicum* was subjected to hydrodistillation using a Clevenger-type apparatus (Clevenger, 1928). Since the oil content was low in quantity, the distillation time was prolonged (5h) and xylene was used as an absorbing medium. The obtained essential oil was stored in sealed glass vial at 4-5°C prior to analysis.

### Gas Chromatography-Mass Spectrometry (GC-MS)

The essential oil was analyzed by GC-MS using a Shimadzu GC-MS-QP 5050A gas chromatograph fitted with a DB1 (methyl phenyl siloxane, 60 m x 0.25 mm i.d., 0.25 µm film thickness) capillary column. The GC was set at the following conditions with helium as the carrier gas; flow rate of 1.3 mL/min; linear velocity: 29.6 cm/s; Split ratio, 1:29; column temperature, 2 min in 60°C, 50-260 °C at 3 °C/min; injector temperature, 240 °C, and 1 µL of volume injection of the essential oil. The MS operating parameters were as follows: ionization potential, 70 eV; ion source temperature; 270 °C; quadrupole 100 °C, Solvent delay 2 min, scan speed 2000 amu/s and scan range 30-600 amu, EV voltage 3000 volts.

### Identification of the compounds

The identification of compounds was based on direct comparison of the retention indices and mass spectral data with those for the standards and by computer matching with the Wiley 229, Nist 107, Nist 21 Library, as well as by comparing the fragmentation patterns of the mass spectra with those reported in the literature.<sup>27</sup> For quantification purpose, relative area percentages were obtained by FID without the use of correction factors, where the FID detector condition was set on a duplicate of the same column applying the same operational conditions.

## Results

The hydrodistillation of the flowering aerial parts of *M. persicum* gave the essential oil in a very small quantity, justifying the categorization of the investigated species in the so-called "oil poor taxa" of the Lamiaceae family. The GC-MS analysis of the essential oil led to the identification of 33 different components,

representing 94.4% of total oil constituents. All the compounds are arranged in order of their elution from the DB1-MS column and the retention time with their percentage composition are summarized in Table 1.

**Table 1.** Chemical constituent of the essential oil from aerial parts of *M. persicum*.

No.	RI <sup>a</sup>	Compounds	Area (%)
1	906	n-Nonane	1.6
2	932	Benzaldehyde	0.9
3	936	α-Pinene	4.6
4	966	1-Octen-3-ol	1.6
5	970	Sabinene	1.3
6	986	Myrcene	0.6
7	1003	Decane	0.2
8	1011	α-Tolualdehyde	2.4
9	1025	Limonene	2.5
10	1036	Acetophenone	14.6
11	1040	m-Tolualdehyde	19.1
12	1052	o-Tolualdehyde	3.5
13	1085	Nonanal	0.5
14	1087	Linalool	0.4
15	1230	Terpinen-4-ol	0.5
16	1250	α-Terpineol	0.3
17	1381	α-Cubebene	0.3
18	1389	β-Bourbonene	0.4
19	1424	β-Caryophyllene	7.4
20	1450	β-Farnesene	6.1
21	1457	α-Humulene	0.8
22	1482	Germacrene D	10.5
23	1497	Bicyclgermacrene	1.3
24	1505	β-Bisabolene	0.7
25	1520	δ-Cadinene	0.4
26	1570	Spathulenol	0.5
27	1577	Caryophyllene oxide	2.1
28	1689	α-Cadinol	0.5
29	1748	α-Bisabolol	1.7
30	1833	Hexahydrofarnesyl acetone	2.9
31	2592	Hexacosane <sup>b</sup>	0.8
32	2793	Octacosane <sup>b</sup>	1.1
33	2892	Nonacosane <sup>b</sup>	2.3
<b>Total</b>			<b>94.4</b>
<b>Non-terpenoids</b>			<b>51.5</b>
<b>Monoterpene hydrocarbons</b>			<b>9</b>
<b>Oxygenated monoterpenes</b>			<b>1.2</b>
<b>Sesquiterpene hydrocarbones</b>			<b>27.9</b>
<b>Oxygenated sesquiterpenes</b>			<b>4.8</b>

<sup>a</sup>RI is the Retention Index relative to C8-C24 n-alkanes on the DB-1 column.  
<sup>b</sup>This compound was compared with an authentic sample.

The essential oil of *M. persicum* was typically a complex mixture of mainly oxygenated non-terpenoids (51.5%), sesquiterpene hydrocarbons (27.9%), oxygenated sesquiterpenes (4.8%), monoterpene hydrocarbons (9%), and oxygenated monoterpenes (1.2%). Among the non-terpenoid compounds, m-tolualdehyde, acetophenone, and  $\alpha$ -tolualdehyde were the chief constituents with 19.1%, 14.6% and 2.4% parts present in the essential oil, correspondingly. The major sesquiterpenes of the oil were identified as germacrene D (10.5%),  $\beta$ -caryophyllene (7.4%),  $\beta$ -farnesene (6.1%), and bicyclogermacrene (1.3%). In the case of identified monoterpenes,  $\alpha$ -pinene, limonene, and sabinene were considered as the most frequent constituents of the oil with the relative percentages of 4.6, 2.5, and 1.3%, respectively. Considering the oxygenated monoterpene and sesquiterpene compounds, terpinen-4-ol with 0.5% and caryophyllene oxide with 2.1% were the dominant components of the groupings. On the whole, the most abundant components of the *M. persicum* essential oil were m-tolualdehyde (19.1%) followed by acetophenone (14.5%), germacrene D (10.5%),  $\beta$ -caryophyllene (7.4%),  $\beta$ -farnesene (6.1%), and  $\alpha$ -pinene (4.6%).

### Discussion

Having reviewed the previously reports of the essential oils from genus *Marrubium*, underlying the fact that most species of *Marrubium* have rather low amounts of aliphatic and non-terpenoid fractions,<sup>16-25</sup> *M. Persicum* not only presented relatively higher amounts of non-terpenoids but also the most prominent constituent of the essential oil was a non-terpenoid compound, m-tolualdehyde. Meanwhile, it would be of note to mention new components, namely acetophenone together with different isomers of tolualdehyde present in the essential oil of *M. persicum*, were reported for the first time in the genus *Marrubium*. Regardless of the essential oil had been characterized with non-terpenoids as the most significant components, *M. persicum* essential oil contained relatively sizeable amounts of germacrene D supporting for the Lawrence concept that oil-poor species of the family Labiatae produce essential oils rich in sesquiterpene hydrocarbons, with germacrene D often being one of the predominant compounds.<sup>28</sup>

Furthermore, it is judicious to be reminiscent of the environmental factors like geography, altitude, temperature, day length, and edaphic conditions of the plant growing wild being notable agents playing key roles in the chemical composition of *M. persicum* oil. Seeing as, these factors could influence the biosynthetic pathways of plants, the relative proportion of the main characteristic compounds of the essential oils as the secondary metabolites would be variable. Based on the biosynthesis pathways of the essential oils, it is suggested that the production of terpenoid and non-isoprenoid compounds diverges early in the

pathway of anabolic plant secondary compound synthesis.<sup>29</sup>

### Conclusion

Regarding the similarities and differences in chemical composition of the oil compared to the other essential oil reports of different species from genus *Marrubium*, it was suggested that the variations in presence or absence and the identity of compounds within these species may arise from intrinsic (genetic, growth stage) and extrinsic conditions such as climatic, seasonal, environmental and distillation processes of these samples. Although, further studies are mandatory and would be of value in determining the origin of the differences detected among these essential oils, the authors hope this study could furnish the background for the basic experiments on the relative studies of this genus or family as it is the first report on the chemical constituents of *Marrubium persicum* essential oil.

### Acknowledgments

The authors would like to thank Drug Applied Research Center of Tabriz University of Medical Sciences for recording the mass spectra. Financial support of this work by the Research Vice-Chancellor of Tabriz University of Medical Sciences is faithfully acknowledged. This article was written based on a data set of PhD. thesis, registered in Tabriz University of Medical Sciences (5/4/6651- NO. 71).

### References

- Hadley SK, Petry JJ. Medicinal herbs: a primer for primary care. *Hosp Pract (1995)* 1999;34(6):105-6, 109-12, 115-6 passim.
- Naghbi F, Mosaddegh M, Mohammadi Motamed S, Ghorbani A. Labiatae Family in folk Medicine in Iran: from Ethnobotany to Pharmacology. *Iran J Pharm Res* 2005;2:63-79.
- Meyre-Silva C, Cechinel-Filho V. A review of the chemical and pharmacological aspects of the genus *marrubium*. *Curr Pharm Des* 2010;16(31):3503-18.
- Zaabat N, Hay AE, Michalet S, Darbour N, Bayet C, Skandrani I, et al. Antioxidant and antigenotoxic properties of compounds isolated from *Marrubium deserti* de Noe. *Food Chem Toxicol* 2011;49(12):3328-35.
- Alkhatib R, Joha S, Cheok M, Roumy V, Idziorek T, Preudhomme C, et al. Activity of ladanein on leukemia cell lines and its occurrence in *Marrubium vulgare*. *Planta med* 2010;76(1):86-7.
- Rigano D, Grassia A, Borrelli F, Aviello G, Piozzi F, Bruno M, et al. Phytochemical and pharmacological studies on the acetonic extract of *Marrubium globosum* ssp. libanoticum. *Planta med* 2006;72(6):575-8.
- Calis I, Hosny M, Khalifa T, Ruedi P. Phenylpropanoid glycosides from *Marrubium alysson*. *Phytochemistry* 1992;31(10):3624-6.

8. Karioti A, Skaltsa H, Heilmann J, Sticher O. Acylated flavonoid and phenylethanoid glycosides from *Marrubium velutinum*. *Phytochemistry* 2003;64(2):655-60.
9. Karioti A, Skopeliti M, Tsitsilonis O, Heilmann J, Skaltsa H. Cytotoxicity and immunomodulating characteristics of labdane diterpenes from *Marrubium cylleneum* and *Marrubium velutinum*. *Phytochemistry* 2007;68(11):1587-94.
10. Rigano D, Aviello G, Bruno M, Formisano C, Rosselli S, Capasso R, et al. Antispasmodic effects and structure-activity relationships of labdane diterpenoids from *Marrubium globosum* ssp. *libanoticum*. *J Nat Prod* 2009;72(8):1477-81.
11. El Bardai S, Lyoussi B, Wibo M, Morel N. Pharmacological evidence of hypotensive activity of *Marrubium vulgare* and *Foeniculum vulgare* in spontaneously hypertensive rat. *Clin Exp Hypertens* 2001;23(4):329-43.
12. El Bardai S, Lyoussi B, Wibo M, Morel N. Comparative study of the antihypertensive activity of *Marrubium vulgare* and of the dihydropyridine calcium antagonist amlodipine in spontaneously hypertensive rat. *Clin Exp Hypertens* 2004;26(6):465-74.
13. Berrougui H, Isabelle M, Cherki M, Khalil A. *Marrubium vulgare* extract inhibits human-LDL oxidation and enhances HDL-mediated cholesterol efflux in THP-1 macrophage. *Life Sci* 2006;80(2):105-12.
14. Boudjelal A, Henchiri C, Siracusa L, Sari M, Ruberto G. Compositional analysis and in vivo anti-diabetic activity of wild Algerian *Marrubium vulgare* L. infusion. *Fitoterapia* 2012;83(2):286-92.
15. Meyre-Silva C, Yunes RA, Schlemper V, Campos-Buzzi F, Cechinel-Filho V. Analgesic potential of marrubiin derivatives, a bioactive diterpene present in *Marrubium vulgare* (Lamiaceae). *Farmaco* 2005;60(4):321-6.
16. Sarikurku C, Tepe B, Daferera D, Polissiou M, Harmandar M. Studies on the antioxidant activity of the essential oil and methanol extract of *Marrubium globosum* subsp. *globosum* (Lamiaceae) by three different chemical assays. *Bioresour Technol* 2008;99(10):4239-46.
17. Laouer H, Yabrir B, Djeridane A, Yousfi M, Beldovini N, Lamamra M. Composition, antioxidant and antimicrobial activities of the essential oil of *Marrubium deserti*. *Nat Prod Commun* 2009;4(8):1133-8.
18. Petrovic S, Pavlovic M, Maksimovic Z, Milenkovic M, Couladis M, Tzakouc O, et al. Composition and antimicrobial activity of *Marrubium incanum* Desr. (Lamiaceae) essential oil. *Nat Prod Commun* 2009;4(3):431-4.
19. Javidnia K, Miri R, Soltani M, Khosravi AR. Constituents of the Essential Oil of *Marrubium astracanicum* Jacq. from Iran. *J Essent Oil Res* 2007;19:559-61.
20. Zarai Z, Kadri A, Ben Chobba I, Ben Mansour R, Bekir A, Mejdoub H, et al. The in-vitro evaluation of antibacterial, antifungal and cytotoxic properties of *Marrubium vulgare* L. essential oil grown in Tunisia. *Lipids Health Dis* 2011;10:161.
21. Argyropoulou C, Skaltsa H. Identification of essential oil components of *Marrubium thessalum* Boiss. & Heldr., growing wild in Greece. *Nat Prod Res* 2012;26(7):593-9.
22. Demirci B, Baser KGC, Kirimer N. Composition of the essential oil of *Marrubium bourgaei* ssp. *caricum* P.H. Davis. *J Essent Oil Res* 2004;16:133-4.
23. Tajbakhsh M, Khalilzadeh A, Rineh A, Balou J. Essential oils of *Marrubium anisodon* C. Koch and *Marrubium propinquum* Fisch. et C.A. Mey., growing wild in Iran. *J Essent Oil Res* 2008:161-2.
24. Nagy M, Svajdlenka E. Comparison of essential oils from *Marrubium vulgare* L. and *M. peregrinum* L. *J Essent Oil Res* 1998;10:585-7.
25. Semnani KM, Saeedi M, Babanezhad E. The essential oil composition of *Marrubium vulgare* L. from Iran. *J Essent Oil Res* 2008;20(6):488-90.
26. Mozaffarian V. A Dictionary of Iranian Plant Names. Tehran, Iran: Farhang Moaser Publication; 2004.
27. Adams RP. Identification of Essential oil Components by Gas Chromatography/ Quadrupole Mass Spectroscopy. Carol Stream, IL: Allured Publ Corp; 2004.
28. Lawrence BM. Essential oils. Carol Strem, IL: Allured Publishing Corp; 1992.
29. Ganjewala D, Luthra R. Essential oil biosynthesis and regulation in the genus *Cymbopogon*. *Nat Prod Commun* 2010;5(1):163-72.