



Research Article



Propylthiouracil-Induced Mitochondrial Dysfunction in Liver and its Relevance to Drug-Induced Hepatotoxicity

Akram Jamshidzadeh^{1,2}, Hossein Niknahad^{1,2}, Reza Heidari^{1*}, Maryam Azadbakht², Forouzan Khodaei², Mohammad Reza Arabnezhad², Omid Farshad^{2,3}

¹Pharmaceutical Sciences Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.

²Department of Pharmacology and Toxicology, School of Pharmacy, Shiraz University of Medical Sciences, Shiraz, Iran.

³Students Research Committee, School of Pharmacy, International Branch, Shiraz University of Medical Sciences, Shiraz, Iran.

Article Info

Article History:

Received: 26 January 2017

Accepted: 5 May 2017

ePublished: 30 June 2017

Keywords:

-Antithyroid drugs
 -Bioenergetics
 -Liver Failure
 -Mitochondrial Dysfunction

ABSTRACT

Background: Propylthiouracil (PTU) administration is associated with several cases of hepatotoxicity, especially in children. The mechanism(s) of PTU-induced hepatotoxicity is obscure. In the current study, we aimed to assess the effect of PTU on hepatocytes mitochondria in different experimental models.

Methods: Mice were treated with PTU (10, 20, 40, 80, and 100 mg/kg, i.p) then, the liver mitochondria were isolated and evaluated. Moreover, liver mitochondria were isolated from normal mice and incubated with increasing concentrations of PTU (10 μ M-1 mM). Mitochondrial dehydrogenases activity, mitochondrial membrane potential, mitochondrial swelling, and mitochondrial adenosine triphosphate (ATP) content were monitored.

Results: PTU hepatotoxicity was biochemically evident in mice by increased serum biomarkers of liver injury. PTU also caused a decrease in mitochondrial dehydrogenases activity, increased mitochondrial swelling, depleted mitochondrial ATP, and caused mitochondrial depolarization both *in vitro* and *in vivo*.

Conclusion: Our data suggest mitochondrial dysfunction as a mechanism for PTU-induced hepatotoxicity.

Introduction

Drug-induced hepatotoxicity is a big challenge in pharmacotherapy and drug development.^{1,2} Antithyroid drugs-induced hepatotoxicity is a clinical complication.^{3,4} Propylthiouracil (PTU) is prescribed against hyperthyroidism especially in situations such as pregnancy.⁵ On the other hand, PTU administration is associated with several adverse drug reactions including taste loss, gastrointestinal disturbances, and lupus-like syndrome.^{6,7} PTU also causes serious adverse effects such as agranulocytosis and liver failure.⁸⁻¹⁰ Several cases of PTU-induced hepatotoxicity have been reported.^{4,11-16} Pediatrics seems to be more vulnerable to PTU-induced hepatotoxicity.¹⁷ There is no precise mechanism for PTU-induced liver injury.^{12,18}

The etiology of drug-induced liver injury sometimes underlies mitochondrial toxicity.¹⁹ Mitochondrial dysfunction could play a pivotal role in cell death and organ failure.¹⁹ Previous studies mention the role of mitochondrial dysfunction in thioamide-based drugs toxicity.^{20,21} PTU is a

thionamide antithyroid drug. Previously, we also found that methimazole as another thionamide antithyroid drug caused mitochondrial injury in isolated rat hepatocytes.^{20,22}

In the current study, we aimed to investigate the effect of PTU on hepatic mitochondria in two different experimental models. The data obtain from this study might help to provide therapeutic/preventive options against PTU-induced hepatotoxicity.

Materials and Methods

Chemicals

3-(N-morpholino)propane sulfonic acid (MOPS), Ethylene diamine tetra acetic acid (EDTA), Dimethyl sulfoxide (DMSO), Sodium succinate, 6-Propyl-2-Thio uracil (PTU), 4,2-Hydroxyethyl,1-piperazineethanesulfonic acid (HEPES), D-mannitol, Rhodamine123 (Rh 123), 2,3-[4,5dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT), Fatty acid-free bovine serum albumin (BSA) fraction V, Sucrose, Coomassie brilliant blue, and Ethylene glycol-bis (2-

*Corresponding Author: Reza Heidari, E-mail: rezaheidari@hotmail.com

©2017 The Authors. This is an open access article and applies the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

aminoethylether)-N,N,N',N'-tetraacetic acid (EGTA) were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Hydroxymethyl aminomethane hydrochloride (Tris-HCl) and Trichloroacetic acid (TCA) were purchased from Merck (Darmstadt, Germany). The luciferase–luciferin-based kit for ATP measurement was from Promega (ENLITEN®). The kits for serum biochemistry were obtained from Pars Azmun® (Tehran, Iran). All other chemicals used for buffer solutions were of analytical grade and obtained from Merck (Darmstadt, Germany).

Animals

Male BALB/c mice (20-25 g, n=84) were obtained from Comparative and Experimental Medicine Research Center, Animal Breeding Department, Shiraz University of medical Sciences, Shiraz Iran. Mice were housed in cages on chip-wood bedding at a constant temperature of 23±1°C and relative humidity of 40%. Animals had free access to tap water and a typical chow diet. Mice received human care and handled according to an animal handling protocol approved by a local ethics committee at Shiraz University of Medical Sciences, Shiraz, Iran (#11453).

In vivo experiments

Mice (n=42) were treated with PTU (10, 20, 40, 80, and 160 mg/kg, i.p). The control animals received PTU vehicle (Normal saline, 5 ml/kg, i.p). Five hours later, mice were sacrificed and their liver mitochondria was isolated based on a previously reported protocol.^{23,24}

Serum biochemistry and liver histopathology

Blood samples were collected from vena cava of the anesthetized animals (Thiopental, 50 mg/kg, i.p). Samples was transferred to standard tubes (Improvacuter®, gel and clot activator-coated tubes; Guangzhou, China) and blood serum was prepared by centrifugation (3000 g, 4°C, 15 min.). The liver tissue was removed and washed in ice-cold sodium chloride solution (0.9% w:v). A Mindray® BS-200 auto-analyzer (Mindray chemistry analyzers for low-volume laboratories, Guangzhou, China) and commercial kits (Pars Azmun®, Tehran, Iran), were used to evaluate serum lactate dehydrogenase (LDH), alanine aminotransferase (ALT), and aspartate amino transferase (AST) in PTU-treated mice.^{25,26} Liver samples were fixed in buffered formalin solution (0.4% sodium phosphate monobasic, NaH₂PO₄, 0.64% sodium phosphate dibasic, Na₂HPO₄, and 10% formaldehyde in distilled water) and used for tissue histopathological assessment.^{27,28} Paraffin-embedded blocks of liver tissue were prepared and stained with haematoxylin and eosin (H&E) before viewing by a light microscope

(Olympus CX21®, Japan).

In vitro experimental design

Increasing concentrations of PTU (10 µM-1 mM) were added to isolated liver mitochondria and several mitochondrial indices were assessed. Control mitochondria received no specific treatment and incubated at the same conditions.

Liver mitochondria isolation

Mice liver mitochondria were isolated by differential centrifugation method based on a previously reported protocol.²³ Briefly, animals were sacrificed by cervical dislocation and their liver was quickly excised and washed with cold (4°C) sodium chloride solution (0.9% w: v).^{23,29} The liver tissue (at a 10:1 buffer to liver ratio) was homogenized in a solution containing 225 mM sucrose, 75 mM mannitol, 2 mM MOPS, 0.1% BSA, and 0.5 mM EGTA (pH = 7.4).²³ Tissue homogenate was centrifuged (1000 g, 4°C, 10 minutes) to remove cell debris, intact cells, and cell nuclei. To precipitate mitochondrial fraction, the supernatants were further centrifuged at 10000 g, 4°C, for 10 minutes.²³ Using fresh buffer medium, the last step was repeated three times. As mentioned, all manipulations during mitochondrial isolation process were performed on ice (4°C) to minimize mitochondrial injury.²³

Mitochondrial dehydrogenase activity (MTT assay)

A colorimetric method based on the 3-(4, 5-dimethylthiazol-2-yl)-2, the 5-diphenyltetrazolium bromide reduction (MTT assay) was used for assessment of mitochondrial dehydrogenases activity.^{30,31} Liver in a buffer containing 320 mM sucrose, 1 mM EDTA, and 10 mM Tris-HCl (pH = 7.4), was incubated with 40 µl of the MTT solution (1 mg/ml) at 37°C for 30 minutes in dark. The product of purple formazan crystals was dissolved in 1 ml dimethyl sulfoxide (DMSO). Then, 100 µl of dissolved formazan was added to 96 well plate and the optical density at 570 nm was measured with an EPOCH plate reader (BioTek® Instruments, Highland Park, USA).^{31,32} Samples protein concentrations were determined by the Bradford method.³³

Mitochondrial membrane potential

Rhodamine 123 was used as the cationic fluorescent probe for assessment of mitochondrial membrane potential.^{23,32} Briefly, fractions (0.5 mg protein/ml) of prepared liver mitochondria were incubated with rhodamine 123 (Final concentration of 10 µM) in a buffer containing 65 mM KCl, 125 mM sucrose, 10 mM HEPES and 5 mM sodium succinate (pH=7.2, 30 min, 37°C). Samples were centrifuged (15,000 g, 10 min, 4 °C), and the

fluorescence intensity of the supernatant was measured using a FLUOstar Omega® multifunctional fluorescent microplate reader ($\lambda_{excitation} = 485 \text{ nm}$ and $\lambda_{emission} = 525 \text{ nm}$).^{23,31}

Mitochondrial Swelling assay

Monitoring the decrease in light absorbance at 540 nm was used as method assess mitochondrial swelling.²³ Briefly, mitochondrial samples (0.5 mg protein/ml) were suspended in swelling buffer (125 mM sucrose, 5 mM sodium succinate, 65 mM KCl, and 10 mM HEPES, pH=7.2). To start mitochondrial swelling, calcium (200 μM) was added after 5 minutes of samples incubation. Light absorbance ($OD_{540 \text{ nm}}$) was assessed using a FLUOstar Omega® multifunctional fluorescent microplate reader with constant shaking (100 rpm orbital shaking before each read) and temperature (30°C).^{23,34} A decrease in light absorbance is directly associated with an increase in mitochondrial volume.²³

Mitochondrial ATP level

A commercial kit based on luciferase–luciferin reaction (Promega, ENLITEN® ATP Assay System, Promega Corporation, Madison, USA) was used to monitor mitochondrial ATP level.^{32,35} Samples and buffer solutions were prepared based on the kit manual and the luminescence intensity was measured using a FLUOstar Omega® multifunctional microplate reader ($\lambda=560 \text{ nm}$).



Statistical analysis

Data are presented as the Mean±SD. The one-way analysis of variance (ANOVA) with Tukey's multiple comparison test as a *post hoc* was used for data comparison. $P < 0.05$ was considered as statistically significant difference between groups.

Results

PTU-induced hepatotoxicity in mice was revealed by significant elevation in serum biomarkers of liver injury (Figure 1). It was found that serum LDH, ALT, and AST levels were significantly elevated in PTU-treated animals (Figure 1). Moreover, liver histopathological changes were evident as inflammation, tissue necrosis, and sinusoidal dilation in PTU-treated animals (Figure 1).

Evaluation of liver mitochondria isolated from PTU-treated animals revealed a decrease in mitochondrial dehydrogenases enzyme activity (MTT assay), and the collapse of mitochondrial membrane potential ($\Delta\Psi$) (Figure 2). In addition, mitochondrial ATP level was decreased in mitochondria isolated from PTU-treated animals (Figure 3). Moreover, mitochondrial swelling was increased in the PTU-treated mice (Figure 4).



	Inflammation	Necrosis	Congestion	Vacuolization	Total
Control	0	0	0	0	Normal
PTU 160 mg/kg	+++	++	++	++	9

Figure 1. Serum biochemistry of PTU-treated mice. PTU: Propylthiouracil. Data are presented as Mean±SD (n=7) as assessed 5 hours after PTU administration. *Indicates significantly different as compared with control ($P < 0.01$). Histopathological evaluation (H&E staining) of mice liver revealed tissue necrosis, sinusoidal dilation and inflammatory cells aggregation in PTU-treated animals (B) in comparison with vehicle-treated group (A). Magnification 40x; Scale bar 1000 μm .

A significant decrease in mitochondrial succinate dehydrogenase activity (SDA) was also evident when isolated mitochondria were incubated with PTU (*In vitro*) (Figure 2). Further assessment of PTU-treated liver mitochondria, revealed a marked collapse of mitochondrial membrane potential

(Figure 2). It was also found that mitochondrial ATP content was depleted in PTU-treated groups (Figure 3). Moreover, when isolated liver mitochondria were incubated with PTU, an increase in mitochondrial swelling was detected (Figure 4).




Figure 2. Effect of PTU on liver mitochondria. PTU: Propylthiouracil. PTU caused a decrease in succinate dehydrogenase (SDA) activity (MTT test) (A and B) and collapse in mitochondrial membrane potential (C and D).

Data are given as Mean±SD (n=7).

Data in colored boxes are significantly different as compared with control (P<0.001).




Figure 3. Effect of PTU on mitochondrial ATP level *in vitro* (A) and *in vivo* (B). PTU: Propylthiouracil. Data are given as Mean±SD (n=7).

*Indicates significantly different as compared with control (P<0.01).




Figure 4. PTU-induced mitochondrial swelling *in vitro* (A) and *in vivo* (B). PTU: Propylthiouracil. Data are given as Mean \pm SD (n=7). *Indicates significantly different as compared with control (P<0.05).

Discussion

Hepatotoxicity is a major clinical complication associated with antithyroid drugs.^{36,37} There is several human cases of PTU-induced hepatotoxicity.^{15,38-39} The mechanism(s) of PTU hepatotoxicity is obscure. In this study, we found that PTU caused mitochondrial dysfunction both *in vitro* and *in vivo*. PTU administration caused mitochondrial depolarization, decreased succinate dehydrogenases activity and ATP level. Moreover, mitochondrial swelling was increased in PTU-treated groups.

Effects of PTU on liver could manifest as a transient elevation of serum transaminases to acute liver failure and death.^{11,13} On the other hand, clinicians advised frequent monitoring of liver function in PTU-treated patients.^{11,13,17,40-44} Hence, finding the mechanism(s) of PTU-induced hepatotoxicity might help to prevent or cure liver injury induced by this drug.

Previously, we found that PTU-induced liver injury was associated with oxidative stress and its associated complications in the liver.²⁴ Oxidative stress might play a role as a cause and/or the consequence of mitochondrial injury.⁴⁵ The mitochondrial injury is involved in the mechanism of hepatotoxicity induced by many xenobiotics.^{19,46,47} Mitochondrial injury could play a pivotal role in cell death and apoptosis process.^{48,49} Hence, xenobiotics-induced mitochondrial injury could lead to cell injury and hepatotoxicity.

It has been found that thionamide-based antithyroid agents such as methimazole caused mitochondrial injury which was revealed by severe mitochondrial depolarization (decreased $\Delta\Psi_m$).^{20,22} Methimazole reactive metabolites might play a role in its adverse

effects toward hepatocytes mitochondria.^{20,22} PTU is also a thionamide-based antithyroid agent with several cases of severe liver injury.^{44,50-52} Mitochondria seem to be a target for PTU-induced toxicity. Interestingly, megamitochondria (big mitochondria in shape and size) is a pathologic finding in PTU-induced hepatotoxicity.⁵³ Mitochondrial membrane fragmentation and lysis of the mitochondrial matrices has been observed in liver biopsies taken from human cases of PTU-induced hepatotoxicity.⁵³ All these findings might indicate that mitochondrial injury might be involved in the mechanism of PTU-induced liver injury.

Although the exact role of reactive metabolites of PTU in the liver has not been fully investigated, but this drug might convert to hepatotoxic metabolites.¹² The association between PTU reactive metabolite(s) and drug-induced hepatotoxicity is not clear. Further investigation on the role of drug metabolism and the effect of PTU reactive metabolite(s) on vital organelles such as mitochondria might shed some light the mechanisms of PTU-induced liver injury.

Several other thionamide-based structures are known as hepatotoxic agents.^{54,55} Thiourea is the parent chemical for many drugs and industrial agents. Some anti-tuberculosis drugs, anti-HIV reverse transcriptase inhibitors, and centrally acting histamine H₃ antagonists, are among thiourea-containing drugs.^{44,50-51,56,57} Interestingly, the clinical administration of the mentioned drugs is associated with drug-induced liver injury.^{55,58} Mitochondrial dysfunction might play a relevant role in the mechanism of liver injury induced by these drugs.^{55,58}

In conclusion, our data suggest that hepatocytes-mitochondria could be a possible target for PTU-induced liver injury. Hence, this study might shed some light on the current knowledge about the mechanism of PTU-induced liver injury. Moreover, these data might help to develop new preventive and therapeutic options against drugs-induced liver injury. The effect of PTU on targets such as mitochondrial respiratory chain enzymes or tricarboxylic acid cycle could be the subject of future studies.

Acknowledgments

The Current investigation was financially supported by the office of Vice-Chancellor of Research Affairs of Shiraz University of Medical Sciences (Grant number: 95-01-36-11453). The authors also gratefully acknowledge the instrumental facility providing of Pharmaceutical Sciences Research Center (PSRC) of Shiraz University of Medical Sciences for this study.

Conflict of interests

The authors claim that there is no conflict of interest.

References

- Grant LM, Rockey DC. Drug-induced liver injury. *Curr Opin Gastroenterol*. 2012;28(3):198-202. doi:10.1097/mog.0b013e3283528b5d
- Hayashi PH, Chalasani NP. Increasing impact of drug-induced liver injury. *Clin Liver Dis*. 2015;5(6):136-8. doi:10.1002/cld.469
- Akmal A, Kung J. Propylthiouracil, and methimazole, and carbimazole-related hepatotoxicity. *Expert Opin Drug Saf*. 2014;13(10):1397-1406. doi:10.1517/14740338.2014.953796
- Woeber KA. Methimazole-induced hepatotoxicity. *Endocr Pract*. 2002;8(3):222-24. doi:10.4158/ep.8.3.222
- Emiliano AB, Governale L, Parks M, Cooper DS. Shifts in propylthiouracil and methimazole prescribing practices: antithyroid drug use in the United States from 1991 to 2008. *J Clin Endocrinol Metab*. 2010;95(5):2227-33. doi:10.1210/jc.2009-2752
- Cooper DS. Antithyroid drugs. *N Eng J Med*. 2005;352(9):905-17. doi:10.1056/NEJMra042972
- Nelson P, Gossain VV. Prolonged treatment with antithyroid drugs: how long to treat safely; months, years, or a lifetime? *Endocrinologist*. 2010;20(5):217-9. doi:10.1097/ten.0b013e3181f47fb2
- Livadas S, Xyrafis X, Economou F, Boutzios G, Christou M, Zerva A, et al. Liver failure due to antithyroid drugs: report of a case and literature review. *Endocrine*. 2010;38(1):24-8. doi:10.1007/s12020-010-9348-y
- Cooper DS. The side effects of antithyroid drugs. *Endocrinologist*. 1999;9:457-78.
- Karras S, Tzotzas T, Krassas GE. Toxicological considerations for antithyroid drugs in children. *Expert Opin Drug Metab Toxicol*. 2011;7(4):399-410. doi:10.1517/17425255.2011.557068
- Carrion AF, Czul F, Arosemena LR, Selvaggi G, Garcia MT, Tekin A, et al. Propylthiouracil-induced acute liver failure: role of liver transplantation. *Int J Endocrinol*. 2010;2010:1-5. doi:10.1155/2010/910636
- Heidari R, Niknahad H, Jamshidzadeh A, Eghbal MA, Abdoli N. An Overview on the proposed mechanisms of antithyroid drugs-induced liver injury. *Adv Pharm Bull*. 2015;5(1):1-11. doi:10.5681/apb.2015.001
- Williams KV, Nayak S, Becker D, Reyes J, Burmeister LA. Fifty Years of Experience with Propylthiouracil-Associated Hepatotoxicity: What Have We Learned? *J Clin Endocrinol Metab*. 1997;82(6):1727-33. doi:10.1210/jc.82.6.1727
- Gallelli L, Staltari O, Palleria C, De Sarro G, Ferraro M. Hepatotoxicity induced by methimazole in a previously health patient. *Curr Drug Saf*. 2009;4(3):204-6. doi:10.2174/157488609789006912
- Yang J, Zhong J, Zhou L-Z, Hong T, Xiao X-H, Wen G-B. Sudden onset agranulocytosis and hepatotoxicity after taking methimazole. *Intern Med*. 2012;51(16):2189-92. doi:10.2169/internalmedicine.51.7845
- Wang M-T, Lee W-J, Huang T-Y, Chu C-L, Hsieh C-H. Antithyroid drug-related hepatotoxicity in hyperthyroidism patients: a population-based cohort study. *Br J Clin Pharmacol*. 2014;78(3):619-29. doi:10.1111/bcp.12336
- Karras S, Memi E, Kintiraki E, Krassas GE. Pathogenesis of propylthiouracil-related hepatotoxicity in children: present concepts. *J Pediatr Endocrinol Metab*. 2012;25(7-8):623-30. doi:10.1515/jpem-2012-0059
- Heidari R, Niknahad H, Jamshidzadeh A, Abdoli N. Factors affecting drug-induced liver injury: antithyroid drugs as instances. *Clin Mol Hepatol*. 2014;20(3):237-48. doi:10.3350/cmh.2014.20.3.237
- Labbe G, Pessayre D, Fromenty B. Drug-induced liver injury through mitochondrial dysfunction: mechanisms and detection during preclinical safety studies. *Fundam Clin Pharmacol*. 2008;22(4):335-53. doi:10.1111/j.1472-8206.2008.00608.x
- Heidari R, Babaei H, Eghbal M. Mechanisms of methimazole cytotoxicity in isolated rat

- hepatocytes. *Drug Chem Toxicol.* 2013;36(4):403-11. doi:10.3109/01480545.2012.749272
21. Sakamoto T, Kondo K, Kashio A, Suzukawa K, Yamasoba T. Methimazole-induced cell death in rat olfactory receptor neurons occurs via apoptosis triggered through mitochondrial cytochrome c-mediated caspase-3 activation pathway. *J Neurosci Res.* 2007;85(3):548-57. doi:10.1002/jnr.21155
 22. Heidari R, Babaei H, Eghbal MA. Ameliorative Effects of taurine against methimazole-induced cytotoxicity in isolated rat hepatocytes. *Sci Pharm.* 2012;80(4):987-99. doi:10.3797/scipharm.1205-16
 23. Caro AA, Adlong LW, Crocker SJ, Gardner MW, Luikart EF, Gron LU. Effect of garlic-derived organosulfur compounds on mitochondrial function and integrity in isolated mouse liver mitochondria. *Toxicol Lett.* 2012;214(2):166-74. doi:10.1016/j.toxlet.2012.08.017
 24. Karamikhah R, Jamshidzadeh A, Azarpira N, Saeedi A, Heidari R. Propylthiouracil-induced liver Injury in mice and the protective role of taurine. *Pharm Sci.* 2015;21(2):94-101. doi:10.15171/ps.2015.23
 25. Heidari R, Esmailie N, Azarpira N, Najibi A, Niknahad H. Effect of ehiol-reducing agents and antioxidants on sulfasalazine-induced hepatic injury in normotermic recirculating isolated perfused rat liver. *Toxicol Res.* 2016;32(2):133-40. doi:10.5487/tr.2016.32.2.133
 26. Heidari R, Sadeghi N, Azarpira N, Niknahad H. Sulfasalazine-induced hepatic injury in an ex vivo model of isolated perfused rat liver and the protective role of taurine. *Pharm Sci.* 2015;21(4):211-9. doi:10.15171/ps.2015.39
 27. Moezi L, Heidari R, Amirghofran Z, Nekooeian AA, Monabati A, Dehpour AR. Enhanced anti-ulcer effect of pioglitazone on gastric ulcers in cirrhotic rats: The role of nitric oxide and IL-1b. *Pharmacol Rep.* 2013;65(1):134-43. doi:10.1016/s1734-1140(13)70971-x
 28. Jamshidzadeh A, Heidari R, Abazari F, Ramezani M, Khodaei F, Ommati MM, et al. Antimalarial drugs-induced hepatic injury in rats and the protective role of carnosine. *Pharm Sci.* 2016;22(3):170-80. doi:10.15171/ps.2016.27
 29. Zhao P, Kalhorn TF, Slattery JT. Selective mitochondrial glutathione depletion by ethanol enhances acetaminophen toxicity in rat liver. *Hepatology.* 2002;36(2):326-35. doi:10.1053/jhep.2002.34943
 30. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods.* 1983;65(1-2):55-63. doi:10.1016/0022-1759(83)90303-4
 31. Niknahad H, Jamshidzadeh A, Heidari R, Hosseini Z, Mobini K, Khodaei F, et al. Paradoxical effect of methimazole on liver mitochondria: In vitro and in vivo. *Toxicol Lett.* 2016;259:108-15. doi:10.1016/j.toxlet.2016.08.003
 32. Jamshidzadeh A, Heidari R, Abasvali M, Zarei M, Ommati MM, Abdoli N, et al. Taurine treatment preserves brain and liver mitochondrial function in a rat model of fulminant hepatic failure and hyperammonemia. *Biomed Pharmacother.* 2017;86:514-20. doi:10.1016/j.biopha.2016.11.095
 33. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976;72(1-2):248-54. doi:10.1016/0003-2697(76)90527-3
 34. Niknahad H, Jamshidzadeh A, Heidari R, Abdoli N, Ommati MM, Jafari F, et al. The postulated hepatotoxic metabolite of methimazole causes mitochondrial dysfunction and energy metabolism disturbances in liver. *Pharm Sci.* 2016;22(4):217-26. doi:10.15171/ps.2016.35
 35. Held P. Luminescent Determination of ATP Concentrations using the clarity™ luminescence microplate reader. *Nat Methods. Application Notes.* 2006. doi:10.1038/an1705
 36. Lee KY, Lee YJ, Hong SH, Jung SK, Lee HE, Seo CJ, et al. Hepatic injury during treatment with antithyroid drugs in patients with hyperthyroidism. *J Korean Soc Endocrinol.* 2000;15(4-5):554-60.
 37. Vitug AC, Goldman JM. Hepatotoxicity from antithyroid drugs. *Horm Res.* 1985;21(4):229-34.
 38. Yang J, Li LF, Xu Q, Zhang J, Weng WW, Zhu Y-j, et al. Analysis of 90 cases of antithyroid drug-induced severe hepatotoxicity over 13 years in China. *Thyroid.* 2015;25(3):278-83. doi:10.1089/thy.2014.0350
 39. Madan R, Goyal RM, Jonklaas J, Burman, K. Methimazole Induced Fulminant Hepatic Failure. *Endocrine Society's 96th Annual Meeting and Expo, Chicago; 2014:SAT-0468.*
 40. Koch L. Therapy: Propylthiouracil use associated with severe hepatotoxicity in children. *Nat Rev Endocrinol.* 2010;6(8):416. doi:10.1038/nrendo.2010.96
 41. Levy M. Propylthiouracil hepatotoxicity. *Clin Pediatr (Phila).* 1993;32(1):25-9. doi:10.1177/000992289303200105
 42. Lock DR, Stoeber ZM. Severe hepatotoxicity on beginning propylthiouracil therapy. *J Clin Gastroenterol.* 1997;24(4):267-9. doi:10.1097/00004836-199706000-00020

43. Malozowski S, Chiesa A. Propylthiouracil-induced hepatotoxicity and death. Hopefully, Never More. *J Clin Endocrinol Metab.* 2010;95(7):3161-3. doi:10.1210/jc.2010-1141
44. Mechcatie E. Propylthiouracil use is associated with hepatotoxicity in Graves' disease. *Clinical Endocrinology News.* 2009;4:22
45. Jaeschke H, McGill MR, Ramachandran A. Oxidant stress, mitochondria, and cell death mechanisms in drug-induced liver injury: lessons learned from acetaminophen hepatotoxicity. *Drug Metab Rev.* 2012;44(1):88-106. doi:10.3109/03602532.2011.602688
46. Chan K, Truong D, Shangari N, O'Brien PJ. Drug-induced mitochondrial toxicity. *Expert Opin Drug Metab Toxicol.* 2005;1(4):655-69. doi:10.1517/17425255.1.4.655
47. Wallace KB, Starkov AA. Mitochondrial targets of drug toxicity. *Annu Rev Pharmacol Toxicol.* 2000;40(1):353-88. doi:10.1146/annurev.pharmtox.40.1.353
48. Orrenius S, Gogvadze V, Zhivotovsky B. Mitochondrial oxidative stress: Implications for Cell Death. *Annu Rev Pharmacol Toxicol.* 2007;47(1):143-83. doi:10.1146/annurev.pharmtox.47.1.20505.105122
49. Ott M, Gogvadze V, Orrenius S, Zhivotovsky B. Mitochondria, oxidative stress and cell death. *Apoptosis.* 2007;12(5):913-22. doi:10.1007/s10495-007-0756-2
50. Koch L. Therapy: Propylthiouracil use associated with severe hepatotoxicity in children. *Nat Rev Endocrinol.* 2010;6(8):416. doi:10.1038/nrendo.2010.96
51. Malozowski S, Chiesa A. Propylthiouracil-induced hepatotoxicity and death. Hopefully, never more. *J Clin Endocrinol Metab.* 2010;95(7):3161-3. doi:10.1210/jc.2010-1141
52. Sequeira E, Wanyonyi S, Dodia R. Severe propylthiouracil-induced hepatotoxicity in pregnancy managed successfully by liver transplantation: A case report. *J Med Case Rep.* 2011;5(1):461. doi:10.1186/1752-1947-5-461
53. Mete UO, Kaya M, Çolakoglu S, Polat S, Tap O, Ozbilgin MK, et al. Ultrastructure of the liver in the propylthiouracil induced hepatitis. *J Islam Acad Sci.* 1993;6(4):268-76.
54. Cashman JR, Hanzlik RP. Oxidation and other reactions of thiobenzamide derivatives of relevance to their hepatotoxicity. *J Org Chem.* 1982;47(24):4645-50. doi:10.1021/jo00145a008
55. Onderwater RCA, Commandeur JNM, Vermeulen NPE. Comparative cytotoxicity of N-substituted N'-(4-imidazole-ethyl)thiourea in precision-cut rat liver slices. *Toxicology.* 2004;197(2):80-90. doi:10.1016/j.tox.2003.11.014
56. Vollinga RC, Menge WMPB, Leurs R, Timmerman H. New analogs of burimamide as potent and selective histamine H3 receptor antagonists: the effect of chain length variation of the alkyl spacer and modifications of the N-thiourea substituent. *J Med Chem.* 1995;38(12):2244-50. doi:10.1021/jm00012a025
57. Nishida CR, Ortiz de Montellano PR. Bioactivation of antituberculosis thioamide and thiourea prodrugs by bacterial and mammalian flavin monooxygenases. *Chem Biol Interact.* 2011;192(1-2):21-5. doi:10.1016/j.cbi.2010.09.015
58. Onderwater RCA, Commandeur JNM, Groot EJ, Sitters A, Menge WMPB, Vermeulen NPE. Cytotoxicity of a series of mono- and di-substituted thiourea in freshly isolated rat hepatocytes: a preliminary structure-toxicity relationship study. *Toxicology.* 1998;125(2-3):117-29. doi:10.1016/s0300-483x(97)00169-8