



Research Article



In Vitro Evaluation of Phenolics Content, Antioxidant and Antimicrobial Activities of Three *Centaurea* L. Species from Iran

Roya Karamian^{1*}, Rose Eghbalian¹

¹Department of Biology, Faculty of Science, Bu-Ali Sina University, Hamedan, Iran.

Article Info

Article History:

Received: 22 January 2017

Accepted: 27 September 2017

ePublished: 30 December 2017

Keywords:

- Antibacterial activity
- Antioxidant
- Centaurea*
- DPPH radical scavenging
- Flavonoid
- Phenol

ABSTRACT

Background: Members of the genus *Centaurea* L. (Astraceae) are used in some countries for betterment of various ailments in the popular medicine. The main group of secondary metabolites in plants are phenolic compounds and flavonoids which are the main subset of phenolic compounds, possess a lot of biological properties such as antioxidant activity. The aim of present research is to determine *in vitro* total phenol and flavonoid contents, antioxidant potency and also antibacterial activity of the methanolic extracts of capitulla and aerial parts of three *Centaurea* species, namely *C. sosnovsky* Grossh., *C. irritans* Wagenitz and *C. kandavanesis* Wagenitz. In addition, any relationships between the phenolics content and their antioxidant properties were discussed.

Methods: To determine total phenol and flavonoid contents of the methanolic extracts, Folin-Ciocalteu and aluminum chloride methods were used, respectively. To evaluate antioxidant activity three different test systems, namely 2, 2-diphenyl-1-picrylhydrazyl free radical scavenging (DPPH), Cu-chelating and β -carotene/linoleic acid bleaching were used. In addition, disc diffusion method was used to assess antibacterial activity of the extracts against 6 gram positive and negative bacteria.

Results: Results indicated that the highest total phenol content (98.91 ± 1.87 mg GAE/g of dry extract) represented by the aerial part extract of *C. kandavanesis*. The highest total flavonoid content (5.382 ± 0.231 mg GAE/g of dry extract) and also DPPH radical scavenging activity (IC_{50} ; 0.20 ± 2.02 mg/ml) were exhibited by the capitulla extract of *C. sosnovsky*. However, the aerial part extract of *C. irritans* had the most potency for β -carotene bleaching (74.74%) among the studied samples. In addition, the extracts had acceptable antibacterial activity against both gram-positive and gram-negative bacteria tested.

Conclusion: Overall, high content of phenolic compounds and proper antioxidant activity of the examined *Centaurea* extracts may suggest them as potent antioxidants for special use in future.

Introduction

One source of medicine in virtually all cultures are medicinal plants.¹ According to WHO, plant medicines are crucial agent for the health of about 80% of the world's population, especially developing countries. Therefore, it is important to develop the research on new drugs and natural products from plants, microorganisms, fungi and animals as sources of innovative and powerful therapeutic agents for newer and affordable medicines.² In various pathological conditions such as tissue injury, inflammation process and neurodegenerative diseases, reactive oxygen species (ROS), play a vital function. However humans body is protected against ROS by antioxidants.³ ROS

compounds can easily ruin phospholipids, nucleic acids, cell membranes and polypeptides chain and play crucial role in development of different diseases.⁴ Scavenger agents like antioxidants will prevent biological systems from the free radical damages.^{5,6} Synthetic antioxidants like Butylated hydroxyl anisole (BHA) and butylated hydroxyl toluene (BHT) were previously used in food industries, but because of their toxic effects on human body using of BHA and BHT are recently restricted.⁷ Tocopherols and tocotrienols, Fat-soluble antioxidants (vitamin E), folic acid (vitamin B9), L-ascorbic acid (vitamin C), flavonoids, carotenoids, phenyl acrylic acids and phenyl methanoic acids are the important antioxidants

*Corresponding Author: Roya Karamian, E-mail: R_karamian@basu.ac.ir

©2017 The Authors. This is an open access article and applies the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

which exist in plants.⁸ The genus *Centaurea* L. belongs to the family Asteraceae and includes 400-700 herbaceous thistle-like flowering species are mainly distributed in Middle East. Ethnopharmacological studies have shown that many *Centaurea* species are used for treatment of diverse ailments in the popular medicine in some countries. For example, in Scottish medicine, *Centaurea cyanus* and *Centaurea scabiosa* are used as diuretic agents.⁹ However; there are few reports to previous works on biological activity of other *Centaurea* species. The goal of current study is to evaluate *in vitro* antioxidant properties and total phenolics content of the methanolic extracts of three *Centaurea* species (*C. sosnovsky*, *C. irritans* and *C. kandavanensis*) in both capitulla and aerial parts. Besides, correlation among total phenol and flavonoid contents and antioxidant properties is determined.

Materials and Methods

Plant materials

Three *Centaurea* species (*Centaurea sosnovsky* Grossh., *Centaurea irritans* Wagenitz and *Centaurea kandavanensis* Wagenitz) were collected from their natural distribution areas; then their aerial and capitulla parts were dissected (Table 1). The Voucher pattern of the species representing these collections have been deposited at the Bu Ali Sina University Herbarium (BASU), Hamedan, Iran.

Extract preparation

Briefly, powdered parts of aerial and capitulla of the species (25 g) were extracted by 250 ml absolute methanol for 12 hat 50 °C using a Soxhlet apparatus and solvents were removed under the vacuum evaporator (Lab Tech, Ev 311, Italy) at temperature below 60 °C to yield a waxy material. Finally, the extracts were lyophilized and kept in 4 °C until used.

Total phenol content estimation

Folin-Ciocalteu reagent was used to determine total phenol content.¹⁰ In brief, 0.5 ml each of the extract (1:10 g/ml) or gallic acid was admixed with 5 ml Folin-Ciocalteu reagent (1:10 distilled water) and 4 ml aqueous Na₂CO₃ (1 M). After 15 min at room temperature the total phenol content of the mixture was determined by colorimetric method at 765 nm (Perkin Elmer UV/visible spectrophotometer, USA). The calibration curve ($y = 0.002x + 0.0177$,

$R^2 = 0.978$) was prepared using gallic acid solution at different concentrations of 0-250 µg/ml in 50% methanol. Total phenol content was calculated from calibration curve and expressed as gallic acid equivalent (mg GAE/g).

Total flavonoid content Estimation

The AlCl₃ method was used for estimating total flavonoid content.¹¹ Briefly, 0.5 ml each of extract (1:10 g/ml) in methanol were mixed with 1.5 ml methanol, 0.1 ml AlCl₃ (10%), 0.1 ml KCH₃COO (1 M) and 2.8 ml distilled water and left at room temperature for 30 min. Absorbance of the mixtures was measured at 415 nm (Perkin Elmer UV/visible spectrophotometer, USA). The calibration curve ($y = 0.0091x + 0.0206$, $R^2 = 0.995$) was prepared using quercetin solution at different concentrations of 12.5-100 µg/ml in methanol. Then, the total flavonoid content was calculated from calibration curve and expressed as quercetin equivalent (mg QE/g extract).

Evaluation of antioxidant activity

Free radical scavenging

The method of Mensor et al was used for measuring free radical scavenging activity of the extracts.¹² In order to obtain dilutions, different extract concentrations were prepared in methanol (0.2-1 mg/ml) and 2.5 ml each of concentration was added to 1 ml DPPH alcohol solution (0.3 mM). After completion the samples were first kept in a dark place at room temperature for 30 min and bleaching of DPPH was read at 517 nm against a blank (methanol). The inhibition in percent for each concentration was calculated according to the following formula:

$$\begin{aligned} \text{DPPH free radical scavenging}(\%) \\ = [1 - (As - Ab)/Ac] * 100 \end{aligned} \quad \text{Eq.(1)}$$

Where, As is absorbance of the reaction mixture containing 2.5 ml of extracts + 1 ml DPPH, Ab is absorbance of the reaction mixture containing 2.5 ml of extracts + 1 ml methanol and Ac is absorbance of control sample containing 1 ml DPPH + 2.5 ml methanol. Also IC₅₀ value, which represented the concentration of the sample that caused 50% inhibition, was determined. Tests were carried out in triplicate and the positive control was ascorbic acid.

Table 1. The collections of investigated *Centaurea* species.

Species	Voucher number/locality	Date	Altitude (m)
<i>C. sosnovsky</i>	BASU 28813/Lorestan	10.5.2015	2100
<i>C. irritans</i>	BASU 28814/Kermanshah	10.5.2015	2100
<i>C. kandavanensis</i>	BASU 29197/Mazandaran	15.7.2015	2140

Cupric ion reducing antioxidant capacity (CUPRAC assay)

The cupric ion reducing capacity was measured due to the method of Apak et al.¹³ 1 ml CuCl₂ (10 mM), 1 ml neocuproine (7.5 mM), and 1 ml NH₄AC buffer (1M, pH 7.0) solutions were added into a test tube. Then, 0.5 ml of extract at different concentrations was mixed and total volume was brought up to 4.1 ml with deionized water. The mixture absorbance was recorded against a blank at 450 nm after 30 min incubation at room temperature.

β-Carotene linoleic acid bleaching

According to the method of Barriere et al, antioxidant capacity is assessed by measuring the inhibition of the volatile organic compounds and the conjugated dienehydroperoxides arising from linoleic acid oxidation.¹⁴ A stock solution of β-carotene/linoleic acid was prepared as follows: first, 0.5 mg β-carotene was dissolved in 1 ml chloroform, then 25 μl linoleic acid and 200 mg Tween 40 were added. The chloroform was subsequently evaporated using a vacuum evaporator. Then 100 ml distilled water saturated with oxygen (30 min at 100 ml/min) were added with vigorous shaking. Aliquots (2.5 ml) of this reaction mixture were transferred to test tubes, and 350 μl portions of the extracts (2 g/l in ethanol) were added before incubating for 48 h at room temperature. The same procedure was repeated with BHT at the same concentration and a blank containing only 350 μl ethanol. After the incubation period, the absorbance of the mixtures was measured at 490 nm. Antioxidant capacity of the samples was compared with that of BHT and blank.

Antibacterial activity assay

The plant extracts were dissolved in DMSO to a final concentration of 2 mg/ml. Antibacterial activity of the extracts at 4 different concentrations (100, 50, 25 and 12.5 mg/ml of extract) was measured against 6 gram positive and negative bacteria, namely *Serratia marcescens* (PTCC 1111),

Enterobacter aerogenes (PTCC 1221), *Salmonella enteric* (PTCC 1236), *Bacillus thuringiensis* (Wild), *Staphylococcus aureus* (Wild) and *Bacillus megaterium* (PTCC 1017) by disc diffusion method.¹⁵ The extracts were dissolved in DMSO to make a 100 mg/ml solution and other concentrations make from this concentration, and then apply on the blank paper discs. Dried discs were placed onto Muller-Hinton agar medium that previously inoculated with a bacterial suspension (1.5 × 10⁸ of bacteria/ml). The cultures were incubated at 37 °C for 24 h. The antibacterial activity against each test organism was quantified by determining mean zone of inhibition. Gentamycin, Penicillin, Nitroflantoin and Neomycin were also used as positive controls.

Statistical analysis

All data are the average of triplicate analyses. Statistical analysis of variance was performed using Student's t-test by SPSS program and *p* value < 0.05 was regarded as significant. Data are expressed as means ± standard deviation.

Results

Determination of total phenol and flavonoid contents

Results proved that the content of total phenols was different in the studied species (from 19.34 ± 3.83 to 98.91 ± 1.87 mg GAE/g of dry extract). The highest total phenol content was found for the aerial part extract of *C. kandavanensis* and the lowest one for the capitulla extract of *C. irritans*. The capitulla extract of *C. sosnovsky* had the highest flavonoid content (Table 2).

DPPH Free radical scavenging

Results from assessment of free radical scavenging capacity of the corresponding extracts are shown in Table 2 and Figure 1. All extracts showed a weaker potency than ascorbic acid for scavenging DPPH free radicals. A lower IC₅₀ represented the higher antioxidant activity.

Table 2. Total phenol and flavonoid contents and DPPH radical scavenging activity of the *Centaurea* species.

Extract	Total phenol (mg/g dw)	Total flavonoids (mg/g dw)	β-Carotene/linoleic acid	DPPH radical scavenging	
				IC ₅₀ (mg/ml)	(%)
<i>C. sosnovsky</i> (aerial)	56.33 ^e ± 1.89	4.558 ^d ± 0.467	49.19 ^{bc} ± 2.55	0.75 ^g ± 2.33	44.40
<i>C. sosnovsky</i> (capitulla)	69.85 ^{de} ± 7.10	5.382 ^b ± 0.231	47.50 ^{bc} ± 3.97	0.2 ^b ± 2.02	78.26
<i>C. irritans</i> (aerial)	86.05 ^{bc} ± 5.26	2.410 ^e ± 0.065	74.74 ^a ± 4.85	0.29 ^{bc} ± 2.9	80.46
<i>C. irritans</i> (capitulla)	19.34 ^f ± 3.83	1.413 ^f ± 0.011	58.76 ^b ± 4.33	0.39 ^{cd} ± 0.88	67.80
<i>C. kandavanensis</i> (aerial)	98.91 ^{ab} ± 1.87	5.083 ^{bcd} ± 0.231	49.07 ^{bc} ± 6.05	0.33 ^c ± 2.95	77.43
<i>C. kandavanensis</i> (capitulla)	66.35 ^e ± 7.12	5.240 ^{bc} ± 0.338	50.95 ^{bc} ± 8.66	0.44 ^d ± 3.2	65.91
Ascorbic acid	-	-	-	0.125 ^a ± 0.8	80.61

Experiment was performed in triplicate and expressed as mean ± SD. Values in each column with different superscripts are significantly different (*P* < 0.05).

The following descending order showed the effectiveness of extracts as DPPH radical scavengers: ascorbic acid (IC_{50} ; 0.143 ± 0.007 mg/ml) < *C. sosnovsky* (capitulla) (IC_{50} ; 0.2 ± 2.02 mg/ml) < *C. irritans* (aerial parts) (IC_{50} ; 0.21 ± 2.09 mg/ml) < *C. kandavanensis* (aerial parts) (IC_{50} ; 0.333 ± 2.95 mg/ml) < *C. irritans* (capitulla) (IC_{50} ; 0.392 ± 0.88 mg/ml) < *C. kandavanensis* (capitulla) (IC_{50} ; 0.444 ± 3.2 mg/ml) < *C. sosnovsky* (aerial parts) (IC_{50} ; 0.754 ± 2.33 mg/ml (Table 2).




Figure 1. Ability of DPPH radical scavenging of the studied *Centaurea* species (S.A: *C. sosnovsky* aerial, S.C: *C. sosnovsky* capitulla, I.A: *C. irritans* aerial, I.C: *C. irritans* capitulla, K.A: *C. kandavanensis* aerial, K.C: *C. kandavanensis* capitulla).

Cupric ion reducing antioxidant capacity

Results from Cu chelating assay showed that the absorbance of extracts linearly depended on concentration (0.2 to 01 mg/ml) and increased by increasing concentration. The aerial part extract of *C. irritans* and capitulla extract of *C. sosnovsky* displayed the highest absorbance values (Figure 2).




Figure 2. Cupric reducing antioxidant capacity of the studied *Centaurea* species at different concentrations (S.A: *C. sosnovsky* aerial, S.C: *C. sosnovsky* capitulla, I.A: *C. irritans* aerial, I.C: *C. irritans* capitulla, K.A: *C. kandavanensis* aerial, K.C: *C. kandavanensis* capitulla).

β -Carotene linoleic acid bleaching

Results from antioxidant activity using β -carotene/linoleic acid model system indicated that the aerial parts extract of *C. irritans* was more active

($74.74 \pm 4.85\%$) than other samples. However, no substantial difference ($P < 0.05$) was observed between other evaluated samples (Table 3).

Table 3. Inhibition percentages of the linoleic acid oxidation by two parts of three *Centaurea* species and BHT.

Extracts	Inhibition (%)
<i>C. sosnovsky</i> (aerial)	49.19bc \pm 2.55
<i>C. sosnovsky</i> (capitulla)	47.50bc \pm 3.97
<i>C. irritans</i> (aerial)	74.74a \pm 4.85
<i>C. irritans</i> (capitulla)	58.76 b \pm 4.33
<i>C. kandavanensis</i> (aerial)	49.07bc \pm 6.05
<i>C. kandavanensis</i> (capitulla)	50.95bc \pm 8.66
BHT	97.29a \pm 1.69
Control -	13.39d \pm 0.44

Experiment was performed in triplicate and expressed as mean \pm SD. Values in each column with different superscripts are significantly different ($P < 0.05$).

Antibacterial activity assay

Results from antibacterial assay indicated that *C. sosnovsky* extract in both parts have strong antibacterial capacity against 6 gram-positive and negative bacteria. Then again, the aerial and capitulla parts of *C. kandavanensis* and also the capitulla part of *C. irritans* showed the weakest antibacterial function versus gram negative bacteria tested. The aerial part extract of *C. irritans* had no effect on *Bacillus thuringiensis* a gram positive bacterium. *Bacillus megaterium* and *Staphylococcus aureus* had the highest sensitivity to the studied extracts (Tables 4-9).

Discussion

Phenolic compounds which act as free radical terminators, are a class of antioxidant agents, and their bioactivities may be related to their capabilities to chelate metals, inhibit lipoxygenase and scavenge free radicals.^{16,17} Activity of multiple cytochrome P₄₅₀ isoforms, lipoxygenases, cyclooxygenase and xanthine oxidase can strongly inhibited by phenolic structures. They also play an important role in plant defense mechanisms to counteract reactive oxygen and nitrogen (ROS and RNS) and inhibit lipid peroxidation.¹⁸⁻²⁰ Antioxidant agents of phenolic compounds are hydroxyl groups that are power hydrogen donors. Flavonoids are big group of antioxidant, which have been classified into seven sub groups: flavanols, flavones, flavanones, isoflavones, catechins, anthocyanins and proanthocyanidins. Because of their potent antioxidant ability, flavonoids have positive effects on human health.²¹ In current research, all extracts indicated a high total phenolic content. The highest content was observed in the aerial part of *C. kandavanensis* (98.91 ± 1.87 mg GAE/g of dry extract) and the lowest one in the capitulla of *C. irritans* (19.34 ± 3.83 mg GAE/g of dry extract).

Table 4. Average inhibition zone of evaluated bacterial strains against aerial parts extract of *C. sosnovskyi*.

Bacterial species	Inhibition zone (mm)				
	Concentration (mg/ml)				
	100	50	25	12.5	DMSO
<i>Bacillus megaterium</i>	7.83±5.3	7.33±2.5	7.33±1.58	7.67±2.33	NA
<i>Staphylococcus aureus</i>	NA	9.00±1.65	8.67±1.11	14.67±1.00	NA
<i>Serratia marcescens</i>	8.51±2.13	8.33±1.51	10.00±1.98	10.17±1.00	NA
<i>Enterobacter aerogenes</i>	6.83±2.65	7.00±2.47	7.33±1.87	8.33±1.15	NA
<i>Salmonella enterica</i>	10.56±1.65	10.00±1.45	8.33±1.65	7.67±2.00	NA
<i>Bacillus thuringiensis</i>	7.00±2.11	8.00±1.9	11.33±1.15	11.17±1.14	NA

NA, No Active.

Experiment was performed in triplicate and expressed as mean ± SD.

Table 5. Average inhibition zone of evaluated bacterial strains against capitulla extract of *C. sosnovskyi*.

Bacterial species	Inhibition zone (mm)				
	Concentration (mg/ml)				
	100	50	25	12.5	DMSO
<i>Bacillus megaterium</i>	7.87±1.22	7.33±0.85	7.33±1.514	7.78±1.45	NA
<i>Staphylococcus aureus</i>	8.33±1.85	9.40±1.50	8.93±1.30	13.67±1.54	NA
<i>Serratia marcescens</i>	7.51±1.45	7.33±1.12	9.00±1.80	9.17±1.78	NA
<i>Enterobacter aerogenes</i>	6.50±2.50	7.00±2.10	7.83±1.21	8.38±1.69	NA
<i>Salmonella enterica</i>	9.56±1.74	10.04±1.31	8.33±1.66	7.00±1.00	NA
<i>Bacillus thuringiensis</i>	6.00±2.00	7.30±1.05	9.53±0.95	10.11±1.81	NA

NA, No Active.

Experiment was performed in triplicate and expressed as mean ± SD.

Table 6. Average inhibition zone of evaluated bacterial strains against aerial parts extract of *C. irritans*.

Bacterial species	Inhibition zone (mm)				
	Concentration (mg/ml)				
	100	50	25	12.5	DMSO
<i>Bacillus megaterium</i>	6.51±1.8	6.87±1.63	7.43±1.12	7.88±2.00	NA
<i>Staphylococcus aureus</i>	5.65±2.12	5.83±2.68	6.70±2.22	7.33±2.5	NA
<i>Serratia marcescens</i>	NA	NA	NA	NA	NA
<i>Enterobacter aerogenes</i>	NA	NA	NA	NA	NA
<i>Salmonella enterica</i>	NA	NA	NA	NA	NA
<i>Bacillus thuringiensis</i>	NA	NA	NA	NA	NA

NA, No Active.

Experiment was performed in triplicate and expressed as mean ± SD.

Table 7. Average inhibition zone of evaluated bacterial strains against capitulla extract of *C. irritans*.

Bacterial species	Inhibition zone (mm)				
	Concentration (mg/ml)				
	100	50	25	12.5	DMSO
<i>Bacillus megaterium</i>	NA	NA	7.00±1.58	7.50±2.15	NA
<i>Staphylococcus aureus</i>	7.00±2.54	8.17±2.15	10.83±1.50	10.88±2.85	NA
<i>Serratia marcescens</i>	7.17±1.50	7.50±3.58	9.83±1.12	8.83±2.20	NA
<i>Enterobacter aerogenes</i>	NA	6.83±2.8	7.83±2.18	12.00±2.50	NA
<i>Salmonella enterica</i>	NA	NA	7.33±1.00	8.67±1.98	NA
<i>Bacillus thuringiensis</i>	7.17±0.55	8.83±1.20	9.33±0.54	14.33±1.50	NA

NA, No Active.

Experiment was performed in triplicate and expressed as mean ± SD.

Table 8. Average inhibition zone of evaluated bacterial strains against aerial parts extract of *C. kandavanensis*.

Bacterial species	Inhibition zone (mm)				
	Concentration (mg/ml)				
	100	50	25	12.5	DMSO
<i>Bacillus megaterium</i>	13.83±1.15	9.33±2.14	6.83±2.50	6.67±2.80	NA
<i>Staphylococcus aureus</i>	8.67±1.33	5.83±2.68	6.70±2.22	7.33±2.5	NA
<i>Serratia marcescens</i>	NA	NA	NA	NA	NA
<i>Enterobacter aerogenes</i>	NA	NA	NA	NA	NA
<i>Salmonella enterica</i>	NA	NA	NA	NA	NA
<i>Bacillus thuringiensis</i>	7.00±1.20	8.17±1.45	8.83±1.6	6.67±1.32	NA

NA, No Active.

Experiment was performed in triplicate and expressed as mean ± SD.

Table 9. Average inhibition zone of evaluated bacterial strains against capitulla extract of *C. kandavanensis*.

Bacterial species	Inhibition zone (mm)				
	Concentration (mg/ml)				
	100	50	25	12.5	DMSO
<i>Bacillus megaterium</i>	12.33±1.12	11.60±1.61	9.00±1.00	8.54±1.54	NA
<i>Staphylococcus aureus</i>	9.34±1.10	6.15±1.10	6.98±2.80	7.50±2.65	NA
<i>Serratia marcescens</i>	NA	NA	NA	NA	NA
<i>Enterobacter aerogenes</i>	NA	NA	NA	NA	NA
<i>Salmonella enterica</i>	NA	NA	NA	NA	NA
<i>Bacillus thuringiensis</i>	7.50±1.50	7.50±1.70	8.00±1.81	8.00±1.20	NA

NA, No Active.

Experiment was performed in triplicate and expressed as mean ± SD.

In another study the phenolic content of some *Centaurea* species, which was assessed varied from 82.273 mg GAE/g in *C. ptosimopappoides* to 175.404 mg GAE/g in *C. cheirollopha*.²² It is popular that phenolic compounds providing health beneficial effects by modifying color, taste, aroma, and flavor. They also play a crucial role in protecting cells from damages of microorganisms, insects, and herbivores.²³ Our results indicated that *C. kandavanensis* in both of aerial and capitulla parts and *C. sosnovsky* in the capitulla part were not notable different ($P < 0.05$) in total flavonoid content. However, the aerial parts of *C. sosnovsky* represented a higher amount of total flavonoids (5.382 ± 0.231 mg QES/g of dry extract). The capitulla extract of *C. irritans* had the lowest content of total flavonoids in comparison with other samples.

Due to the injurious role of free radicals in foods and biological systems, radical scavenging activity is very important.²⁴ Free radicals are generated as byproducts of biological reactions. *In vitro* antioxidant content of the phenolic compounds on a molar basis is more than vitamins E and C.²⁵ DPPH scavenging activity of the extracts and ascorbic acid as a synthetic antioxidant was assessed by using IC₅₀. There is no significant announcement to previous works on the antioxidant activity (as DPPH free radical scavenging) of *C. sosnovsky*, *C. irritans* and *C. kandavanensis*. The studied extracts

here showed more amount of IC₅₀ than ascorbic acid that means their lower antioxidant activity. The capitulla extract of *C. sosnovsky* and the aerial part of *C. irritans* represented the lowest IC₅₀ values. To the best of our knowledge, several studies were fulfilled on radical scavenging activity of some members of the genus *Centaurea*, such as *C. cyanus*, *C. scabiosa*,²⁶ *C. calolepis* and *C. cadmea*.^{27,28} However, the methods for DPPH assay, which were used in these studies was different, so their results are no directly comparable to ours.

Many reports revealed that there is a direct interaction between antioxidant activity and reducing power of plant extract.^{29,30} Cupric reducing capacity procedure, is depended on the assessment of absorbance at 450nm by the organization of the reliable complex between neocuproine and copper(I). The latter is formed by the reduction of copper(II) in the presence of neocuproine.¹³ According to our results, there was no relative relevance between cupric reducing capacity of the extracts and total phenol or flavonoid content. The aerial part extract of *C. irritans* and the capitulla extract of *C. sosnovsky* represented the highest absorbance and the lowest IC₅₀ values in DPPH scavenging activity.

Discoloration of β-carotene by antioxidants is the base of β-Carotene/linoleic acid test system. the extent of β-carotene degradation can be hindered by neutralizing the linoleate free radical and any other

free radicals formed within the system.³¹ The aerial parts extract of *C. irritans* was effectively inhibited the oxidation of linoleic acid ($74.74 \pm 4.85\%$), followed by the capitulla extract of *C. irritans* ($58.76 \pm 4.33\%$). In addition, *C. kandavanensis* and *C. sosnovsky* in both of aerial and capitulla parts were not remarkable distinct ($P < 0.05$) (Table 2). There was a moderate correlation ($R^2 > 0.56$) between total flavonoid amount of the extracts and β -carotene bleaching inhibition potential, offering that up to 56% of β -carotene bleaching inhibition can be related to flavonoid compounds. Based on our results the studied *Centaurea* extracts had high flavonoid content and represented good antioxidant activity.

Our outcomes demonstrated that all tested bacteria the aerial parts extract of *C. sosnovsky* had a strong antibacterial activity against all tested bacteria. Also, the aerial and capitulla extracts of *C. kandavanensis* showed the weakest antibacterial function chiefly versus gram negative bacteria. Our results showed that the studied extracts have high antibacterial function especially against gram positive bacteria that is in parallel with Cansaran et al.³² It could be due to the presence of multilayer membrane, which covered each gram-negative bacteria cell, whereas, gram-positive bacteria consist of a single layer. It is the reason why gram-negative bacteria are more resistant to antibacterial than gram-positive bacteria.³³

Conclusion

Our current research prepared usable knowledge on phenolic content and antioxidant activity of three *Centaurea* species by ordinary, simple and reliable protocols. Our observations mentioned that difference in the antioxidant activity between the extracts cannot adequately describe by quantitative difference in their total phenolic contents. Hence, the methanolic extracts of *Centaurea* species contain major phytochemicals that may be candidates to develop new drugs, which might be practical in betterment of diseases caused by different free radicals. However, more researches are needed to figure out the *in vivo* antioxidant potential of these extracts in animal models.

Acknowledgments

The authors would like to thank Bu-Ali Sina University for financial support of this work.

Conflict of interests

The authors claim that there is no conflict of interest.

References

1. Baquar SR. The role of traditional medicine in rural environment. In: Issaq S editor. Traditional Medicine in Africa. Nairobi: East Africa Educational Publishers Ltd; 1995.

2. Cooper EL. Drug discovery, CAM and natural products. Evid Based Complement Alternat Med. 2004;1(3):215-7.
3. Aruoma OI, Cuppette SL. Antioxidant methodology: *in vivo* and *in vitro* concept. Champaign: IL, AOCS Press; 1997. p. 142-69.
4. Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. Int J Biochem Cell Biol. 2007;39(1):44-84. doi:10.1016/j.biocel.2006.07.001
5. Zhang HY, Yang DP, Tang GY. Multipotent antioxidants: from screening to design. Drug Discov Today. 2006;11(15-16):749-54. doi:10.1016/j.drudis.2006.06.007
6. Yu L, Perret J, Davy B, Wilson J, Melby CL. Antioxidant properties of cereal products. J Food Sci. 2002;67(7):2600-3. doi:10.1111/j.1365-2621.2002.tb08784.x
7. Prior RL, Wu X, Schaich K. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. J Agric Food Chem. 2005;53(10):4290-302. doi:10.1021/jf0502698
8. Sun L, Zhang J, Lu X, Zhang L, Zhang Y. Evaluation to the antioxidant activity of total flavonoids extract from persimmon (*Diospyros kaki* L.) leaves. Food Chem Toxicol. 2011;49(10):2689-96. doi:10.1016/j.fct.2011.07.042
9. Kumarasamy Y, Byres M, Cox PJ, Jaspars M, Nahar L, Sarker SD. Screening seed of some Scottish plants for free radical scavenging activity. Phytother Res. 2007;21(7):615-21. doi:10.1002/ptr.2129
10. McDonald S, Prenzler PD, Autolovich M, Robards K. Phenolic content and antioxidant activity of olive extracts. Food Chem. 2001;73(1):73-84. doi:10.1016/S0308-8146(00)00288-0
11. Chang C, Yang M, Wen H, Chern J. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. J Food Drug Anal. 2002;10(3):178-82.
12. Mensor L, Menezes FS, Leitao GG, Reis AS, Santos TS, Coube CS. Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method. Phytother Res. 2001;15(2):127-30. doi:10.1002/ptr.687
13. Apak R, Guclu K, Ozyurek M, Karademir SE, Ercag E. The cupric ion reducing antioxidant capacity and polyphenolic content of some herbal teas. Int J Food Sci Nutr. 2006;57(5-6):292-304. doi:10.1080/09637480600798132
14. Barriere C, Centeno D, Lebert A, Leroy Setrin S, Berdague JL, Talon R, et al. Roles of superoxide dismutase and catalase of *Staphylococcus xylosum* in the inhibition of linoleic acid

- oxidation. FEMS Microbiol Lett. 2001;201(2):181-5. doi:10.1016/s0378-1097(01)00271-3
15. Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by single disk method. Am J Clin Pathol. 1966;45(4):493-6.
 16. Kahkonen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala TS, et al. Antioxidant activity of plant extracts containing phenolic compounds. J Agric Food Chem. 1999;47(10):3954-62. doi:10.1021/jf990146l
 17. Rice Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. Free Radic Biol Med. 1996;20(7):933-56. doi:10.1016/0891-5849(95)02227-9
 18. Shahidi F, Wanasundara PK. Phenolic antioxidants. Critic Rev Food Sci Nutr. 1992;32(1):67-103. doi:10.1080/10408399209527581
 19. Valentao P, Fernandes E, Carvalho F, Andrade PB, Seabra RM, Bastos ML. Hydroxyl radical and hypochlorous acid scavenging activity of small centaury (*Centaureum erythraea*) infusion. A comparative study with green tea (*Camellia sinensis*). Phytomedicine. 2003;10(6-7):517-22. doi:10.1078/094471103322331485
 20. Parr AJ, Bolwell JP. Phenols in the plant and in man. The potential for possible nutritional enhancement of the diet by modifying the phenols content or profile. J Sci Food Agric. 2002;80(7):985-1012. doi:10.1002/(sici)1097-0010(20000515)80:7<985::aid-jsfa572>3.0.co;2-7
 21. Miller AL. Antioxidant Flavonoids: Structure, Function and Clinical Usage. Altern Med Rev. 1996; 1(2): 103-11.
 22. Karamenderes C, Khan S, Tekwani BL, Jacop MR, Khan IA. Antiprotozoal and antimicrobial activities of *Centaurea* species growing in Turkey. Pharm Biol. 2006;44(7):534-9. doi:10.1080/13880200600883080
 23. Vaya J, Belinky PA, Aviram M. Antioxidant constituents from licorice roots: Isolation, structure elucidation and antioxidative capacity toward LDL oxidation. Free Radic Biol Med. 1997;23(2): 302-13. doi:10.1016/s0891-5849(97)00089-0
 24. Peng Y, Ye J, Kong J. Determination of Phenolic Compounds in *Perilla frutescens* L. by Capillary Electrophoresis with Electrochemical Detection. J Agric Food Chem. 2005;53(21):8141-7. doi:10.1021/jf051360e
 25. Rice Evans C, Miller N, Paganga G. Antioxidant properties of phenolic compounds. Trends Plant Sci. 1997;2(4):152-9. doi:10.1016/s1360-1385(97)01018-2
 26. Kumarasamy Y, Byres M, Cox PJ, Jaspars M, Nahar L, Sarker SD. Screening seed of some Scottish plants for free radical scavenging activity. Phytother Res. 2007;21(7): 615-21. doi:10.1002/ptr.2129
 27. Karamenderes C, Bedir E, Pawar R, Baykan S, Khan IA. Elemanolide sesquiterpenes and eudesmane sesquiterpene glycosides from *Centaurea hierapolitana*. Phytochemistry. 2007;68(5):609-15. doi:10.1016/j.phytochem.2006.10.013
 28. Karamenderes C, Konyalioglu S, Khan S, Khan IA. Total phenolic contents, free radical scavenging activities and inhibitory effects on the activation of NF-kappa B of eight *Centaurea* L. species. Phytother Res. 2007;21(5):488-91. doi:10.1002/ptr.2097
 29. Yildirim A, Mavi A, Kara AA. Determination of antioxidant and antimicrobial activities of *Rumex crispus* L., Extracts. J Agric Food Chem. 2001;49(8):4083-9. doi:10.1021/jf0103572
 30. Koleva II, Van Beek TA, Linszen JPH, De Groot A, Evstaieva LN. Screening of plants extracts for antioxidant activity: a comparative study on three testing methods. Phytochem Anal. 2002;13(1): 8-17. doi:10.1002/pca.611
 31. Jayaprakasha GK, Singh RP, Sakariah KK. Antioxidant activity of grape seed (*Vitis vinifera*). Food Chem. 2001;73(3):285-90. doi:10.1016/s0308-8146(00)00298-3
 32. Ali Shtayeh MS, Yaghmour RM, Faidi YR, Salem K, Al Nuri MA. Antimicrobial activity of 20 plants used in folkloric medicine in the palestinian area. J Ethnopharmacol. 1998;60(3):265-71. doi:10.1016/s0378-8741(97)00153-0
 33. Cansaran A, Mercan Dogan N, Öztekin M, Acar G. Antimicrobial activity of various extracts of *Centaurea cankiriense* A. Duran and H. Duman. Afr J Microbiol Res. 2010; 4(8): 608-12