



Chemical Composition and Antibacterial Activity of the Essential Oil of *Zosimia absinthifolia* Growing in East Azarbaijan (Iran)

Fariba Heshmati Afshar^{1,2}, Mohammad Bakhshandeh², Masood Mohamadzadeh², Solmaz Asnaashari³, Laleh Khodaie^{1,4*}

¹Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

²Department of Pharmacognosy, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

³Biotechnology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

⁴Department of Phytopharmacy, Faculty of Traditional Medicine, Tabriz University of Medical Sciences, Tabriz, Iran.

Article Info

Article History:

Received: 17 January 2017

Accepted: 14 June 2017

ePublished: 30 December 2017

Keywords:

- Antibacterial
- Apiaceae
- Disc diffusion method
- Essential oil
- Zosimia absinthifolia*

ABSTRACT

Background: *Z. absinthifolia*, with local name of “Zarak-e-Kuhi” has been consumed as a spice in some regions of Iran.

Methods: In this study, the essential oil obtained from the aerial parts of *Zosimia absinthifolia* (Apiaceae) (ZaeM) was analyzed by GC-MS. Also, the chemical profile of the oil was compared with the same species collected from different localities. Moreover, the essential oil was investigated for its antimicrobial activity by disc diffusion method.

Results: Analysis of ZaeM by GC-MS resulted nineteen components consisting 98.22% of the total oil. The essential oil was predominantly made up of esters (68.48%) and alcohols (26.04%). The most abundant components of the oil were octyl acetate (47.29%), n-octanol (25.79%), octyl butyrate (10.15%) and octanoic acid octyl ester (7.9%). The relative amounts of the main compounds of *Z. absinthifolia* essential oils growing in different places were comparable with each other. Both qualitative and quantitative variability of chemical composition of *Z. absinthifolia* essential oils were possibly related to the individual genotypes or geographical origin. Disc diffusion method was employed for the determination of antimicrobial activity of ZaeM. The results showed that the essential oil just had inhibitory activity against *B. subtilis*, a gram positive strain. The inhibitory activity could be attributed to the long chain alcohol, octanol (25.79%), which had previously shown antimicrobial activity.

Conclusion: As a conclusion, ZaeM could have antimicrobial potentials on *B. subtilis*. Further investigations are needed to isolate and identify antimicrobial compounds from this volatile oil.

Introduction

The family Apiaceae (Umbelliferae) consists of 150 genera and 3000 species largely exist in temperate regions.¹ The genus *Zosimia* includes two species in Iran: *Zosimia radianse* and *Zosimia absinthifolia*.² The seeds of *Z. absinthifolia*, with local name of “Zarak-e-Kuhi” have been consumed as spice in some regions of Iran.³ The reported phyto constituents from methanolic extract of roots and fruits of *Z. absinthifolia*, were lactones of coumarin group, zosimin and deltoin. Moreover, two flavonoids as well as one chalcone have been identified from the methanolic extract of aerial parts of this species.⁴ In another study, three flavonoids were separated from the methanolic extract of seeds of *Z. absinthifolia*.³ Furthermore, GC and GC/MS

analysis of the n- hexan extract from the seeds of this herb resulted n-octylacetate, octylpropanoate, decane, angelicin, umbelliferone as the major chemical compounds. Since Apiaceae (Umbelliferae) was selected as the source of essential oil containing herbs, so the volatile oil of the genus *Zosimia* would be a good candidate to investigate both in terms of the composition and biological activities. In the previous manuscript the essential oil composition of *Z. absinthifolia* collected from different geographical locations has been reported.⁵⁻⁷ Even though the chemical composition of *Z. absinthifolia* essential oil (Zae) has been previously reported, in this case, due to the effect of harvesting time and season, geographical location, altitude and climate on the yield and

*Corresponding Author: Laleh Khodaie, E-mail: khodaie@gmail.com

©2017 The Authors. This is an open access article and applies the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

composition of volatile oils,⁸⁻⁹ this study aimed to determine firstly, the chemical composition of Zae growing in Mishoodaghi Mount in East-Azərbayjan Province of Iran (Zae M); secondly, to compare the main phytoconstituents of Zae M with Zae growing in the different regions with the different conditions. To the best of our knowledge there was no report on antibacterial activity of this herb so this paper will focus on the determination of antibacterial activity of ZaeM by disc diffusion method as well as the discussion of the connection between antibacterial activity and Zae phytoconstituents.

Materials and Methods

Plant Material

Aerial parts of *Zosimia absinthifolia* (Vent.) Link were collected from a place near Marand (Misho mountain) at E: 45° 47', N: 38° 19', altitude of 2036 in East-Azərbayjan Province, Iran in June 2014. Voucher specimen was deposited in the herbarium of Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

Essential oil Extraction

The fresh aerial parts of *Z. absintifolia* were shade-dried at the room temperature for 10 days. Then dried aerial parts of the herb (90 g) were subjected to hydrodistillation using a Clevenger-type apparatus) for 4 h. Then the oil was dried over anhydrous sodium sulfate, measured and stored in a dark glass at 4 °C for further studies.

GC-MS and GC-FID Analysis

The volatile oil was analyzed using a Shimadzu GCMS-QP5050A Gas Chromatography Mass Spectrometer (GC-MS) fitted with a fused silica DB-1 capillary column (60 m × 0.25 mm i.d., 0.25 μm film thickness). Helium was used as the carrier gas at a flow rate of 1.3 ml/min. The oven temperature was kept at 50°C for 3 min, and programmed to rise to 270°C at a rate of 5°C/min and then kept constant for 4 minutes. The injector temperature was 250°C and split ratio was adjusted at 1:24. The Mass Spectral (MS) data obtained at the following conditions; ionization potential 70 eV; ion source temperature 200°C; quadrupole temperature 100°C; solvent delay 3 minutes and EM voltage 3000 volts. Identification of the essential oil components was performed by the comparison of their mass spectra with the NIST 21, NIST 107 and WILEY229 libraries, the spectral data banks, which make computer matching available. The comparison of the fragmentation patterns of mass spectra of unknown compounds with the reported components was carried out as well. Kovats indices (KI) of the components were calculated using standard n-alkanes (C₈-C₂₀) injection, under the same chromatographic conditions.

For the quantitation (area %), the GC analyses were

performed on a Shimadzu QP-5050A GC series apparatus fitted with a FID detector. The FID detector temperature was 300 °C. To obtain the same elution order as GC-MS, the simultaneous auto-injection was performed duplicate on the same column, applying the same operational conditions. The relative percentage amounts of separated compounds were calculated from the FID chromatograms. Moreover, flame ionization detector (FID) which was operated in ionization potential mode at 70 eV and used the same program reported above, was applied for quantification purpose and calculating the relative area percentage (area %) without the use of correction factors.

Antimicrobial assay

Disc diffusion method was employed for the determination of antibacterial activity of Zae M. The examined organisms were included three gram negative bacteria, *Pseudomonas aeroghinosa* (ATCC 9027), *Salmonella paratyphi* (ATCC 4420) and *Proteus mirabilis* (ATCC 29906), two gram positive species including *Bacillus subtilis* (ATCC 9372) and *Staphylococcus aureus* (ATCC 6538) and a fungus, *Candida albicans* (ATCC 10231) which were purchased in lyophilized culture from Pasture institute, Tehran, Iran. The activated microorganisms were transferred to Muller Hinton Broth medium (Merck, Germany) and were incubated overnight at 37°C. The turbidity was corrected until standard of 0.5 McFarland's [108 colony forming units (CFUs) per ml] by adding the sterile distilled water. About 10 ml of the prepared inoculums (106CFUs per ml) was spread over the autoclaved Muller Hinton Agar Medium. Then the sterile discs of Whatman paper with 6 millimeters diameter, which were impregnated with 10 μl of the essential oil and dissolved in 10 cc of 50% aqueous DMSO, were placed on the surface of the media. The plates were incubated for 30 min in the refrigerator to allow the essential oil to diffuse. Then they were incubated at 37°C for 24 h. Finally, the inhibition zones around the sterile discs were measured. In order to compare the potency of the antimicrobial activity of the volatile oil, standard discs of Amikacin (the concentration) as a positive control were placed on the seeded plates.¹⁰⁻¹¹ All of the experiments were performed in triplicate. Mean Inhibition Zone Diameter (MIZD) as well as the SD were calculated as well. The plates that indicated significant antibacterial activities were further assayed for their Minimum Inhibitory Concentration (MIC) at the tested concentration. Serial two-fold dilutions of sample were prepared in the broth. Cultures containing only sterile nutrient broth with no influence on bacterial growth were used as controls. An equal volume of the adjusted inoculums were added to the each test tube. After the incubation at 37°C for 24 h, the MIC was read. The MIC was

defined as the lowest concentration of a fraction which was able to completely inhibit the growth of the each bacterial strain.¹²

Results

The essential oil obtained by the hydrodistillation of aerial parts of *Z. absinthifolia* yielded a lemon yellow colored volatile oil of 0.15% (v/w). Table 1 illustrated phyto constituents of the ZaeM listed in the order of their elution from DB1 column. The listed components were qualitatively and quantitatively analyzed by GC-Mass and GC-FID.

Nineteen components consisting 98.22% of ZaeM were identified while 1.78% of the essential oil remained unidentified. Among identified phytochemicals, octyl acetate was the main compound (47.29%). Moreover, percentage of n-octanol (25.79%), octyl butyrate (10.15%) and octanoic acid octyl ester (7.9%) were more than other identified components. Apart from the major volatiles, only (E)-5-decenyl acetate (2.25%) and octanal (1.19%) exceeded a content of 1% of the total oil, while the remaining volatiles (n=13) were present in low amounts.

Table 1. Percentage of chemical components of Zae M.

| No. | Compound | %Area | RI ^a | Molecular formula | Identification method |
|-------------------------|----------------------------|-------|-----------------|--|-----------------------|
| 1 | n-Hexanal | 0.13 | 781 | C ₆ H ₁₂ O | RI+MS |
| 2 | α-Pinene | 0.53 | 928 | C ₁₀ H ₁₆ | RI+MS |
| 3 | Camphene | 0.1 | 940 | C ₁₀ H ₁₆ | RI+MS |
| 4 | β-Pinene | 0.06 | 967 | C ₁₀ H ₁₆ | RI+MS |
| 5 | Octanal | 1.19 | 978 | C ₈ H ₁₆ O | RI+MS |
| 6 | 1-Hexyl acetate | 0.14 | 1003 | C ₈ H ₁₆ O ₂ | RI+MS |
| 7 | 1,8-Cineol | 0.61 | 1018 | C ₁₀ H ₁₈ O | RI+MS |
| 8 | (Z)-3-Octen-1-ol | 0.25 | 1042 | C ₈ H ₁₆ O | RI+MS |
| 9 | n-Octanol | 25.79 | 1059 | C ₈ H ₁₈ O | RI+MS |
| 10 | Nonanal | 0.11 | 1081 | C ₉ H ₁₈ O | RI+MS |
| 11 | (E)-5-decenyl acetate | 2.25 | 1177 | C ₁₂ H ₂₂ O ₂ | RI+MS |
| 12 | Octyl acetate | 47.29 | 1199 | C ₁₀ H ₂₀ O ₂ | RI+MS |
| 13 | Bornyl acetate | 0.66 | 1268 | C ₁₂ H ₂₀ O ₂ | RI+MS |
| 14 | n-octyl propionate | 0.09 | 1283 | C ₁₁ H ₂₂ O ₂ | RI+MS |
| 15 | Octyl butyrate | 10.15 | 1372 | C ₁₂ H ₂₄ O ₂ | RI+MS |
| 16 | β-bourbonene | 0.37 | 1383 | C ₁₅ H ₂₄ | RI+MS |
| 17 | β-caryophyllene | 0.42 | 1417 | C ₁₅ H ₂₄ | RI+MS |
| 18 | Caryophyllene oxide | 0.18 | 1571 | C ₁₅ H ₂₄ O | RI+MS |
| 19 | Octanoic acid, octyl ester | 7.9 | 1761 | C ₁₆ H ₃₂ O ₂ | RI+MS |
| Total compounds | | 19 | | | |
| Esters | | 68.48 | | | |
| Alcohol | | 26.04 | | | |
| Aldehyde | | 1.43 | | | |
| Monoterpenes | | 1.30 | | | |
| sesquiterpenes | | 0.97 | | | |
| Total Identified | | 98.22 | | | |
| Not identified | | 1.78 | | | |
| Total identified | | 98.22 | | | |

^aCompounds listed in order of elution from a DB-1 column, ^bIdentification method: (RI= Retention indices, MS=Mass spectroscopy).




Figure 1. The identified chemical groups from the essential oils of the aerial parts of ZaeM.

Table 2. The percentage of the major compounds of different *Z. absinthifolia* volatile oils growing in different areas.^{3,5-7}

| No | Compounds | ZaeM ¹ | ZaeF ² | ZaeL ³ | ZaeKh ⁴ |
|----|---------------------|-------------------|-------------------|-------------------|--------------------|
| 1 | n-Hexanal | 0.13 | 0.2 | - | - |
| 2 | α-Pinene | 0.53 | 1.1 | 5.3 | - |
| 3 | Camphene | 0.1 | 0.2 | - | - |
| 4 | β-Pinene | 0.06 | 0.1 | - | - |
| 5 | Octanal | 1.19 | 0.2 | - | - |
| 6 | 1,8-Cineol | 0.61 | - | 0.2 | 0.3 |
| 7 | n-Octanol | 25.79 | 9.6 | - | 4.1 |
| 8 | Octyl acetate | 47.29 | 3.8 | 24.7 | 12.2 |
| 9 | Bornyl acetate | 0.66 | 1.5 | - | - |
| 10 | Octyl butyrate | 10.15 | 19.2 | - | - |
| 11 | β-Bourbonene | 0.37 | - | 3.3 | 0.4 |
| 12 | β-Caryophyllene | 0.42 | 13.9 | 22.2 | 14.6 |
| 13 | Caryophyllene oxide | 0.18 | 5.7 | - | 2.4 |

¹Z. absinthifolia essential oil gathered from Mishoodaghi mount in East-Azarbaijan Province of Iran. ²Z. absinthifolia essential oil gathered from northern mountains of Firozabad in Fars province of Iran. ³Z. absinthifolia essential oil gathered from Lonbar in Khalkhal, northwest of Iran. ⁴Z. absinthifolia essential oil gathered from north Khoramabad, Lorestan province of Iran.

As it could be seen in the Table 1 and Figure 1, phytochemicals of ZaeM were predominantly made up of firstly, esters (68.48%), secondly, alcohols (26.04%) and thirdly, Aldehydes (1.43%). Also the percentage of monoterpene and sesquiterpenes were 1.30% and 0.97 %, respectively. Table 2 exhibited the percentages of major constituents of ZaeM, *Z. absinthifolia* growing in northern mountains of Firozabad in Fars province (ZaeF), *Z. absinthifolia* growing in Lonbar in Khalkhal, in northwest of Iran (ZaeL) and north Khoramabad, Lorestan province of Iran (ZaeKh) successively.

The results of the antimicrobial test demonstrated that the only susceptible strain to Zae M was *B. subtilis*, a gram positive bacterium, with MIZD±SD of 22.95±1.90 mm and MIC of 25 mg/ml. The MIZD±SD (mm) of Amikasin was 42.3±1.50 mm.

Discussion

There are several reports about qualitative and quantitative changes of chemical composition of the essential oils obtained from different species of *Z. absinthifolia* due to the effects of the harvesting time and season, the geographical location, the altitude, the climate and the other factors including the chemo type, the reproductive stage, the cutting height, the drying condition, which could extremely affect the yield and composition of volatile oils of the same species.^{8,13} These findings could have important implications for collecting herbs from the best place and time to obtain the highest quality essential oils.

As it was illustrated in Table 2, the major constituents of Zae grown in different regions have shown significant differences. As it was evident in this table, the main constituents of the essential oils were not identical. The main similar chemical constituents were n-octanol, octyl acetate, octyl butyrate and β-caryophyllene, whereas ZaeM obtained the highest amount of n-octanol and octyl

acetate among remaining ones. Likewise, ZaeF and ZaeKh contained the highest percentages of octyl butyrate and β-caryophyllene, respectively. So these findings further support the effect of the mentioned factors on the yield and chemical composition of the essential oils.

The plants of the Apiaceae family possess a range of compounds with many biological activities. Some of the previously reported pharmacological activities were their ability to induce apoptosis,¹⁴ antibacterial,¹⁵ hepatoprotective¹⁶ and vaso-relaxant,¹⁷ cyclooxygenase inhibitory effects and antitumor properties.¹⁸ In the recent years, there has been an increasing interest in screening herbs for their antibacterial activity due to development of new types of effective antimicrobial phyto compounds such as extracts of herbs and their essential oils, for using as the food preservatives or other goals.¹⁹ Moreover, screening of the volatile oils as rich source of biologically active compounds have been carried out to indicate antibacterial activities of these natural products. So, it could be reasonable to expect a variety of phyto constituents in these oils with the antibacterial activity and antibiotic potential.²⁰ To the best of our knowledge, this was the first report on the antimicrobial activity of ZaeM. The results obtained from the *in vitro* antimicrobial test, showed that ZaeM indicated the inhibitory activity against *B. subtilis*, a gram positive strain. As seen in Table 1, octanol, as a long chain alcohol and a fundamental phyto constituent that includes almost a quarter of ZaeM (25.79%), has been reported to possess antimicrobial activity.²¹

Conclusion

As a conclusion, ZaeM could have antimicrobial potentials against *B. subtilis*. Further investigations are needed to isolate antimicrobial compounds from this volatile oil. Moreover, ZaeM worth to the further investigations for exploring further

biological activities as well as the isolation and identification of biologically active components.

Acknowledgment

This research is a part of the thesis of a Pharm D student at Pharmacy Faculty of Tabriz University of Medical Sciences. Hereby, assistance, cooperation and funding of Drug Applied Research Center as well as research deputy of Tabriz University of Medical Sciences and the participants would be appreciated.

Conflict of interests

The authors claim that there is no conflict of interest.

References

1. Welkie GW. Taxonomic distribution of dicotyledonous species capable of root excretion of riboflavin under iron deficiency. *J Plant Nutr.* 2000;23(11-12):1819-31. doi:10.1080/01904160009382145
2. Mozaffarian V. Identification of medicinal and aromatic plants of Iran. 2nd ed. Tehran: Farhang Mo'aser; 2015. p. 1232.
3. Jahangiri S, Shafaghat A. Chemical Constituents of Methanol and n-Hexane Extracts of *Zosimia absinthifolia* (Seeds). *Nat Acad Sci Lett.* 2014;37(4):325-30. doi:10.1007/s40009-014-0227-6
4. Shafaghat A, Salimi F, Shoaie Z, Aslaniyan N. Flavonoids from *Zosimia absinthifolia* (Vent.) Link. *Middle East J Sci Res.* 2011;7(6):864-8.
5. Shafaghat A. Comparison of Chemical Composition of Essential Oil and n-Hexane Extracts of *Zosimia absinthifolia* (Vent.) Link. *J Essent Oil Bear Plants.* 2011;14(4):490-3. doi:10.1080/0972060x.2011.10643606
6. Javidnia K, Miri R, Soltani M, Khosravi AR. Constituents of the Oil of *Zosimia absinthifolia* (Vent.) Link. from Iran. *J Essent Oil Res.* 2008;20(2):114-6. doi:10.1080/10412905.2008.9699968
7. Esmaili A, Amiri H, Rustaiyan A, Masoudi S, Anaraki MT. The Essential Oils of Two Umbelliferae, *Zosimia absinthifolia* (Vent.) Link. and *Smyrniopsis aucheri* Boiss. Growing Wild in Iran. *J Essent Oil Bear Plants.* 2010;13(1):73-7. doi:10.1080/0972060x.2010.10643793
8. Nikbakht MR, Sharifi S, Emami SA, Khodaie L. Chemical composition and antiproliferative activity of *Artemisia persica* Boiss. and *Artemisia turcomanica* Gand. essential oils. *Res Pharm Sci.* 2014;9(2):155-63.
9. Oreizi E, Rahiminejad MR, Asghari G. Influence of Environment on Glandular Trichomes and Composition of Essential Oil of *Perovskia abrotanoides* Karel. *Jundishapur J Nat Pharm Prod.* 2014;9(4):e16432. doi:10.17795/jjnpp-16432
10. Essawi T, Srour M. Screening of some Palestinian medicinal plants for antibacterial activity. *J Ethnopharmacol.* 2000;70(3):343-9. doi:10.1016/s0378-8741(99)00187-7
11. Sharififar F, Moshafi MH, Mansouri SH, Khodashenas M, Khoshnoodi M. In vitro evaluation of antibacterial and antioxidant activities of the essential oil and methanol extract of endemic *Zataria multiflora* Boiss. *Food Control.* 2007;18(7):800-5. doi:10.1016/j.foodcont.2006.04.002
12. Khodaie L, Delazar A, Lotfipour F, Nazemiyeh H. Antioxidant and Antimicrobial activity of *Pedicularis sibthorpii* Boiss. and *Pedicularis wilhelmsiana* Fisch ex. *Adv Pharm Bull.* 2012;2(1):89-92. doi:10.5681/apb.2012.012
13. Temime SB, Campeol E, Cioni PL, Daoud D, Zarrouk M. Volatile compounds from Chétoui olive oil and variations induced by growing area. *Food Chem.* 2006;99(2):315-25. doi:10.1016/j.foodchem.2005.07.046
14. Pae H, Oh H, Yun Y, Oh G, Jang S, Hwang KM, et al. Imperatorin, a furanocoumarin from *Angelica dahurica* (Umbelliferae), induces cytochrome c-dependent apoptosis in human promyelocytic leukaemia, HL-60 cells. *Pharmacol Toxicol.* 2002;91(1):40-8. doi:10.1034/j.1600-0773.2002.910107.x
15. Chou S, Everngam M, Sturtz G, Beck J. Antibacterial activity of components from *Lomatium californicum*. *Phytother Res.* 2006;20(2):153-6. doi:10.1002/ptr.1828
16. Ahmed B, Alam T, Varshney M, Khan S. Hepatoprotective activity of two plants belonging to the Apiaceae and the Euphorbiaceae family. *J Ethnopharmacol.* 2002;79(3):313-6. doi:10.1016/s0378-8741(01)00392-0
17. Senejoux F, Demougeot C, Cuciureanu M, Miron A, Cuciureanu R, Berthelot A, et al. Vasorelaxant effects and mechanisms of action of *Heracleum sphondylium* L. (Apiaceae) in rat thoracic aorta. *J Ethnopharmacol.* 2013;147(2):536-9. doi:10.1016/j.jep.2013.03.030
18. Oroojalian F, Kasra Kermanshahi R, Azizi M, Bassami MR. Phytochemical composition of the essential oils from three Apiaceae species and their antibacterial effects on food-borne pathogens. *Food Chem.* 2010;120(3):765-70. doi:10.1016/j.foodchem.2009.11.008
19. Shan B, Cai Y, Brooks J, Corke H. The in vitro antibacterial activity of dietary spice and medicinal herb extracts. *Int J Food Microbiol.* 2007;117(1):112-9. doi:10.1016/j.ijfoodmicro.2007.03.003

20. Prabuseenivasan S, Jayakumar M, Ignacimuthu S. In vitro antibacterial activity of some plant essential oils. *BMC Complement Altern Med.* 2006;6(1):1-8. doi:10.1186/1472-6882-6-39

21. Kubo I, Muroi H, Himejima M. Antibacterial activity against *Streptococcus mutans* of mate tea flavor components. *J Agric Food Chem.* 1993;41(1):107-11. doi:10.1021/jf00025a023