RELAXANT EFFECTS OF OCIMUM BASILICUM ON GUINEA PIG TRACHEAL CHAINS AND ITS POSSIBLE MECHANISM(S)

MOHAMMAD HOSSEIN BOSKABADY, SAHAR KIANI, BEHNIA HAGHIRI

Department of Physiology, Ghaem Medical Centre, Mashhad University of Medical Sciences, Mashhad, Iran

ABSTRACT

Therapeutic effects of Ocimum basilicum on respiratory diseases especially dyspnea have been reported in Iranian ancient medical books. In the present study, the relaxant effects of macerated and soxhlet extracts of this plant on tracheal chains of guinea pigs were evaluated. The relaxant effects of 4 cumulative concentrations of macerated and soxhlet extracts (0.25, 0.5, 0.75 and 1.0 W/V) in comparison with saline as negative control and 4 cumulative concentrations of theophylline (0.25, 0.5, 0.75, and 1.0 mM) as positive control were examined on precontracted tracheal chains of two groups of 6 guinea pig by 60 mM KCl (group 1) and 10 µM methacholine (group 2). Decrease in contractile tone of tracheal chains was considered as relaxant effect. In group 1 experiments only the last two higher concentrations of theophylline showed significant relaxant effect compared to that of saline (p<0.001 for both concentrations), which were significantly greater than those of macerated and soxhlet extracts (p<0.001 for all cases) and in group 2 experiments both macerated and soxhlet extracts showed concentrationdependent relaxant effects compared to that of saline (p < 0.05 to p < 0.001 for both extracts). There were significant differences between the relaxant effects of both extracts with those of theophylline in group 2 experiments (p<0.01 to p<0.001). The relaxant effects of macerated and soxhlet extracts in group 1 were significantly lower than those of groups 2. These results showed a potent relaxant effect of Ocimum basilicum on tracheal chains of guinea pigs which were lower than theophylline at concentrations used.

Keywords: Ocimum basilicum, Respiratory diseases Dyspnea, Relaxant

INTRODUCTION

Ocimum basilicum L. (labiatae) is a grassy and annual plant. The leaves of this plant are oval with a sharp tip and the flowers are yellow, white and pink. It is a native plant of Iran, Afghanistan and India (1,2,3). This plant is a popular culinary herb and its essential oils have been used for many years in food perfumery and dental produts (4) in Iranian traditional medicine (5,6).

According to the literature, about 45 compounds are found in volatile oil of this plant and the major compounds are linalol, eugenol, methylchavicol, methylcinnamat, linolen, ocimene, pinene, cineol, anethol, estragol, thymol, citral and comphor (1,7,8).

Different parts of Ocimum basilicum have diuretic (1,3), antipyretic (1) and antitussive effects (1,2) and have been used to treat gastritis, stomach-ache, flatulence and constipation (1-3,9-11), vomiting and hiccup (1). The seeds and the boiled extract are used for treatment of diarrhea (1). This plant also has therapeutic effects for nasal polyps (1), upper respiratory tract diseases and dyspnea (1,2), inflammation of the urinary tract (2) and it has been used as a bathing solution for treatment of ulcers (2,9) and increase in lactation (3). Inhalation of the fragrance of this

plant is useful to alleviate headache (1), and also it produces a favorable impression on the mental activity (12). In the Previous studies, the antiinflammatory (10,13,14), anti-HIV (15), antioxidant (5,16-18), antibacterial and antifungal (4,19-21) effects of this plant have been reported. In the present study, the relaxant effect of macerated and soxhlet extracts of Ocimum basilicum on guinea pig tracheal chains was examined.

MATERIALS AND METHODS

Plant and extracts

Ocimum basilicum was Collected from the School of Pharmacy and identified by Mr. Joharchi and the specimen number of the plant is 19484. The soxhlet extract was prepared as follows: Fifty grams of the chopped, dried plant leaves were extracted with 300 ml distilled water by suxhelat apparatus. For the preparation of the macerated extract, the same amount of plant was macerated with 300 ml distilled water (on a shaker) for 48 hr. The solvent of both extracts were then removed under reduced pressure at 50 °C and distilled water were added to residues in such a way that plant ingredient concentration in the final soxhlet extracts were 10% W/W.

Correspondence: M.H. Boskabady, Department. of Physiology, Ghaem Medical Centre, Mashhad, Post Code 91735, Iran, Email: m-boskabady@mums.ac.ir

Tissue preparations

Male guinea pigs (400-700 g) were killed by a blow on the neck and tracheas were removed. Each trachea was cut into 10 rings (each containing 2-3 cartilaginous rings). The cartilages of all rings were then cut open opposite to the trachealis muscle, and sutured together to form a tracheal chain (22). Tissue was then suspended in a 10 ml organ bath (organ bath 61300, Bio-Science Palmer-Washington, Sheerness, Kent U.K.) containing Krebs-Henseliet solution of the following composition (mM): NaCl 120, NaHCO₃ 25, MgSO₄ 0.5, KH₂PO₄ 1.2, KCl 4.72, CaCl₂ 2.5 and dextrose 11.

The Krebs solution was kept at 37° C under stream of 95% O₂ and 5% CO₂ gases. Tissue was suspended under an isotonic tension of 1 g and allowed to equilibrate for at least 1 h while it was washed with Krebs solution every 15 min.

Protocols

The relaxant effects of four cumulative concentrations of macerated and soxhlet extracts (0.25, 0.5, 0.75 and 1.0 g/100 ml), four cumulative concentrations of theophylline anhydrous (Sigma Chemical Ltd UK) (0.25, 0.5, 0.75, and 1.0 mM) as positive control, and saline as negative control were examined. For preparation of different concentrations in the case of macerated and soxhlet extracts, 0.25 ml of 10% W/V of the concentrated extracts and in the case of theophylline, 0.25 ml of 10 mM solution were added to the organ bath. The consecutive volumes were added to organ bath at 5 min intervals.

In each experiment the effect of four cumulative volumes of each extract, four cumulative volumes of theophylline, or saline on contracted tracheal smooth muscle were determined after exposure of tissue to the solution for 5 min. A decrease in tone was considered as a relaxant (broncho-dilatory) effect and expressed as positive percentage change in proportion to the maximum contraction and an increase in tone was considered as a contractile (bronchoconstrictory) effect which was expressed as negative percentage change (23).

The relaxant effect of different solutions were tested with two different experimental designs (n=6 for each group) as follows:

- 1. On tracheal chains contracted by 60 mM KCl (group 1 experiments).
- 2. On non-incubated tracheal chains contracted by 10 μ M methacholine hydrochloride (Sigma Chemical Ltd UK), (group 2 experiments).

The relaxant effects in two groups of experiments, were examined in two different series of tracheal chains. All experiments were performed randomly with a 1 h resting period of

tracheal chains between each consecutive experiments while washing the tissues every 15 min with Krebs solution. In all experiments responses were recorded on a kymograph (ET8 G-Boulitt, Paris) and were measured after fixation.

Statistical analysis

All data were expressed as mean \pm SEM. Data of relaxant effects of different concentrations of extracts were compared with the results of negative and positive control using ANOVA with Tukey Kramer post test. The data of relaxant effect obtained in two groups of experiments were also compared using unpaired t test. The relaxant effect of two extracts and theophylline were related to the concentrations using least square regression. Significances were accepted for p<0.05.

RESULTS

Relaxant (bronchodilatory) effect

In group 1 experiments, only the two higher concentrations of theophylline showed significant relaxant effects compared to those of saline (p<0.001 for both cases). In addition, the effects of the last two higher concentrations of both extracts were significantly lower than those of theophylline (p<0.001 for both cases). There was no significant difference between the effects of different concentrations of the two extracts (Table 1).

In groups 2 experiments, both extracts from Ocimum basilicum and theophylline showed relatively potent and concentration-dependent relaxant effects on tracheal chains of guinea pig. The relaxant effects of the most concentrated extracts and theophylline were significantly higher than those of saline (p < 0.05 to p < 0.001). Only the low concentration of theophylline did not show significant relaxant effect compared to that of saline (Table 2). However , the effects of two higher concentrations of macerated extract and only the highest concentration of soxhlet extract in group 2 were significantly lower than those of theophylline (p<0.01 to p<0.001), (Table 2). In contrast, the effects of the lowest concentrations of both extracts were significantly greater than that of theophylline (p<0.01 and p<0.001 for macerated and soxhlet extract respectively), (Table 2). In addition, the effects of the different concentrations of soxhlet extract in this group were greater than those of macerated extracts but only for the lower concentration, which were statistically significant (p<0.05), (Table 2).

Concentration	Saline	Macerate	d Soxhlet	Theophylline
		extract	extract	
		1.67±1.3	2.83±1.83	-5.20±2.16
0.25	-	NS	b NS bb nS	NS
		2.17±1.64	4.83±3.37	13.65±6.15
0.5	-	NS 1	ns NS ns nS	NS
		3.17±2.32	2 7.00±5.00	49.00±5.77
0.75	-	NS b	bb NS bbb nS	aaa
		4.17±2.99	9.50±6.70	85.83±6.40
1	0.90 ± 0.640	NS b	bb NS bbb nS	aaa

Table 1. Relaxant effect of two different extracts of *Ocimum basilicum* in comparison with negative control (saline) and positive control (theophylline) in group 1 experiments (contracted tracheal chains by 60 mM KCl).

Values are presented as mean±SEM. Statistical differences between the effect of extracts and theophylline with negative control (saline); NS: non-significant difference, aaa: p<0.001. Statistical differences between the effect of extracts and positive control (theophylline); ns: non-significant difference, b: p<0.05, bb: p<0.01, bbb: p<0.001. Statistical differences between the effect of two extracts; nS: non-significant difference. The unit of concentration for extracts was g/100 ml and for theophylline was mM.

Table 2. Relaxant effect of two different extracts of *Ocimum basilicum* in comparison with negative control (saline) and positive control (theophylline) in group 2 experiments (contracted tracheal chains by 10 µM methacholine).

Concentration	Saline	Macerated	Soxhlet	Theophylline
		extract	extract	
		7.00±2.31	13.83±2.32	-2.01 ± 0.34
0.25		a bb	aaa bbb c	NS
		11.33±2.89	19.5±4.05	12.32±2.22
0.5		aa ns	aaa ns nS	aa
		15.00±3.16	25.5±4.72	33.21±4.12
0.75		aa bb	aaa ns nS	aaa
		19.50±3.22	30.83±5.24	74.88±6.09
1	0.90±0.640	aa bbb	aaa bbb nS	aaa

Values are presented as mean±SEM. Statistical differences between the effect of extracts and theophylline with negative control (saline); NS: non-significant difference, a: p<0.05, aa: p<0.01, aaa: p<0.001. Statistical differences between the effect of extracts and positive control (theophylline); ns: non-significant difference, bb: p<0.01, bbb: p<0.001. Statistical differences between the effect of two extracts; nS: non-significant difference, c: p<0.05. The unit of concentration for extracts was g/100 ml and for theophylline was mM.

Table 3	Correlation	(r) of th	e relaxant	effects	of two	different	extracts	of	Ocimum	basilicum	and
theophyl	line with their	r concentr	ations in th	ree grou	ups of ex	periments	5.				

DifferentSubstances	Group 1	Group 2
Macerated extract	0.192 NS	0.576**
Soxhlet extract	0.234 NS	0.558**
Theophylline	0.935***	0.930***

Statistical significances; NS: non-significant difference, **: p<0.01, ***: p<0.001.



Figure 1. Concentration response curves of the relaxant effect of theophylline in two groups of experiments (group 1; KCl induced contraction (o) and group 2; methacholine induced contraction of tracheal chains (\Box , n=6 for each group). Statistical differences in the relaxant effect between group 1 and group 2; NS: non-significant difference.



Figure 3. Concentration response curves of the relaxant effect of soxhlet extract from *Ocimum basilicum* in two groups of experiments (group 1; KCl induced contraction (o) and group 2; methacholine induced contraction of tracheal chains (\Box , n=6 for each group). Statistical differences in the relaxant effect between group 1 and group 2; NS: non-significant difference, *; p<0.05, **: p<0.01.

Comparison of the relaxant effect between two groups of experiments

The relaxant effects of higher concentrations of both extracts in group 2 were statistically greater



Figure 2. Concentration response curves of the relaxant effect of macerated extract of *Ocimum basilicum* in two groups of experiments (group 1; KCl induced contraction (o) and group 2; methacholine induced contraction of tracheal chains (\Box , n=6 for each group). Statistical differences in the relaxant effect between group 1 and group 2; NS: non-significant difference, *; p<0.05, **: p<0.01.

than those of group 1 experiments (p<0.05 to p<0.01). However, there were no significant differences in the effects of all concentrations of theophylline between groups 1 and 2 (Fig. 1, 2, and 3)

Correlation between concentrations of solutions and their relaxant effects

There were significant positive correlations between the relaxant effects of both extracts and theophylline with concentrations of the solutions in group 2 experiments (p<0.01 to p<0.001), (Table 3). The correlation between the relaxant effect and concentration of theophylline in group 1 was also significant (p<0.001).

DISCUSSION

In this study the relaxant (bronchodilatory) effects of macerated and soxhlet extracts of Ocimum basilicum in comparison with saline as negative control and theophylline as positive control were studied. The relaxant effect of both extracts and theophylline were concentration dependent. There were positive correlations between increase in concentrations and the relaxant effects of both extracts in group 2, and theophylline in groups 1 and 2 experiments. The relaxant effects of all concentrations of macerated extract were nonsignificantly smaller than those of soxhlet extract in both groups of experiments. The absence of any obvious relaxant effect of macerated and soxhlet extract of Ocimum basilicum in group 1 and relatively potent effect of the extracts in group 2 experiments may be due to the opening of potassium channels which has been demonstrated previously (27). If the extracts from Ocimum basilicum open potassium channels, they should not have any effect on tracheal chains contracted by KCl, while they should have effects on the tracheal chain which were contracted by metacholine. In fact, the results of groups 2 may support this hypothesis. Since KCl affect calcium channels (28) and calcium channel blockers are bronchodilator (29,30), another explanation for results of the experiment is the absence of blocking effects of this extract on calcium channels.

The effects of different concentrations of soxhlet extract in both groups of experiments were greater than those of macerated extract. Which may suggest that the effective substances in soxhlet extract are higher than that of macerated extract. The other possible mechanisms for broncho-dilatory effect of Ocimum basilicum include: stimulation of β -adrenergic receptors (23,24), blocking of histamine H₁ receptors (25), anticholinergic activity (26), stimulation of inhibitory nerve other than adrenergic or cholinergic system (NANC) or inhibition of stimulatory NANC (31), methyl xanthin like activity (32), due to inhibition of phosphodiestrase (33). However, contributions of these mechanisms to bronchodilatory effect of extracts of Ocimum basilicum on tracheal chains of guinea pigs (s) should be clarified in further studies.

In addition Ocimum basilicum might also have an anti-inflammatory effect, which will contribute to the therapeutic effect of this plant on asthma. While anti-inflammatory (10,13,14) and anti-oxidant (5,16,17,18) effects of this plant have been reported, the effect of, Ocimum basilicum on airway inflammation which is present in asthma should be investigated in further studies.

REFERENCES

- 1. Mirheidar H. Herbal knowledge. 1st Edition. Islamic Culture Publication; 1990. pp. 85-91.
- 2. Volak J, Jiri S. Plantes Medicinales. Interpreter; Zamani S. 3rd Edition. Grund Edition; 1997. p. 240.
- 3. Zargary A. Medicinal plants. 5th Edition.Tehran: Tehran University Publications; 1990. pp. 312-314.
- 4. Suppakul P, Miltz J, Sonneveld K, Bigger SW. Antimicrobial properties of basil and its possible application in food packaging. J Agiric Food Chem 2003; 51:3197-207.
- Dasgupta T, Rao AR, Yadava PK. Chemomodulatory efficacy of basil leaf (Ocimum basilicum) on drug metabolizingand antioxidant enzymes, and on carcinogen-induced skin and forestomach papillomagenesis. Phytomedicine 2004; 11:139-51.
- Javanmardi J, Khalighi A, Kashi A, Bais HP, Vivanco JM. Chemical characterization of basil (Ocimum basilicum L.) found in local accessions and used in traditional medicines in Iran. J Agric Food Chem 2002; 50:5878-83.
- 7. Wang T, Cui SY, Hu XL, Lu RH. Study on the constituents of volatile oil from Ocimum basilicum. Zhongguo Zhong Yao Za Zhi 2003; 28:740-2.
- 8. Grayer RJ, Kite GC, Goldstone FJ, Bryan SE, Paton A, Putievsky E. Infraspecific taxonomy and essential oil chemotypes in sweet basil, Ocimum basilicum. Phytochemistry 1996; 43:1033-9.
- 9. Avesina. Law in Medicine. Interpreter; Sharafkhandy A. Theran: Ministry of Guidance publications; 1990. p. 302.
- 10. Singh S. Evaluation of gastric anti-ulcer activity of fixed oil of Ocimum basilicum Linn.and its possible mechanism of action. Indian. J Exp Biol 1999; 37:253-7.
- 11. Akhtar MS, Munir MJ. Evaluation of the gastric antiulcerogenic effects of Solanum nigrum, Brassica oleracea and Ocimum basilicum in rats. J Ethnopharmacol 1989; 27:163-76.
- 12. Satoh T, Sugawara Y. Effects on humans elicited by inhaling the fragrance of essential oils: sensory test, multi-channel thermometric study and forehead surface potential wave measurement on basil and peppermint. Anal Sci 2003; 19:139-46.
- 13. Singh S. Mechanism of action of antiinflammatory effect of fixed oil of Ocimum basilicum Linn. Indian. J Exp Biol 1999; 37:248-52.
- 14. Singh S. Comparative evaluation of antiinflammatory potential of fixed oil of different species of Ocimum and its possible mechanism of action. Indian J Exp Biol 1998; 36:1028-31.
- Yamasaki K, Nakano M, Kawahata T, Mori H, Otake T, Ueba N, Oishi I, Inami R, Yamane M, Nakamura M, Murata H, Nakanishi T. Anti-HIV-1 activity of herbs in Labiatae. Biol Pharm Bull 1998; 21:829-33.
- 16. Jayasinghe C, Gotoh N, Aoki T, Wada S. Phenolics composition and anti oxidant activity of sweet basil (Ocimum basilicum L.). J Agric Food Chem 2003; 51:4442-9.

- 17. Yun YS, Nakajima Y, Iseda E, Kunugi A. Determination of anti oxidant activity of herbs by ESR. Shokuhin Eiseigaku Zasshi 2003; 44:59-62.
- 18. Lee KG, Shibamoti T. Determination of antioxidant potential of volatile extracts isolated from various herbs and spices. J Agric Food Chem 2002; 50:4947-52.
- 19. Edris AE, Farrag ES. Antifungal activity of peppermint and sweet basil essential oils and their major aroma constituents on some plant pathogenic fungi from the vapor phase. Nahrung 2003; 47:117-21.
- Opalchenova G, Obreshkova D. Comaparative studies on the activity of basil --an essential oil from Ocimum basilicum L.-against multi-drug clinical isolates of the genera staphylococcus, Enterococcus and Pseudomonas by using different test methods. J Mocrobiol Methods 2003; 54:105-10.
- 21. Lachowicz KJ, Jones GP, Briggs DR, Bienvenu FE, Wan J, Wilcock A, Coventry MJ. The synergistic preservative effects of the essential oils of sweet basil(Ocimum basilicum L.) against acid-tolerant food microflora. Lett Appl Microbiol 1998; 26:209-14.
- 22. Holroyde MC. The influence of epithelium on the responsiveness of guinea-pig isolated trachea. Br J Pharmacol 1986; 87:501-507.
- 23. Martin CAE, Naline E, Bakdach H, Advenier C. Beta3 adrenoceptor agonists, BRL 37344 and SR 58611A do not induce relaxation of human, sheep and guinea-pig airway smooth muscle in vitro. Eur Respir J 1994; 7:1610-1615.
- 24. Linden A, Bergendal A, Ullman A, Skoogh BE, Lofdahl CG. Salmetrol, formetrol, and salbutamol in the isolated guinea-pig trachea: differences in maximum relaxant effect and potency but not in functional antagonism. Thorax 1993; 48:547-553.
- 25. Popa VT, Somani P, Simon P, Simon V. The effect of inhaled verapamil on resting bronchial tone and airway constriction by histamine and acetylcholine in normal and asthmatic subjects. Am Rev Respir Dis 1984; 130:106-113.
- 26. Lronards B, Rampart M, Herman AG. Selective M₃ muscarinic receptor inhibit smooth muscle contraction in rabit trachea without increasing the release of acetylcholine. J Pharmacol Exp Ther 1992; 263:773-770.
- 27. Buckle DR, Arch JRS, Boering NE, Foster KA, Taylor JF, Taylor SG, Shaw DJ. Relaxation effect of potassium channel activators BRL 38227 and Pinacidil on guinea-pig and human airway smooth muscle, and blockade of their effects by Glibenclamide and BRL 31660. Pulmon Pharmacol 1993; 6:77-86.
- 28. Perez-Guerrero C, Suarez, J, Herrera MD, Marhuenda E. Spasmolytic effect of tetrazepam on rat duodenum and guinea pig ileum. Pharmacol Res 1997; 35:493-494.
- 29. Miyahara Y, Kizawa Y, Sano M, Murakami H. Effect of organic and inorganic Ca²⁺ antagonists on acetylcholine induced contraction in molluscan (Mytilus edulis) smooth muscle. Gen Pharmacol 1993; 24:1419-1423.
- 30. McCaig D, DeJonckeere S. Effect of two Ca²⁺ modulator in normal and albumin sensitised guinea pig trachea. Eur J Pharmacol 1993; 249:53-63.
- Linden A, Lofdahl CG, Ullman A, Skoogh BE. Non adrenergic non cholinergic responses stabilize smooth muscle tone with and without parasympathetic activation in guinea pig isolated airways. Eu Respir J 1993; 6:425-433.
- 32. Meini S, Ballati L, Evangelista S, Manzini S, Isbufulline, a xanthine derivative, inhibits bronchoconstrictor responses produced by stimulation of capsaicin sensitive nerves in guinea-pig, in vitro and in vivo evidence. Pulmon Pharmacol 1993; 6:279-286.
- 33. Van Amsterdam RGM, Meurs H, Brouwer F, Postema JB, Timmermans A, Zaagsma J. Role of phosphoinosited metabolism in functional antagonism of airway smooth muscle contraction by betaadrenoceptor agonist. Eur J Pharmacol 1989; 172:175-183.