SOLUBILITY PREDICTION OF SULFONAMIDES AT VARIOUS TEMPERATURES USING A SINGLE DETERMINATION

¹JALAL HANAEE, ²ABOLGHASEM JOUYBAN, ¹SIAVOUSH DASTMALCHI, ¹KHOSRO ADIBKIA, ¹AHMAD MIRZAZADEH, ¹MOHAMMAD BARZEGAR-JALALI

¹School of Pharmacy, Tabriz University of Medical Sciences, ²Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

ABSTRACT

Solubility of sulphamethoxazole, sulphisoxazole and sulphasalazine in six solvents namely water, methanol, ethanol, 1-propanol, acetone and chloroform were determined at 15, 25, 37 and 45 °C. Two models derived from the Hildebrand solubility approach are proposed for solubility prediction at different temperatures using a single determination. The experimental data of the present work as well as data gathered from the literature have been employed to investigate the accuracy and prediction capability of the proposed models. The overall percent deviations between the predicted and experimental values were 10.78 and 14.63% which were comparable to those of the classical two and three parameter models. The proposed models were much superior to the two pure predictive models i.e., the ones which do not require experimental solubility determination, as the overall percent deviations produced by the latter models were 150.09 and 161.00%.

Keywords: Solubility; Sulphonamides; Modeling; Temperature; Prediction

INTRODUCTION AND THEORY

Solubility is one of the most important physicochemical properties of drugs. As a general rule, water-insoluble compounds may not act as a specific drug. The solubility of a solute could be affected by factors like pH, pKa, polarity etc. The modeling of solubility data provides not only a means of screening experimental data sets for outliers which require of possible redetermination, but also facilitates interpolation at other points falling between measured data. Solubility prediction employing models without a curve-fit parameter or the minimum number of curve-fit parameters is the final goal of the solubility data modeling. By using such models, researchers are able to estimate the unmeasured solubilities.

Some models were proposed for mathematical representation of solubility in mixed solvents at a constant (1-10) and different temperatures (11-16). Knowledge of solubility at various temperatures is useful in the physical stability

studies of liquid dosage forms, in processes involving temperature changes and in preformulation stage of a new drug where only small amount of the drug is available.

In the present report, the solubility of three sulphonamides, i.e. sulphamethoxazole, sulphisoxazole and sulphasalazine, in water, methanol, ethanol, 1-propanol, acetone and chloroform at 15-45 °C were measured. Sulfonamides have already been used by others (13) as model drugs for solubility investigations. Two models based on the Hildebrand solubility approach which can predict the solute solubility in the range of ±25 °C with acceptable error using a single experimental data determined at 25 °C are proposed and the superiority of these models to previously published models (13) in which no experimental data has been used to train the model, are shown. The errors of the proposed models in comparison with other models employing more than one data point, were satisfactory.

Five models for solubility prediction at different

Correspondence: M. Barzegar-Jalali, Department of Pharmaceutic, School of Pharmacy, Tabriz University of Medical Sciences, Tabriz 51664, Iran, Email: barzegar_jalali@yahoo.com

temperatures, i.e. the van't Hoff (17), the Hildebrand (17), the Grant-Prankerd (11, 12) and the extended Hildebrand-Scatchard models (13) which are expressed by the respective Eqs. I - V, have been reported.

$$\ln X_2 = \frac{-\Delta H_s}{RT} + A \tag{1}$$

$$\ln X_2 = \frac{\Delta S_s \ln T}{R} + B \tag{II}$$

$$\ln X_2 = C\ln T + \frac{D}{T} + E \tag{III}$$

$$-\ln X_{2} = \frac{-\Delta H_{f}(T_{m} - T)}{RTT_{m}} + \frac{V_{2}\phi_{1}^{2}(\delta_{1} - \delta_{2})^{2}}{RT\ln[\varepsilon\delta_{1}(\delta_{1} - \delta_{2})^{2}]} \quad (IV)$$

$$-\ln X_{2} = \frac{-\Delta H_{f}(T_{m} - T)}{RTT_{m}} + \frac{V_{2}\phi_{1}^{2}(\delta_{1} - \delta_{2})^{2}}{RT\ln[ET_{(30)}\delta_{1}(\delta_{1} - \delta_{2})^{2}]}(V)$$

In the above equations X_2 is the mole fraction solubility of the solute, ΔH_s and ΔS_s are the enthalpy and the entropy of solution, ΔH_f is enthalpy of solute fusion, R is the molar gas constant (1.9872 cal mol⁻¹ K⁻¹), T and T_m are the solution temperature and melting point of the solute (in K), A, B, C, D and E stand for the models constants, V₂ and δ_2 are the molar volume and solubility parameter of the solute, \ddot{o}_1 and δ_1 denote the volume fraction and solubility parameter of the solvent, (represents dielectric constant and ET(30) is the solvatochromic polarity parameter of the solvent. The value of \ddot{o}_1 which is very close to 1 was assumed to be equal to one (18-21).

Eqs. I and II are two parameter models and a minimum number of two experimental solubilities which are determined at two different temperatures are required in order to predict solubilities at other temperatures. The constant terms of Eq. III are calculated by using a minimum number of three solubilities at three different temperatures. Eqs. IV and V are pure predictive models and require no experimental solubility data but require parameters related to solute structure and solvent which are easily calculated (22,23) or found in literature (24-27). The proposed models are as follows:

$$-\ln X_{2} = -\ln X_{2}^{i} + \ln \left[\frac{V_{2} \phi_{1}^{2} (\delta_{1} - \delta_{2})^{2}}{RT} \right] + F \quad (VI)$$

$$-\ln X_{2} = -\ln X_{2}^{i} + \left[\frac{V_{2}\phi_{1}^{2}(\delta_{1} - \delta_{2})^{2}}{RT}\right]^{G} \quad (VII)$$

Where X_2^{i} is ideal mole fraction solubility of the solute, F and G are the models constants which are calculated via a single solubility determination at 25 °C by the following equations:

$$F = -\ln X_{2} + \ln X_{2}^{i} - \ln \left[\frac{V_{2} \phi_{1}^{2} (\delta_{1} - \delta_{2})^{2}}{RT} \right] \quad (VIII)$$

$$G = \frac{\ln \left(\ln \frac{X_{2}^{i}}{X_{2}} \right)}{\ln \left[\frac{V_{2} \phi_{1}^{2} (\delta_{1} - \delta_{2})^{2}}{RT} \right]} \quad (IX)$$

Eq. VII is applicable only to those cases where X_2^{i} > X_2 at 25 °C.

The value of $-\ln X_2^i$ in Eqs. VI and VII is given by Eq. X (28):

$$-\ln X_{2}^{i} = \frac{-\Delta H_{f}(T_{m} - T)}{RTT_{m}} \qquad (X)$$

 ΔH_f is calculated by Eq. XI:

$$\Delta H_{f} = \frac{0.02303 (T_{m} - T)RT_{m}}{\ln(\frac{T_{m}}{T})}$$
(XI)

Substitution of ΔH_f from Eq. XI into Eq. X yields Eq. XII.

$$- \ln X_{2}^{i} = \frac{0.02303 (T_{m} - T)^{2}}{T \ln \left(\frac{T_{m}}{T}\right)}$$
(XII)

The value of T in the latter equation is equal to 298.15 K (23). The value of X_2^i in the proposed models is calculated by means of Eq. XII.

To assess the prediction capability of the models, percent deviation (PD) for each data set, mean percent deviation(MPD) and number of sets with minimum PD values which are expressed as percentage of all sets(denoted as percent best adherence, PBA) are calculated. The latter two criteria are employed for overall comparison of the models:

$$PD = \frac{100}{N} \sum \left(\frac{\left| X_2^{calculated} - X_2^{Observed} \right|}{X_2^{Observed}} \right) \qquad (XIII)$$

$$MPD = \frac{\sum PD}{N_s}$$
(XIV)

$$PBA = \frac{N_{ba}}{N_{s}}.100 \qquad (XV)$$

where N denotes the number of experimental data in each set , N_{ba} is number of sets with the best adherence for a model and N_{s} is the total number of sets.

In the present work we compared the accuracy and predictability of the proposed models (Eqs. VI and VII) with the classical two and three parameter models (Eqs. I-III) using the solubility data of experimental three sulphonamides in six solvents at various temperatures. Also the proposed models were compared with the pure predictive models (Eqs. IV and V). In addition to our data, the solubility data for different solutes with diverse structures in variety of solvents at various temperatures which are collected from the literature were employed to demonstrate the suitably and applicability of the proposed models.

MATERIALS AND METHODS

Chemicals:

All solvents were reagent grade from Merck (Germany). Double-distilled water was used throughout this investigation.

The solutes (sulphamethoxazole, sulphisoxazole and sulphasalazine) were kindly supplied by Daru-Pakhsh, Tehran, Iran. The purity of drugs was checked by infrared spectrophotometer (Shimadzu, Japan) and the IR spectra of the samples were the same as those of the references (29) and were above 99.3 %. Melting point of the compounds were 169, 194 and 253 °C for sulphamethoxazole, sulphisoxazole and sulphasalazine, respectively (29).

Solubility determinations:

The solubilities of the sulphonamides were determined by equilibrating an excess amount of

solid with particular solvent at 15, 25, 37 and 45 °C using a thermostat incubator (± 1.7 °C). After reaching equilibrium (5 days), the solutions were filtered using filters of pore size 0.5 μ m and diluted with solvents. The sulphonamide concentrations were quantified by a double-beam UV spectrophotometer (Shimadzu, Japan) at different wavelengths given in table 1. In order to convert molar concentrations into the mole fractions, the densities of the saturated solutions were determined in 5 ml pycnometer.

RESULTS AND DISCUSSION

Table 1 shows the natural logarithm of mole fraction solubility of three sulphonamides in individual solvents at different temperatures together with λ_{max} values used for determination of the solutes concentration. For the all solvent systems, the higher temperature means higher solubility. The solutes exhibited maximum solubility in acetone and minimum in water.

Table 2 presents names of solutes and solvents used in our experiment as well as those which were taken from the literature, their physicochemical characteristics and the range of temperatures used in °C and number of data in each set (N). The mentioned parameters and quantities with the exception of N were used for calculation of the solutes solubility by means of Eqs. IV and V. Also some of these parameters are necessary for the solubility calculations via the proposed models.

In table 3 the values of PD, MPD and PBA for Eqs. I - VII calculated by Eqs. XIII and XIV together with the parameters of the proposed models i.e. F and G are listed. This table contains not only the results of our experimental data but also includes the results of data collected from the literature (12, 13, 26, 27, 30 -32). In the application of Eqs. I and II the solubility data at 25° and 37 °C were used in order to predict solubilities at other temperatures. In the data sets where the solubility value at 37 °C was not available, the value at the nearest temperature (35° or 40 °C) was used. The constant terms of Eq. III were calculated by solubility data at 25°, 37°, 50 °C or 0 °C in the absence of 50 °C datum. As mentioned above in the application of Eqs. IV and V no solubility data were required. The errors associated with the predicted solubilities by the latter models were generally very high and above the acceptable 30% level (33, 34). The corresponding MPD values were150.09 and 161.00%.

Eq. VI produced only one error higher than 30%. In contrast Eq. VII produced six PD values above 30% level. The number of PDs above 30% for Eqs. I – V were 2, 2, 4, 38 and 41, respectively. However, the MPD values for Eqs. VI and VII were 10.78 and 14.63, respectively which were comparable with those of two and three parameter models I - III (the corresponding MPD values were 9.64, 9.91 and 13.83%). Considering PBA criterion Eq. VII was the best as its PBA was 32.7% while suffering from the disadvantage that it could not be applied to 8 out of 51 sets because of its mathematical character. The corresponding criteria for Eqs. VI and I were 19.61%. This can be an advantage for Eq.VI which requires only a single solubility datum for predictions. In addition, Eq. VI dose not have the limitation of Eq.VII in that it is applicable for all data sets. Eqs. IV and V gave the lowest PBA values (1.96% and 0%) while the corresponding values of Eqs. II and III were 15.69 and 16.67% which were slightly less than those of Eq. I and VI.

CONCLUSION

Solubility prediction based upon the minimum number of experiments is valuable from practical point of view, especially in the preformulation studies of a new drug that only small amount of the drug is available and in the processes involving temperature changes. The solubility predictions at different temperatures using the solubility at 25 °C are proposed employing Eqs. VI and VII, which have a single constant term and their accuracy and prediction capability were much superior to those of pure predictive models (Eqs. IV and V) but overall comparable to those of two and three parameter models (Eqs. I – III) requiring two and three experimental solubilities.

	T(°C)	Acetone	Chloroform	Ethanol	Methanol	1-Propanol	Water	
	1.5	2.52	1 .25	5.25	1.(2	(01	11.07	
0.1.1	15	-2.53	-7.35	-5.35	-4.62	-6.21	-11.37	
Sulphamethoxazole	25	-2.26	-7.15	-5.01	-4.24	-5.86	-10.83	
	37	-1.96	-6.79	-4.66	-3.78	-5.42	-10.22	
	45	-a	-6.39	-4.43	-3.58	-5.21	-9.91	
$\lambda_{max} (nm)$		209.2	266.6	270.4	269.2	269.8	267.8	
	15	-9.39	-a	-9.83	-10.67	-10.02	-13.33	
Sulphasalazine	25	-9.22	-a	-9.54	-9.82	-9.85	-12.23	
	37	-8.83	- a	-9.32	-9.51	-9.47	-11.77	
	45	-a	-a	-9.06	-9.26	-9.20	-11.64	
λ _{max} (nm)		349.4	-	349.2	349.2	349.2	350.0	
	15	-3.59	-8.42	-5.92	-5.39	-6.53	-11.95	
Sulphisoxazole	25	-3.22	-8.12	-5.51	-4.97	-6.20	-11.45	
	37	-2.96	-7.75	-5.09	-4.55	-5.74	-10.93	
	45	_ ^a	-7.45	-4.83	-4.16	-5.47	-10.54	
λ_{max} (nm)		209.4	266.6	270.8	269.6	271.2	263.2	

Table 1 Natural logarithm of experimental mole fraction solubility data of sulphonamides in individual solvents at different temperatures and the corresponding λ_{max} values

^aNot determined.

Solute	Solvent	t(°C)	Ν	δ_1	ε ^b	ET ₍₃₀₎	δ_2	V_2	$T_m(K)$	$\Delta H_{\rm f}$	Reference
Sulphamethoxazole	Acetone	15-45	3	9.07 ^a	20.56	42.2	13 19 ^c	152.10°	169.0	7393 ^d	This work
Sulphamethoxazole	Chloroform	15-45	4	8 77 ^a	4.81	39.1	13.19°	152.10°	169.0	7393 ^d	This work
Sulphamethoxazole	Ethanol	15-45	4	12.58^{a}	24.55	51.9	13.19°	152.10°	169.0	7393 ^d	This work
Sulphamethoxazole	Methanol	15-45	4	13.77^{a}	32.66	55.4	13 19 [°]	152.10°	169.0	7393 ^d	This work
Sulphamethoxazole	1-Propanol	15-45	4	11.84 ^a	20.45	50.7	13.19°	152.10 [°]	169.0	7393 ^d	This work
Sulphamethoxazole	Water	15-45	4	24.52 ^a	78.54	63.1	13.19 ^c	152.10 [°]	169.0	7393 ^d	This work
Sulphasalazine	Acetone	15-45	3	9.07 ^a	20.56	42.2	15.40 ^c	207.30 °	253.0	9666 ^g	This work
Sulphasalazine	Ethanol	15-45	4	12.58 ^a	24.55	51.9	15.40 ^c	207.30 °	253.0	9666 ^g	This work
Sulphasalazine	Methanol	15-45	4	13.77 ^a	32.66	55.4	15.40 °	207.30 °	253.0	9666 ^g	This work
Sulphasalazine	1-Propanol	15-45	4	12.58 ^a	20.45	50.7	15.40 °	207.30 °	253.0	9666 ^g	This work
Sulphasalazine	Water	15-45	4	24.52 ^{-a}	78.54	63.1	15.40 ^c	207.30 ^c	253.0	9666 ^g	This work
Sulphisoxazole	Acetone	15-45	3	9.07 ^a	20.56	42.2	12.87 °	166.60 ^c	194.0	8035 ^d	This work
Sulphisoxazole	Chloroform	15-45	4	8.77 ^a	4.81	39.1	12.87 °	166.60 ^c	194.0	8035 ^d	This work
Sulphisoxazole	Ethanol	15-45	4	12.58 ^a	24.55	51.9	12.87 ^c	166.60 ^c	194.0	8035 ^d	This work
Sulphisoxazole	Methanol	15-45	4	13.77 ^a	32.66	55.4	12.87 °	166.60 °	194.0	8035 ^d	This work
Sulphisoxazole	1-Propanol	15-45	4	11.84 ^a	20.45	50.7	12.87 °	166.60 °	194.0	8035 ^d	This work
Sulphisoxazole	Water	15-45	4	24.52 ^a	78.54	63.1	12.87 °	166.60 °	194.0	8035 ^d	This work
Sulfanilamide	Water	20-75	7	24.52	78.54	63.1	12.98	110.40	438.54	7304	13
Acetylsulfanilamide	Water	20-75	7	24.52	78.54	63.1	13.58	134.20	454.28	7702	13
Sulfapyridine	Water	20-75	7	24.52	78.54	63.1	13.05	158.60	464.53	7978	13
Sulfadiazine	Water	20-75	7	24.52	78.54	63.1	13.85	151.10	528.28	9723	13
Sulfatiazole	Water	20-75	7	24.52	78.54	63.1	13.84	145.80	474.31	8231	13
Sulfamerazine	Water	20-75	7	24.52	78.54	63.1	13.48	164.60	509.11	9185	13
Sulfafurazole	Water	20-75	7	24.52	78.54	63.1	12.87	166.60	469.69	8111	13
Sulfamethyltiazole	Water	20-75	7	24.52	78.54	63.1	13.46	160.30	523.24	9579	13

Table 2 The details of solutes and solvents characteristics used for solubility calculations at different temperatures

Sulfadimidine

Sulfamethoxypyridazine

Water

Water

20-75

20-75

24.52

24.52

78.54

78.54

63.1

63.1

13.16

13.50

179.10

168.40

471.12

454.52

No.

Table 2	2 continued											
28	Sulfaproxyline	Water	20-75	7	24.52	78.54	63.1	12.55	222.90	460.31	7859	13
29	Sulphamethoxypyridazine	Ethyl acetate	20-40	5	8.85	6.02	38.1	13.50 ^d	168.40 ^d	454.52 ^d	7720 ^d	26
30	Sulphamethoxypyridazine	Ethanol	20-40	5	12.58 ^a	24.55	51.9	13.50 ^d	168.40 ^d	454.52 ^d	7720 ^d	26
31	Sulphamethoxypyridazine	Water	20-40	5	24.52	78.54	63.1	13.50 ^d	168.40 ^d	454.52 ^d	7720 ^d	26
32	Paracetamol	Ethyl acetate	20-40	5	8.85 ^a	6.02	38.1	13.40 ^e	118.66 ^e	442.28	6274	27
33	Paracetamol	Ethanol	20-40	5	12.58 ^a	24.55	51.9	13.40 ^e	118.66 ^e	442.28	6274	27
34	Paracetamol	Water	20-40	5	24.52	78.54	63.1	13.40 ^e	118.66 ^e	442.28	6274	27
35	Phenacetin	Water	20-40	5	24.52	78.54	63.1	11.55	148.40	407.40	6523 ^g	30
36	Phenacetin	Dioxane	20-40	5	10.01 ^a	2.21	36.0	11.55	148.40	407.40	6523 ^g	30
37	Naphthalene	Hexane	0-50	6	7.30 ^a	1.88	31.0	9.89 ^a	111.50 ^a	353.15	5250 ^g	32
38	Naphthalene	Carbon	0-50	6	8.71 ^a	2.23	32.4	9.89 ^a	111.50 ^a	353.15	5250 ^g	32
39	Naphthalene	Toluene	0-50	6	8.88 ^a	2.38	33.9	9.89 ^a	111.50 ^a	353.15	5250 ^g	32
40	Naphthalene	Benzene	0-50	6	9.06 ^{-a}	2.27	34.3	9.89 ^a	111.50 ^a	353.15	5250 ^g	32
41	Naphthalene	Chloroform	0-50	6	8.77 ^a	4.81	39.1	9.89 ^a	111.50 ^a	353.15	5250 ^g	32
42	Naphthalene	Chlorobenzene	0-50	6	9.59 ^a	5.62	36.8	9.89 ^a	111.50 ^a	353.15	5250 ^g	32
43	Naphthalene	Carbon disulfide	0-50	6	10.00 ^a	2.64	32.8	9.89 ^a	111.50 ^a	353.15	5250 ^g	32
44	Methyl paraben	Water	0-58	13	24.52	78.54	63.1	13.31 ^f	113.90^{f}	404.15	6143	12
45	Ethylparaben	Water	0-58	14	24.52	78.54	63.1	12.82 ^f	130.00^{f}	389.15	6334	12
46	Propyl paraben	Water	0-50	12	24.52	78.54	63.1	12.42^{f}	146.10^{f}	369.65	6692	12
47	Butylparaben	Water	0-35	8	24.52	78.54	63.1	12.09 ^f	$162.20^{\rm f}$	341.65	5760	12
48	5-5-Diethyl barbituric acid	Water	10-50	9	24.52	78.54	63.1	14.20 ^f	125.80 ^f	463.15	5710	12
49	5-5-Diisopropyl barbituric	Water	10-50	9	24.52	78.54	63.1	13.02 ^f	158.60 ^f	496.15	7978	12
50	acid 5-Ethyl 5-phenyl barbituric acid	Water	25-50	6	24.52	78.54	63.1	14.42 ^f	147.60 ^f	450.15	6058	12
51	Paracetamol	Water	5-70	17	24.52	78.54	63.1	13.40 ^e	118.66 ^e	443.15	7720	12

^a The values were taken from reference (25), ^b The values of ε and ET(30)were taken from reference (24), ^c The values of δ_2 and V₂ were taken from reference (26), ^d Taken from reference (13), ^e Taken from reference (31)

	The models	constants		PD°							
No. ^a	F٥	G⁵	Eq. IV	Eq. V	Eq. VI	Eq. VII	Eq. I ^c	Eq. II ^c	Eq. III ^c		
1	-3.2721	_c	95.14	94.62	9.52	_e	0.09	1.07	_f		
2	1.4705	0.6982	298.73	499.48	7.21	10.87	13.81	12.98	16.39		
3	3.2992	0.0215	163.48	165.19	8.83	5.16	1.88	2.11	4.48		
4	2.6226	0.7155	17.75	18.21	2.88	2.29	6.17	5.96	15.60		
5	2.5555	-0.7708	336.54	346.8	4.20	4.13	5.536	5.34	13.63		
6	3.2705	0.5470	104.125	86.26	11.51	2.65	4.04	4.99	11.19		
7	-0.4839	0.2910	76.64	69.31	14.77	13.89	16.45	15.40	_f		
8	1.4505	0.8851	563.29	614.17	27.44	32.81	10.62	10.81	32.97		
9	2.8221	-13.8535	1442.43	1476.33	28.63	125.50	41.32	41.91	93.54		
10	1.7612	1.0007	414.23	486.59	18.66	25.63	9.46	9.12	11.28		
11	1.7954	0.4872	23.64	27.59	32.65	35.38	59.49	61.52	55.34		
12	-3.0937	_c	93.89	93.25	10.05	- ^e	14.66	15.63	- ^t		
13	1.6588	0.7513	414.35	667.27	8.75	13.95	4.90	3.95	7.90		
14	4.3414	0.1378	90.87	91.48	4.90	2.67	1.81	2.95	3.00		
15	1.5377	1.9219	5.19	5.27	2.12	3.32	8.20	8.53	27.57		
16	2.4986	-0.2099	228.44	234.29	4.56	2.36	4.84	4.64	10.40		
17	2.8937	0.5155	31.66	38.28	5.54	2.64	4.86	5.23	14.54		
18	0.0087	0.3641	73.20	75.22	17.51	16.93	9.56	13.83	11.31		
19	1.0984	0.4489	40.08	45.04	11.38	7.37	6.04	3.31	5.07		
20	2.4684	0.5045	33.11	37.41	19.86	11.25	20.67	26.62	15.39		
21	1.7000	0.4818	33.35	32.10	10.01	16.05	21.12	25.76	25.54		
22	1.8178	0.4917	9.86	11.72	15.52	9.51	3.13	1.30	1.21		
23	1.4440	0.4550	60.67	64.66	9.71	10.34	17.65	16.59	24.16		
24	3.0471	0.5218	34.78	36.22	5.30	10.94	13.44	17.62	17.20		
25	1.3064	0.4486	65.30	68.74	9.82	4.75	1.57	2.22	1.16		
26	1.7917	0.4631	64.03	68.07	24.30	35.87	5.92	4.40	16.23		
27	2.1727	0.4922	39.59	44.02	10.79	5.56	12.04	16.16	10.43		
28	5.1353	0.5545	56.28	55.25	10.58	40.29	7.37	10.15	6.30		
29	-0.1051	0.2957	21.96	17.94	3.98	3.79	5.69	5.38	1.49		
30	3.4854	-0.5075	693.03	702.27	11.18	6.10	10.44	10.40	18.26		
31	2.1792	0.4925	47.72	53.33	6.80	6.44	11.54	11.65	13.80		
32	-0.5712	-0.1135	24.48	64.50	12.57	9.58	2.42	2.62	2.37		
33	0.8509		40.80	40.39	15.40	- "	3.87	3.90	5.38		
34	-1.0185	0.2444	81.74	83.16	7.18	6.23	5.65	5.70	7.06		
35	2.7149	0.4985	65.68	69.91	2.74	2.95	2.84	2.91	5.03		
36	1.6331	-0.2042	109.96	141.64	3.97	6.62	3.04	3.10	5.28		
37	0.4743	-1.4862	20.91	42.59	19.67	30.56	1.44	3.03	5.45		
38	1.2985		20.27	13.42	1.65		3.84	2.04	10.36		
39	1.7399	1.4625 c	14.03	8.50	8.13	7.88 e	19.10	19.57	36.23		
40	1.9006	 c	23.07	18.55	5.35	- e	1.88	1.10	6.98		
41	1.1601	- c	35.31	32.18	10.93	- e	3.50	2.17	7.80		
42	3.8501	c	23.27	22.23	9.25	- e	1.99	0.77	5.71		
43	5.9322	-	12.60	13.34	2.88	-	2.23	4.89	4.15		
44	2.1910	0.5262	104.03	103.17	3.60	0.14	7.50	0.20	12.95		
45	3.4148	0.5642	245.25	213.01	9.17	8.31	11.38	9.89	11.31		
40	4.9321	0.5973	302.73	200.04	15.70	10.01	23.11	21.30	22.09		
4/ /0	0.0001	0.0043	1 C.00C 20 72	301.00	0.3Z	15.23	2 02	10.00	0.24 0.00		
40 70	-0.0212	0.2921	20.13	33.92 76.02	10.40	10.01	5.0Z	∠.31 1 50	2.20		
49 50	1 6/02	0.4177	216 06	10.33	6 99	13.10	2.40	4.09	4.00		
50	-0 0465	0.4902	210.00 88 70	190.00 80 FF	0.00	13.10 8 56	15 75	2.02	16.25		
	-0.3400	0.2040	460.00	161.00	10.02	14.00	0.04	0.04	10.20		
	-	-	150.09	161.00	10.78	14.63	9.64	9.91	13.83		
PBA''	-	-	1.96	0.00	19.61	32.56	19.61	15.69	16.67		

Table 3 Parameters of the proposed models(Eqs. VI and VII), percent deviation (PD) for each set, mean percent deviation(MPD) of all sets and percent best adherence(PBA) for Eqs. I - VII

^a The set numbers are the same as Table 2, ^b The models constants obtained employing experimental solubility at 25°C and known T_m values by means of Eqs. VIII and IX, ^c The values of G could not be calculated due to $X_2^i < X_2$ at 25 °C. ^d Calculated with the aid of Eq. XIII, ^eNot determined because of $X_2^i < X_2$ at 25 °C, ^f Prediction was not possible due to scarcity of data, ^gCalculated by means of Eq. XIV, ^h Calculated by Eq. XV

REFERENCES

- 1. Yalkowsky SH, Roseman TJ. Solubilization of drugs by cosolvents. In: Yalkowsky SH, ed., Techniques of solubilization of drugs, New York; Marcel Dekker, 1981. p. 91-134.
- 2. Martin A, Newburger J, Adjei A. Extended Hildebrand solubility approach: Solubility of theophylline in polar binary solvents. J Pharm Sci 1980; 69: 487-491.
- 3. Williams NA, Amidon GL. Excess free energy approach to the estimation of solubility in mixed solvent systems: I. Theory. J Pharm Sci 1984; 73: 9-13.
- 4. Acree WE Jr, McCargar JW, Zvaigzne AI, Teng IL. Mathematical representation of thermodynamic properties. Carbazole solubilities in binary alkane + dibutyl ether and alkane + tetrahydropyran solvent mixtures. Phys Chem Liq 1991; 23: 27-35.
- 5. Acree WE Jr. Mathematical representation of thermodynamic properties. Part 2. Derivation of the combined nearly ideal binary solvent (NIBS) /Redlich-Kister mathematical representation from a two-body and three-body interactional mixing model. Thermochimica Acta 1992; 198: 71-79.
- 6. Barzegar-Jalali M, Hanaee J. Model for solubility estimation in mixed solvent systems. Int J Pharm 1994;109: 291-295.
- 7. Barzegar-Jalali M, Jouyban-Gh A. Models for calculating solubility in binary solvent systems. Int J Pharm 1996; 140: 237-246.
- 8. Barzegar-Jalali M, Jouyban-Gh A. A general model from theoretical cosolvency models. Int J Pharm 1997; 152: 247-250.
- 9. Jouyban-Gh A. The modified Wilson model and predicting drug solubility in water cosolvent mixtures. Chem Pharm Bul 1998; 46: 1058-1061.
- 10. Jouyban-Gh A, York P, Hanna M, Clark BJ. Solubility prediction of salmeterol xinafoate in waterdioxane. Int J Pharm 2001; 216: 33-41.
- 11. Grant DJW, Mehdizadeh M, Chow AHL, Fair-Brother JE. Non-linear van't Hoff solubility temperature plots and their pharmaceutical interpretation. Int J Pharm 1984; 18: 25-38.
- 12. Prankerd RJ. Solid-state properties of drugs. I. Estimation of heat capacities for fusion and thermodynamic functions for solution from aqueous solubility-temperature dependence measurements. Int J Pharm 1992; 84: 233-244.
- 13. Regosz A, Pelplinska T, Kowalski P, Thiel Z. Prediction of solubility of sulfonamides in water and organic solvents based on the extended regular solution theory. Int J Pharm 1992; 88: 437-442.
- 14. Yu X, Zipp GL, Davidson III GWR. The effect of temperature and pH on the solubility of quinolone compounds: Estimation of heat of fusion. Pharm Res 1994; 11: 522-527.
- 15. Jouyban-Gh A, Barzegar-Jalali M. Predicting solubility in mixed solvents at various temperatures. AIMECS 97, Seoul, S. Korea, 1997 July 12.
- 16. Jouyban-Gh A, Acree WE Jr. Comparison of models for describing multiple peaks in solubility profiles. Int J Pharm 1998; 167: 177-182.
- Martin AN, Bustamante P, Chun AHC. Physical pharmacy: Physical chemistry principles in the pharmaceutical sciences, 4th ed., Philadelphia, Lea and Febiger, 1993.
- Chertkoff MJ, Martin AN. The solubility of benzoic acid in mixed solvents. J Pharm Sci 1960; 49: 444-447.
- 19. Yalkowsky SH, Amidon GL, Zografi G, Flynn GL. Solubility of nonelectrolytes in polar solvents III: Alkyl p-aminobenzoates in polar and mixed solvents. J Pharm Sci 1975; 64: 48-52.
- 20. Acree WE Jr, Rytting JH. Solubility in binary solvent systems: III. Predictive expressions based on molecular surface area. J Pharm Sci 1983; 72: 292-296.
- 21. Bustamante P; Escalera B; Martin A, Selles E. A modification of the extended Hildebrand approach to predict the solubility of structurally related drugs in solvent mixtures. J Pharm Pharmacol 1993; 45: 253-257.
- 22. Fedors RF. Method of estimating both the solubility parameter and molar volumes of liquids. Poly Eng Sci 1974; 14: 147-154.
- 23. Yalkowsky SH, Valvani SC. Solubility and partitioning I: Solubility of nonelectrolytes in water. J Pharm Sci 1980; 69: 912-922.
- 24. Reichardt C. Solvents and solvent effects in organic chemistry, 2nd ed., Weinheim, Verlag, 1990; 365-371, 408-410.
- 25. Hansen CM, Beerbower A. Solubility parameters. In: Encyclopedia of chemical technology, Suppl. Vol. 1, New York; Wiley; 1970. 324-328.
- Bustamante P, Escalera B. Enthalpy and entropy contributions to the solubility of sulfamethoxypyridazine in solvent mixtures showing two solubility maxima. J Pharm Pharmacol 1995; 47: 550-555.

- 27. Bustamante P, Romero S, Reillo A. Thermodynamics of paracetamol in amphiprotic and amphiproticaprotic solvent mixtures. Pharm. Sci 1995; 1: 505-507.
- 28. Martin A, Miralles MJ. Extended hildebrand solubility approach: Solubility of tolbutamide, acetohexamide, and sulfisomidine in binary solvent mixtures. J Pharm Sci 1982; 71: 439-442.
- 29. Moffat A., Clark's isolation and identification of drugs. 2nd Ed., London; Pharmaceutical Press; 1986. p 986, 988, 993.
- 30. Bustamante C, Bustamante P. Nonlinear enthalpy-entropy compensation for the solubility of phenacetin in dioxane-water solvent mixtures. J Pharm Sci 1996; 85: 1109-1111.
- 31. Subrahmanyam C, Sreenivasa Reddy M, Venkata Rao J, Gaundu Rao P. Irregular solution behaviour of paracetamol in binary solvents. Int J Pharm 1992; 78: 17-24.
- 32. Hildbrand J.H., Jenks C.A., Solubility. IV. Solubility relations of naphthalene and iodine in the various solvents, included a method for evaluating solubility data, J Am Chem Soc 1920; 42: 2180-2189.
- 33. Beerbower A, Wu PL, Martin A. Expanded solubility parameter approach I: naphthalene and benzoic acid in individual solvents. J Pharm Sci 1984; 73: 179-188.
- 34. Reillo A, Cordoba M, Escalera B, Selles, E., Cordoba M Jr. Prediction of sulfamethizole solubility in dioxane-water mixtures. Pharmazie 1995; 50: 472-475.