Antioxidant properties and prevention of cell cytotoxicity of *Phlomis* persica Boiss.

¹Moein S., ^{*1}Farzami B., ¹Khaghani S., ²Moein MR., ³Larijani B.

¹Department of Medical Biochemistry, Medical Sciences/University of Tehran, Tehran,
 ²Department of Pharmacognosy, Medical Sciences/University of Shiraz, Shiraz,
 ³Department of Endocrinology, Medical Sciences/University of Tehran, Tehran, Iran.

Received: 7 Jun 2006; Revised: 6 Aug 2006; Accepted: 27 Aug 2006

ABSTRACT

The free radical scavenging capacity, reducing power and inhibition of β -carotene peroxidation of *Phlomis persica (PP)* extracts was investigated. In addition, the effect of this extract on reduction of the hydrogen peroxide-induced cytotoxicity in non-immortalized fibroblast was examined. The extracts showed free radical scavenging capacity, and the ethyl acetate extract showed marked effect on inhibition of lipid peroxidation similar to that of gallic acid. These results were further supported by a protective effect of *Phlomis* crude extract on H₂O₂-induced cytotoxicity in human non-immortalized fibroblasts.

Keywords: Antioxidant activity; *Phlomis persica;* DPPH scavenging; β -Carotene bleaching test; Cell cytotoxicity.

INTRODUCTION

Free radicals are electrically charged molecules which have unpaired electron and tendency to capture electrons from other substances in order to be neutralized. Although the initial reactions result in neutralization of the free radical, but another free radical is formed

in the process, which by itself causes a chain of reactions to occur and as a result, thousands of free radicals reactions can occur within seconds of the initial reaction (1).

Cell damages caused by free radicals appear to be a major contributor to aging cancer, cardiovascular disease, cataracts, immune system decline, and brain dysfunction (2). Fortunately, free radicals formation is controlled naturally by various beneficial compounds known as antioxidants (3, 4).

Antioxidants are a group of substances when present at low concentrations compared to oxidized substrates significantly inhibit or delay oxidative processes, while being oxidized themselves (5). Phenolic compounds such as flavonoids are ubiquitous in the plant kingdom and approximately 3000 flavonoid substances have been described (6). In addition to an antioxidant effect. flavonoids mav exert protection against heart disease through inhibition of lipooxygenase and cycloxygenase activities in platelets and macrophages (7). Flavonoids have demonstrated to have also been antiinflammatory, anti-allergenic, anti-viral, antiaging and anti-carcinogenic activity (6-9).

The plants of Labiatae's family contain flavonoid compounds (10).One of the labiatae family species is *Phlomis persica*, and in this study the antioxidant activity of its extracts and prevention of cell cytotoxicity of crude extract was investigated.

MATHERIALS AND METHODS

Chemicals

Butylated Hydroxy Toluene (BHT), 1, 1 Diphenyl -2-Picryl Hydrazyl radical (DPPH), Gallic acid, and 3-(4, 5-dimethyl-iazol-2yl), 2.5-diphenyl tetrazolium bromide (MTT) were obtained from Sigma Chemical Co. All other reagents were from Merck Chem. Co. Human non-immortalized fibroblasts (HFFF- P16) were obtained from Pasteur Institute of Iran.

Plant materials

Aerial parts of *Phlomis persica* were collected in March 2005, from the area around Shahre- Kord, Iran, identified and authenticated (Voucher No 125) in the Department of Pharmacognosy, Shaheed Beheshti University of Medical Sciences, Tehran, Iran. The leaves were separated, dried at room temperature and grounded into powder.

Extraction and fractionation

Aerial parts of *Phlomis persica* (3000g) were extracted with ethanol (38.4 L) to give 618.9 g of concentrated crude extract. Four hundreds and thirty grams of the crude extract was then

Measurement of reducing power

The reducing power of extracts of Phlomis persica was determined using the method described by Hartwell (11). A series of concentrated extract of Phlomis persica (12.5, 25, 50, 100 and 200 μ g/ ml) in methanol were made and mixed with 0.2 M phosphate buffer of pH, 6.6 containing 1% ferrocyanate. The mixture was incubated at 50 °C for 20 minutes. To 5 ml of this mixture was added 2.5 ml of 10% trichloroacetic acid and the mixture was centrifuged at 3000 g for 10 minutes. The supernatant was separated and mixed with 2.5 ml of distilled water containing 0.5 ml of 1% ferric chloride. The absorbance of this mixture which could be a measure of antioxidant activity of the extract was determined at 700 nm (11).

Determination of antioxidant using DPPH

The antioxidant activity of plant extract and the standard antioxidants were assessed on the basis of radical scavenging effect of the stable DPPH free radical. In a modified assay (12), 200 μ l of a 100 mM solution of DPPH radical in methanol was mixed with 20 μ l of 12.5-400 μ g/ml extracts, BHT and gallic acid respectively and solutions were left at room temperature for 30 minutes. The DPPH radical inhibition was measured at 490 nm by using a micro-plate reader model Panasonic KX-P108 athos 2020.

The IC_{50} of each sample (concentration in μ g/ml required to inhibit DPPH radical formation by 50%) was calculated by Matlab software. Tests were carried out in triplicates.

The antioxidant activity (AOA) is given by:

100- [(A) sample-(A) blank) \times 100/(A) control]

Where "A" is the absorbance of the samples in wells. DPPH (without plant extract) and methanol were used as control and blank respectively.

β - Carotene bleaching test

For this assay modification of a reported method (13) was employed. One miligram of β - carotene was dissolved in 5 ml of chloroform and to 1 ml of this solution was added 25 µl of linoleic acid and 200 mg of tween 40. Following evaporation of chloroform under vacuum at 40°C, the residue was treated with 50 ml of oxygenated pure water (oxygenation was performed by bubbling air through water for 15 min) and the mixture was vigorously shaken. This emulsion was prepared freshly before each experiment. Stock solution of extract (0.1%) and antioxidants as standards

(0.01%) were prepared in methanol .To each well of 96 well microliter was added 250 μ l of β carotene emulsion in linoleic acid, and 30 μ l of solution of extracts in methanol. An equal amount (30 μ l) of methanol was used as the blank. The microtiter plates were incubated at 55°C for 105 min and the absorbance of samples were measured by a micro plate reader (Panasonic Kx-P108 athos 2020) at 492 nm (14). The antioxidant activity coefficient (ACC) was given by a modified version of the formula of Chevolleau et.al (15)

$$ACC = [(A_{A}105 - A_{B}105)/(A_{B}0 - A_{B}105)]$$

In which $A_A 105$ and $A_B 105$ are the absorbancies of the test and blank samples at zero and 105 min respectively and $A_B 0$ is the absorbance of the blank sample at t=0 min.

Determination of the phenolic compounds

The content of total phenolic compounds in methanolic solutions of the crude extract was determined by a described method (16). For the preparation of calibration curve, 1ml aliquots of 0.024, 0.075, 0.105 and 0.3mg/ml gallic acid solutions in methanol were mixed with 5ml of folin-ciocalteu reagent (diluted ten fold) and 4ml (75g/l) of sodium carbonate. The absorption was read after 30 min at 20 °C at 765 nm and the calibration curve was constructed. One ml of methanolic solution of the crude extract (10 g/l) was mixed with the same reagents as described above, and after 1 hour the absorption was measured for determination of plant phenolic componds. All determinations were performed in triplicates. Total content of the phenolic compounds in methanolic solution of the crude extract equivalent to gallic acid (GAE) was calculated from the formula:

C = c.V/m'

Where C is total content of phenolic compounds in mg/g of the extract, equivalent to GAE; c is the concentration of gallic acid which is found from the calibration curve in mg /ml; V is the volume of extract in ml; and m is the weight of the plant crude extract.

Cell culture and treatments

Human non –immortalized fibroblast (HFFF-P16) cell were obtained from Pasteur institute. The cells were grown at 37° C in humidified 5% CO₂ 95% air mixture in DMEM (Dulbecco's modified Eagle medium) supplemented with 15% fetal calf serum(FCS), 1m of L-glutamine, 100U/ml of penicillin and streptomycin. The medium was replaced 3 times in a week until the cells were near confluency. The crude extract of *Phlomis persica* was dissolved in the minimum amount of

dimethyl sulphoxide (DMSO) in a way that the final DMSO concentration was never higher than 0.05% when mixed with culture medium. DMSO at this concentration has not been reported toxic for the cells. The solution of crude extracts in DMSO (12.5-25 μ g/ml) were added to the culture medium and kept at incubator for 24 hours, solution of gallic acid in DMSO (6.25 μ g/ml) was used as a standard. After 24 hours the cells in plates (32×10⁴/ml) were treated with 0.1 mM H₂O₂ for 30 minutes and then washed with buffer phosphate.

MTT bioassay

MTT bioassay measures the conversion of tetrazolium salt to the colored product, the concentration of which can be measured spectrophotometrically (17). Twenty μ l of 0.5% MTT in PBS was added to each micro well. The microplate was put in incubator (37°C) for 6 hours, the supernatant was removed and replaced with 200 μ L of acidified isopropranol and 20 μ L of 3% (W/V) dodecyl sulphate in water.

The optical density of each sample was measured with a micro plate spectrophotometer reader

(Panasonic Kx-P108 athos 2020) at 570 nm. Five replicates were performed for each sample. Reduction in H_2O_2 cytotoxicity were determined and expressed in percent by crude extract and gallic acid as standard.

Statistical evaluation

Mean ±SD are given. The data were analyzed for statistical significance using one way ANOVA followed by Tukey or Dunnett post test. P values less than 0.05 were considered significant.

RESULTS

Reducing power capacity

Fig (1) shows the reducing power of the crude, butanolic and ethyl acetate extracts of *Phlomis persica*, and BHT and gallic acid as standards at equal concentrations of $12.5 \,\mu\text{g}$ /ml to $200 \,\mu\text{g}$ /ml.

Radical scavenging activity

Fig (2) depicts radical scavenging effect of the crude, buthanolic and ethyl acetate extracts of *Phlomis persica* (PP) and BHT and gallic acid as standards in concentrations of 12.5 to 400 µg/ml. The IC₅₀ of BHT and gallic acid were found to be 68 and 2.49 µg/ml respectively. The IC₅₀ of the ethyl acetate extract (116.2±1.5 µg/ml) was less than butanolic extract (148.1±3.5 µg/ml).

β -carotene bleaching test

The results of β - carotene bleaching test are illustrated in table 2. The ACC of ethyl acetate

extract of *Phlomis* was more than *Phlomis* crude extract and butanolic fraction (P < 0.001).

Table 1.	IC ₅₀ of	antioxidan	t compounds	and	Phlomis
extracts co	mpared	with BHT	and Gallic ac	id	

Antioxidant compounds	IC ₅₀ µg/ml ^a
Gallic acid	$2.49 \pm .6$
BHT	63.17±.9
Phlomis ethyl acetate extract	116.2±1.5
Phlomis butanolic extract	148.1 ± 3.5
Phlomis crude extract	169.8 ± 5

a: Each value reperesents the mean \pm SD

The IC_{50} of ethyl acetate extract was less than butanolic extract and crude extract (P < 0.05).

Table 2. Antioxidant activity of Phlomis extracts	in
comparison with BHT and gallic acid as measured	by
the BCBT ^a method.	

Antioxidant compound	ACC ^b
ВНТ	866.6±15.7
gallic acid	754.03±44.28
Phlomis crude extract	218.27±26.5
Phlomis butanolic extract	350.6±24.4
Phlomis ethyl acetate extract	656.54±30.56

a) BCBT (β -Carotene Bleaching Test).

b) Antioxidant activity coefficient by BCBT method given as ACC.

Table 3. Effects of *Phlomis* extracts on H_2O_2 induced cytotoxicity in non immortalized fibroblasts. MTT (% of control) % Inhibition of H_2O_2 cytotoxicity

Control	100
H_2O_2 (0.1mM)	$45.2 \pm 2.9^{\mathrm{a}}$
Gallic acid 6.25 µg/ml	92.4 ± 2.8^{b}
Phlomis extract	
12.5 µg/ml	72.8 ± 3.3 ^b
25.0 µg/ml	$80.2 \pm 4.5^{ m b}$

^a Each value represents the mean \pm SD of five experiments significant vs control untreated fibroblasts and ^b vs H₂O₂ treated cell (P< 0.01).

Cell culture

The results of H_2O_2 cytotoxicity in fibroblasts cells was evaluated on the basis of its effects on cell growth factor (MTT test) and are listed in Table 3. Cell growth was significantly suppressed in the H_2O_2 treated cells. Addition of the crude extract (12.5 and 25µg) before exposure to peroxide mitigated the cytotoxic response and this effect was concentration dependent. Under the same conditions gallic acid (6.25 µg/ml) reduced the cytotoxicity of H_2O_2 . The extract of *Phlomis persica* at concentrations of 12.5, 25, µg/ml, were not cytotoxic.

DISCUSSION

Species of labiatae's due to their distribution and diversity in the Mediterranean region and their essential oils are a point of interest from the botanic, taxonomic, chemical composition and ecological aspects (18).



Figure 1. Reducing power of *Phlomis persica* compared with antioxidant standards BHT and gallic acid (gallic acid is diluted 1:10).



Figure 2. Radical scavenging activity of *Phlomis* extracts compared with BHT and gallic acid (gallic acid is diluted 1:10). Radical scavenging activity of *Phlomis* ethyl acetate extract is more than its butanolic extract (P < 0.05).

Phlomis is one of the labiatae species and the genus *Phlomis* is composed of about 100 species, of which 17 species are described in flora of Iran, among them 10 are endemic (19, 20). Recently isolation of irridiosl (21) and phenyl propanoid glycoside (22) from the oil of *Phlomis fruticosa* (21), have been reported. Of these compounds Phenyl propanoid glycoside has shown antioxidant activity (25).

In other investigations, 4 flavonol glycosides were isolated from a methanolic extract of *Phlomis spectabilis* (26). In this investigation the antioxidant activity of *Phlomis persica* was investigated.

The antioxidant potential of a compound can be attributed to its reducing power, radical scavenging ability and prevention of β -carotene bleaching. These properties may serve as

significant indicators of its potential as antioxidant. However the activity of putative antioxidants has been attributed to various mechanisms, among which are prevention of chain initiation, binding to transition metal catalysis, decomposition of peroxides (27). Our experiments using DPPH and reducing power tests showed that there is a strict proportionality between the concentration of extracts as antioxidant and BHT and gallic acid as standards. The reducing power of *Phlomis persica* and its fractions increased by increase in their concentrations (Fig.1). However their reducing powers were lower than BHT and gallic acid. Table 1 shows that IC_{50} of the ethyl acetate extract was less than butanolic and crude extracts. It is generally accepted that phenolic compounds

which are present in fractions of plant extract

inhibit the oxidation by donation of hydrogen atoms to scavenge free radicals (28).Furthermore, it is reported that the ethyl acetate extract is more active than other fractions in radical scavenging properties (28). Crude extract of *Phlomis* and its fractions (containing phenolic compounds) can inhibit β - carotene bleaching and there is a significant difference in ACC between standards and butanolic extract (P<0.001, F=424.06). Also a significant difference in ACC between BHT and ethyl acetate extract was observed (P<0.001, F=424.06).

The amount of phenolic compounds of *Phlomis* crude extract and its ethyl acetate extract were 28.8 ± 0.8 and 40 ± 0.1 mg/g respectively. The higher antioxidant activity of the ethyl acetate extract may be related to its higher phenolic compounds.

In this study, the cytotoxic effects of H_2O_2 on non -immortalized fibroblast cells were evaluated by inhibition of the cell growth which was significantly suppressed in the H_2O_2 -treated cells (Table-3). Addition of *Phlomis persica* extract before exposure to H_2O_2 mitigated the cytotoxic response and it was observed that this effect was concentration dependent. In other words a significant difference (P<0. 01, F=168.89) was observed between non-immortalized fibroblasts which were treated only with H_2O_2 and the cells which were treated with crude extract or gallic acid before exposure to H_2O_2 .

In conclusion, the results of this study indicate that this extract exhibits interesting antioxidant properties expressed by its capacity to scavenge free radicals (DPPH), prevention of lipid peroxidation, reducing ability and reduction of H_2O_2 –induced cytotoxicity in human non immortalized fibroblast cells.

ACKNOWLEDGMENT

The financial support of the Endocrinology and Metabolism Reseach Center is greatly appreciated. Also thanks are due to Mr. Kamali – Nejad (Department of pharmacognosy Shaheed Beheshti University of Medical Sciences, Tehran, Iran) for his assistance in identifying the species of plant (*Phlomis*) in this research.

REFERENCES

- 1. Percival M. Antioxidants. Clin Nutr Insights 1998; 1-4.
- 2. Sies H. Antioxidant functions of vitamins. Ann Ny Acad Sci 1992; 669:7-20.
- 3. Langseth L. Antioxidants and diseases of the brain. Antioxidant vitamins. Newslett 1993; 4: 30.
- 4. Halliwell B. Free radicals, antioxidants and human disease: curiosity, cause, or consequence. Lancet 1994; 344:721-724.
- 5. Vaya J, Aviram M. Nutritional antioxidant: Mechanism of action, analysis of activities and medical applications.Phytother Res 2003; 8(6):1-17.
- 6. Middleton COV, Harbored, EJB. Plant flavonoid in biology and medicine- biochemical, pharmacological and structure-activity relationships, Alan R. Liss, Newyork; 1986. P. 28.
- 7. Havsteen B. Flavonoids, a class of natural products of high pharmacological potency. Biochem Pharm 1983; 32(7): 1141-1148
- Kuhana J. The flavonoids; a class of semi –essential food components: their role in human nutrition. Nutr Diet 1976; 24: 117-191
- 9. Middleton E. The flavonoids, Trends in Pharmaceut. Science 1984; 5: 335-380
- 10. Gill FA, Barberan F F, Tomas F. Distribution of flavonoid glucosides in members of the subfamily lamioideae. Lamiales newslett1992; 1: 14
- 11. Hartwell JL. Plants used against cancer. loydia 1969; 32: 247-290
- 12. Bruits M, Asres k, Bucar F. The antioxidant activity of the essential oils of Artemisia Afra, Artemisia byssinica and juniperus procera. Phytother Res 2001; 15: 103- 108.
- 13. Dapkevicius A, venskutonis R, Van Beek TA, linssen JPH. Antioxidant activity of extracts obtained by different isolation. J Sci Food Agric 1998; 77: 140- 146.
- 14. Pratt, DE. Natural antioxidants of soybeans and other oil seeds, In Autooxidation in Food and Biological Systems, Simic MG, Karel M (eds). Plenum:New York. 1980. P. 283-293.
- 15. Chevolleau S, Mallet JF, Ucciani E, Gamisans J, Gruber M. Antioxidant activity in leaves of soma Mediterranean plants. J Am Oil Chem Soc 1992; 691: 1269- 1271.
- 16. Miliauskas G, Venskutonis PR, van Beek TA. Screening of radical scavenging activity of some medicinal plants and aromatic plant extract. Food Chem 2004; 85: 231-237.
- 17. Mosmann T. Rapid colorimetric assay for cellular growth and survival, application to proliferation and cytotoxicity assay. J Immunol Meth 1983; 65: 55- 63.
- 18. Gonales MR, Tejero J, Molero M, Casares MP. The labiatea family in popular medicine eastern, Andalucia. Lamiales newslett. 1992; 1: 27.
- 19. Rechinger KH. Flora Iranica. Graz: Akademische Druck Verlagsanstalt ;1982. P. 298-310.

- 20. Mozafarian V. A dictionary of Iranian plants name. Tehran: Farhang Mo'aser publisher; 1996. P. 450.
- 21. Bucar F, Ninov S, Ionkova IK, Schubert-Zsilavecza T M, Asenov I. Flavonoids from Phlomis nissolii. Phytochemistry 1998; 48: 573-574.
- 22. Kasai R, Katagiri M, Ohtani k, Yamasaki, Yang CR, Tanaka O. Iridoid glucosides from Phlomis vounghusbandii roots. Phytochemistry 1994; 36: 967-970.
- 23. Tsitsimi E, Loukis, verykokidou E. Composition of the essential oil of the flowers of phlomis fruticosa L.from Greece. J Essent Oil Res 2000; 12: 355-356.
- 24. Calis IAA, Basaran I, Saracoglu O, Sticher p, Plinosides R. Phenyl propanoid glycosides and iridoids Phenyl propanoid glycosides and iridoids from Phlomis Linearis. Phtyochemistry 19991; 30: 3075-3075.
- 25. Grace SC, Logan BA, Adams WW. Seasonal differences in foliar content of chlorogenic acid, a phenyl propanoid antioxidant, Mahonia repens Plant. Cell Environ. 1998; 21: 513- 521.
- 26. Rajive K, Satesh B, Ashok K. Flavonol glycoside of Phlomis spectabilis. Phytochemistry. 1985; 24(5):1124-1125.
- 27. Mensor LL, Menezes SF, Leitao GG, Reis S A, Santos C T, Coube SC. Antioxidant activity in Brazilian plants. Phytother Res. 2001; 15: 127-130.
- 28. Weng XC. Antioxidant activity and components of Salvia plebeian R.Br. a Chinese herb. Food Chem. 2001; 73: 299-305.