



RESEARCH ARTICLE

Open Access

Antinociceptive properties of new coumarin derivatives bearing substituted 3,4-dihydro-2H-benzothiazines

Masoumeh Alipour¹, Mehdi Khoobi², Saeed Emami³, Saeed Fallah-Benakohal², Seyedeh Farnaz Ghasemi-Niri², Mohammad Abdollahi⁴, Alireza Foroumadi² and Abbas Shafiee^{2*}

Abstract

Background: Coumarins are an important class of widely distributed heterocyclic natural products exhibiting a broad pharmacological profile. In this work, a new series of coumarins bearing substituted 3,4-dihydro-2H-benzothiazines were described as potential analgesic agents. The clinical use of NSAIDs as traditional analgesics is associated with side effects such as gastrointestinal lesions and nephrotoxicity. Therefore, the discovery of new safer drugs represents a challenging goal for such a research area.

Results: The target compounds 3-(3-methyl-3,4-dihydro-2H-benzo[6][1,4]thiazin-3-yl)-2H-chromen-2-ones **2a-u** were synthesized and characterized by spectral data. The antinociceptive properties of target compounds were determined by formalin-induced test and acetic acid-induced writhing test in mice. Among the tested compounds, compound **2u** bearing 2-(4-(methylsulfonyl)benzoyl)- moiety on benzothiazine ring and 4-(methylsulfonyl)phenacyloxy- group on the 7 position of coumarin nucleus showed better profile of antinociception in both models. It was more effective than mefenamic acid during the late phase of formalin-induced test as well as in the acetic acid-induced writhing test.

Conclusion: Considering the significant antinociceptive action of phenacyloxy coumarin derivatives, compound **2u** prototype might be further used as model to obtain new more potent analgesic drugs.

Keywords: Analgesic activity, Antinociception, Coumarin, Benzothiazine, Formalin test, Writhing test

Introduction

Pain is an uncomfortable sensation that alerts the human organs about a current or potential damage to tissues [1]. It has been accepted that pain can widely affect human life quality, and its management is considered as a main challenge in pharmacotherapy [2]. NSAIDs are one of major classes of traditional analgesics for treatment of pain. The clinical use of NSAIDs is associated with side effects such as gastrointestinal lesions and nephrotoxicity [3]. Therefore, the discovery of new safer drugs represents a challenging goal for such a research area.

Coumarins are an important class of widely distributed heterocyclic natural products exhibiting a broad pharmacological profile [4]. Several coumarin derivatives have been synthesized with diverse biological activities [5-9] especially analgesic/anti-inflammatory activity [10-13]. Recently, the synthesis and anti-inflammatory/analgesic activities of several coumarin derivatives with various substitutions on 3-position of coumarin nucleus have been reported [14-16]. On the other hand, benzothiazine derivatives are also important heterocyclic compounds with wide spectrum of biological activities [17,18]. In view of the above facts and in continuation of our research program on the synthesis of biologically active heterocyclic compounds [19,20], we introduced herein the new coumarin derivatives bearing substituted 3,4-dihydro-2H-benzothiazines as analgesic agents. The antinociceptive properties of target compounds were determined by formalin-induced paw licking test and

* Correspondence: ashafiee@ams.ac.ir

²Department of Medicinal Chemistry, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medical Sciences, Tehran 14176, Iran

Full list of author information is available at the end of the article

acetic acid-induced writhing test in mice. Indeed, the formalin-induced paw licking method is used to investigate both peripheral and central mechanisms whereas the acetic acid test is believed to demonstrate the involvement of peripheral mechanisms in the control of pain [21,22].

Materials and methods

Chemistry


The target compounds 3-(3-methyl-3,4-dihydro-2*H*-benzo[*b*][1,4]thiazin-3-yl)-2*H*-chromen-2-ones **2a-r** (Additional file 1: Table S1) were synthesized according to the pathway outlined in Scheme 1 [23]. All reagents and chemicals were commercially available and used as received. Alumina-supported potassium fluoride (KF/Al₂O₃) was prepared by literature method [24]. The dihydrobenzothiazole derivatives **1** were prepared as reported method by us [19,20]. The synthesis of compounds **2a-d**, **2f-i** and **2p-r** was described in our previous paper [23]. Column chromatography was carried out on silica gel (70–230 mesh). TLC was conducted on silica gel 250 micron, F254 plates. Melting points were measured on a Kofler hot stage apparatus and are uncorrected. The IR spectra were taken using Nicolet FT-IR Magna 550 spectrographs (KBr disks). ¹H NMR spectra were recorded on a Bruker 400 or 500 MHz NMR instruments. The chemical shifts (δ) and coupling constants (*J*) are expressed in parts per million and Hertz, respectively. Mass spectra of the products were obtained with an HP (Agilent technologies) 5937 Mass Selective Detector. Elemental analyses were carried out by a CHN-Rapid Heraeus elemental analyzer. The results of elemental analyses (C, H, N) were within ± 0.4% of the calculated values.

General procedure for the synthesis of compounds **2**

A suspension of dihydrobenzothiazole derivatives **1** (1.0 mmol), KF/Al₂O₃ (0.7 g), and quinine hydrochloride (10 mol%) in ethanol (3.0 mL) was stirred at room temperature for 5 min. Then, appropriate phenacyl halide (1.2 mmol) was added to the mixture and stirring was continued. After completion of the reaction (3–5 h), the solvent was removed under reduced pressure. The residue was mixed with ethyl acetate (5 mL) and the catalyst was filtered and washed with ethyl acetate (3 × 5 mL). After evaporation of the solvent at reduced pressure, the crude product was purified by column chromatography (*n*-hexane/ethyl acetate, 9:1) and crystallized from ethanol for further purification.

3-(2-(3,4-Dichlorobenzoyl)-3-methyl-3,4-dihydro-2*H*-benzo[*b*][1,4]thiazin-3-yl)-2*H*-chromen-2-one (**2e**)

Yellow solid (361 mg, 75%); *syn*-isomer; mp 91–93°C; IR (KBr, cm⁻¹) 3382 (NH), 1708 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 1.94 (s, 3H, CH₃ benzothiazine), 4.49 (s, 1H, NH), 5.77 (s, 1H, C-H benzothiazine), 6.75 (dt, *J* = 7.2 and 1.2 Hz, 1H, H₇ benzothiazine), 6.95 (m, 2H, H_{5,8} benzothiazine), 7.14 (dt, *J* = 7.2 and 1.2 Hz, 1H, H₆ benzothiazine), 7.22 (t, *J* = 8.0 Hz, 1H, H₆ chromene), 7.28 (dd, *J* = 8.0 and 1.9 Hz, 1H, H₅ benzoyl), 7.33 (d, *J* = 8.0 Hz, 1H, H₆ benzoyl), 7.40 (m, 2H, H_{5,8} chromene), 7.43 (d, *J* = 1.9, 1H, H₃ benzoyl), 7.49 (dt, *J* = 8.0 and 1.2 Hz, 1H, H₇ chromene), 7.77 (s, 1H, H₄ chromene); ¹³C NMR (125 MHz, CDCl₃) δ 24.3, 42.9, 58.0, 110.9, 116.1, 117.1, 119.1, 119.3, 124.3, 127.0, 127.1, 128.4, 128.7, 130.1, 130.8, 131.2, 131.4, 132.0, 136.9, 137.1, 139.8, 141.0, 153.2, 160.9, 192.6; Anal. calcd for



Scheme 1 Synthesis of coumarin based dihydrobenzothiazines **2a-u**. Reagents and conditions: (a) phenacyl halide (1.2 mmol), KF/Al₂O₃ (0.7 g), quinine hydrochloride (10 mol%), EtOH (3 mL), r.t. (b) phenacyl halide (2.5 mmol), KF/Al₂O₃ (1.5 g), quinine hydrochloride (10 mol%), EtOH (3 mL), r.t.

C₂₅H₁₇Cl₂NO₃S: C, 62.25; H, 3.55; N, 2.90. Found: C, 62.41; H, 3.67; N, 3.15.

3-(2-(4-Fluorobenzoyl)-3-methyl-3,4-dihydro-2H-benzo[b][1,4]thiazin-3-yl)-2H-chromen-2-one (2j)

Yellow solid (336 mg, 78%); *syn*-isomer; mp 161–163°C; IR (KBr, cm⁻¹) 3413 (NH), 1708 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 1.76 (s, 3H, CH₃ benzothiazine), 4.50 (s, 1H, NH), 5.97 (s, 1H, C-H benzothiazine), 6.73 (t, *J* = 7.4 Hz, 1H, H₇ benzothiazine), 6.93 (m, 2H, H_{5,6} benzothiazine), 7.14 (m, 3H, H₈ benzothiazine and H_{3,5} benzoyl), 7.28 (t, *J* = 7.4 Hz, 1H, H₆ chromene), 7.35 (d, *J* = 7.4 Hz, 1H, H₈ chromene), 7.40 (d, *J* = 7.4 Hz, 1H, H₅ chromene), 7.50 (t, *J* = 7.4 Hz, 1H, H₇ chromene), 7.79 (s, 1H, H₄ chromene), 8.05 (m, 2H, H_{2,6} benzoyl); ¹³C NMR (125 MHz, CDCl₃) δ 24.6, 37.5, 57.6, 111.9, 115.7, 115.9, 116.1, 119.1, 119.4, 124.4, 126.7, 128.4, 128.5, 131.2, 131.3, 131.5, 133.2, 139.5, 141.1, 153.3, 161.3, 164.7, 166.7, 191.3; Anal. calcd for C₂₅H₁₈FNO₃S: C, 69.59; H, 4.20; N, 3.25. Found: C, 69.42; H, 4.03; N, 3.47.

3-(3-Methyl-2-(thiophene-2-carbonyl)-3,4-dihydro-2H-benzo[b][1,4]thiazin-3-yl)-2H-chromen-2-one (2k)

Yellow solid (356 mg, 85%); as mixture of diastereomers (*anti/syn*: 15/85); IR (KBr, cm⁻¹) 3389 (NH), 1707 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 1.77_{syn} (s, CH₃ benzothiazine), 1.87_{anti} (s, CH₃ benzothiazine), 4.48_{syn} (s, NH), 4.53_{anti} (s, NH), 5.50_{anti} (s, C-H benzothiazine), 5.87_{syn} (s, C-H benzothiazine), 6.77_{syn} (t, *J* = 8.0, H₇ benzothiazine), 6.81_{anti} (t, *J* = 8.0, H₇ benzothiazine), 6.92_{anti} (d, *J* = 8.0 Hz, H₅ benzothiazine), 6.95_{syn} (d, *J* = 8.0 Hz, H₅ benzothiazine), 7.07–7.10_{anti} (m, H_{6,8} benzothiazine), 7.10–7.13_{syn} (m, H_{6,8} benzothiazine), 7.20_{syn} (t, *J* = 7.6 Hz, H₇ chromene), 7.25_{syn} (t, *J* = 7.6 Hz, H₆ chromene), 7.25–7.29_{anti} (m, H₄ thiophene and H₇ chromene), 7.38_{syn} (d, *J* = 7.6 Hz, H₅ chromene), 7.42_{syn} (d, *J* = 7.6 Hz, H₈ chromene), 7.48–7.54 (m, H_{5,7,8} chromene (*anti*) and H₄ thiophene (*syn*)), 7.56_{anti} (d, *J* = 4.0, H₃ thiophene), 7.74_{syn} (d, *J* = 4.0, H₃ thiophene), 7.79_{anti} (d, *J* = 4.0, H₅ thiophene), 7.80_{syn} (s, H₄ chromene), 7.97_{syn} (d, *J* = 4.0, H₅ thiophene), 8.15_{anti} (s, H₄ chromene); ¹³C NMR (*syn*-isomer, 125 MHz, CDCl₃) δ 27.5, 45.3, 54.2, 116.1, 117.0, 118.6, 119.0, 120.1, 124.5, 126.2, 127.1, 128.2, 128.4, 131.4, 131.7, 132.2, 134.4, 140.2, 140.9, 143.5, 153.1, 160.1, 186.8; Anal. calcd for C₂₃H₁₇NO₃S₂: C, 65.85; H, 4.08; N, 3.34. Found: C, 65.92; H, 3.91; N, 3.29.

3-(2-(5-Bromothiophene-2-carbonyl)-3-methyl-3,4-dihydro-2H-benzo[b][1,4]thiazin-3-yl)-2H-chromen-2-one (2l)

Yellow solid (378 mg, 77%); as mixture of diastereomers (*anti/syn*: 26/74); IR (KBr, cm⁻¹) 3390 (NH), 1712 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 1.75_{syn} (s, CH₃ benzothiazine), 1.87_{anti} (s, CH₃ benzothiazine), 4.45_{syn} (s, NH), 4.55_{anti} (s, NH), 5.48_{anti} (s, C-H benzothiazine), 5.78_{syn} (s, C-H benzothiazine), 6.76_{syn} (t, *J* = 8.0, H₇ benzothiazine),

6.81_{anti} (t, *J* = 8.0, H₇ benzothiazine), 6.92_{syn} (d, *J* = 8.0 Hz, H₅ benzothiazine), 6.94_{anti} (d, *J* = 8.0 Hz, H₅ benzothiazine), 6.92_{anti} (d, *J* = 8.0 Hz, H₈ benzothiazine), 6.96_{syn} (d, *J* = 8.0 Hz, H₈ benzothiazine), 7.09_{anti} (t, *J* = 8.0 Hz, H₆ benzothiazine), 7.12_{syn} (t, *J* = 8.0 Hz, H₆ benzothiazine), 7.16_{syn} (d, *J* = 4.0 Hz, H₄ thiophene), 7.23_{anti} (d, *J* = 4.0 Hz, H₄ thiophene), 7.24_{syn} (t, *J* = 7.5, H₆ chromene), 7.27–7.29_{anti} (m, H_{6,8} chromene), 7.36_{syn} (d, *J* = 7.5 Hz, H₈ chromene), 7.40_{syn} (d, *J* = 7.5 Hz, H₅ chromene), 7.50_{syn} (t, *J* = 7.5 Hz, H₇ chromene), 7.51–7.53_{anti} (m, H_{5,7} chromene), 7.56_{anti} (d, *J* = 4.0 Hz, H₃ thiophene), 7.69_{syn} (d, *J* = 4.0 Hz, H₃ thiophene), 7.77_{syn} (s, H₄ chromene), 8.15_{anti} (s, H₄ chromene); ¹³C NMR (*syn*-isomer, 125 MHz, CDCl₃) δ 24.8, 38.7, 57.4, 112.3, 116.1, 116.9, 119.1, 119.5, 123.6, 124.4, 126.6, 128.1, 128.4, 130.8, 131.5, 131.6, 132.4, 139.2, 141.0, 145.9, 153.3, 161.2, 185.6; MS, *m/z* (%) 499 ([M + 2]⁺, 40%), 497 (M⁺, 37%), 375 (47%), 373 (44%), 308 (100%), 294 (51%), 280 (84); Anal. calcd for C₂₃H₁₆BrNO₃S₂: C, 55.43; H, 3.24; N, 2.81. Found: C, 55.22; H, 3.47; N, 2.73.

3-(3-Methyl-2-(thiophene-3-carbonyl)-3,4-dihydro-2H-benzo[b][1,4]thiazin-3-yl)-2H-chromen-2-one (2m)

Yellow solid (335 mg, 80%); as mixture of diastereomers (*anti/syn*: 32/68); IR (KBr, cm⁻¹) 3374 (NH), 1708 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 1.77_{syn} (s, CH₃ benzothiazine), 1.87_{anti} (s, CH₃ benzothiazine), 4.50_{anti} (s, NH), 4.55_{syn} (s, NH), 5.48_{anti} (s, C-H benzothiazine), 5.81_{syn} (s, C-H benzothiazine), 6.74_{syn} (t, *J* = 7.5, H₇ benzothiazine), 6.80_{anti} (t, *J* = 7.5, H₇ benzothiazine), 6.93_{anti} (d, *J* = 7.5 Hz, H₅ benzothiazine), 6.97_{syn} (d, *J* = 7.5 Hz, H₅ benzothiazine), 7.08_{anti} (d, *J* = 7.5 Hz, H₈ benzothiazine), 7.12_{syn} (t, *J* = 7.5 Hz, H₆ benzothiazine), 7.13_{syn} (t, *J* = 7.2 Hz, H₆ chromene), 7.24–7.25_{syn} (m, H₈ benzothiazine and H₈ chromene), 7.26–7.29_{anti} (m, H₆ benzothiazine and H_{6,8} chromene), 7.35–7.37_{syn} (m, H₄ thiophene and H_{5,7} chromene), 7.37–7.40_{anti} (m, H₆ chromene and H₄ thiophene), 7.50–7.53_{anti} (m, H₇ chromene and H₅ thiophene), 7.58_{syn} (d, *J* = 5.0 Hz, H₅ thiophene), 7.79_{anti} (s, H₂ thiophene), 8.04_{syn} (s, H₂ thiophene), 8.14_{anti} (s, H₄ chromene), 8.25_{syn} (s, H₄ chromene); ¹³C NMR (*syn*-isomer, 125 MHz, CDCl₃) δ 27.8, 45.7, 54.4, 116.1, 116.8, 119.1, 119.9, 124.6, 126.3, 126.5, 127.3, 128.2, 128.4, 130.6, 131.5, 131.7, 132.6, 140.0, 140.2, 141.2, 153.1, 160.2, 187.7; MS, *m/z* (%) 419 (M⁺, 68%), 404 (12%), 386 (12%), 308 (97%), 295 (100%), 280 (64%), 111 (63); Anal. calcd for C₂₃H₁₇NO₃S₂: C, 65.85; H, 4.08; N, 3.34. Found: C, 65.98; H, 3.82; N, 3.60.

8-Methoxy-3-(3-methyl-2-(4-methylbenzoyl)-3,4-dihydro-2H-benzo[b][1,4]thiazin-3-yl)-2H-chromen-2-one (2n)

Yellow solid (343 mg, 75%); *syn*-isomer; mp 145–147°C; IR (KBr, cm⁻¹) 3360 (NH), 1700 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 1.77 (s, 3H, CH₃ benzothiazine), 2.43 (s, 3H, CH₃ benzoyl), 3.90 (s, 3H, O-CH₃ chromene), 4.52 (s, 1H, NH), 6.00 (s, 1H, C-H benzothiazine), 6.78 (dt, *J* = 7.5 and

1.3 Hz, 1H, H₇ benzothiazine), 6.93 (dd, $J = 8.0$ and 1.3 Hz, 1H, H₇ chromene), 7.02-7.11 (m, 4H, H_{5,6,8} benzothiazine and H₆ chromene), 7.18 (m, 3H, H₅ chromene and H_{3,5} benzoyl), 7.74 (d, $J = 8.3$ Hz, 2H, H_{2,6} benzoyl), 8.13 (s, 1H, H₄ chromene); ¹³C NMR (125 MHz, CDCl₃) δ 21.6, 27.9, 43.6, 54.6, 56.1, 113.3, 114.8, 118.9, 119.4, 119.7, 119.9, 124.3, 126.3, 127.4, 128.5, 129.3, 131.1, 133.7, 140.0, 140.2, 142.8, 143.8, 146.7, 159.5, 192.9 cm⁻¹; Anal. calcd for C₂₇H₂₃NO₄S: C, 70.88; H, 5.07; N, 3.06. Found: C, 70.64; H, 5.23; N, 3.22.

3-(2-(4-Fluorobenzoyl)-3-methyl-3,4-dihydro-2H-benzo[b][1,4]thiazin-3-yl)-8-methoxy-2H-chromen-2-one (2o)

Yellow solid (323 mg, 70%); *syn*-isomer; mp 236–238°C; IR (KBr, cm⁻¹) 3398 (NH), 1690 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 1.78 (s, 3H, CH₃ benzothiazine), 3.98 (s, 3H, O-CH₃ chromene), 4.51 (s, 1H, NH), 5.98 (s, 1H, C-H benzothiazine), 6.74 (t, $J = 7.4$ Hz, 1H, H₇ benzothiazine), 6.93 (m, 2H, H_{5,6} benzothiazine), 6.98 (d, $J = 8.0$ Hz, 1H, H₇ chromene), 7.05 (m, 2H, H_{5,6} chromene), 7.16 (m, 3H, H₈ benzothiazine and H_{3,5} benzoyl), 7.77 (s, 1H, H₄ chromene), 8.05 (m, 2H, H_{2,6} benzoyl); ¹³C NMR (125 MHz, CDCl₃) δ 24.5, 37.4, 56.3, 57.6, 111.9, 113.3, 115.7, 115.9, 116.9, 119.3, 119.8, 124.2, 126.7, 128.4, 131.2, 131.3, 131.4, 133.1, 139.5, 141.2, 146.8, 160.7, 164.7, 166.7, 191.2; Anal. calcd for C₂₆H₂₀FO₄S: C, 67.67; H, 4.37; N, 3.04. Found: C, 67.43; H, 4.18; N, 3.25.

3-(2-(4-Bromobenzoyl)-3-methyl-3,4-dihydro-2H-benzo[b][1,4]thiazin-3-yl)-7-(2-(4-bromophenyl)-2-oxoethoxy)-2H-chromen-2-one (2s)

Yellow solid (507 mg, 72%); as mixture of diastereomers (*anti/syn*: 18/82); IR (KBr, cm⁻¹) 3382 (NH), 1697 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 1.74_{syn} (s, CH₃ benzothiazine), 1.85_{anti} (s, CH₃ benzothiazine), 4.48_{anti} (s, NH), 4.51_{syn} (s, NH), 5.29_{anti} (s, O-CH₂), 5.31_{syn} (s, O-CH₂), 5.60_{anti} (s, C-H benzothiazine), 5.91_{syn} (s, C-H benzothiazine), 6.71_{anti} (t, $J = 7.5$, H₇ benzothiazine), 6.73_{syn} (t, $J = 7.5$, H₇ benzothiazine), 6.80_{anti} (s, H₈ chromene), 6.82_{syn} (s, H₈ chromene), 6.87_{syn} (d, $J = 8.5$ Hz, H₆ chromene), 6.90_{anti} (d, $J = 8.5$ Hz, H₆ chromene), 6.93_{syn} (d, $J = 8.0$ Hz, H_{3,5} phenyl-2-oxoethoxy), 7.05_{anti} (d, $J = 8.0$ Hz, H_{3,5} phenyl-2-oxoethoxy), 7.07_{anti} (t, $J = 7.5$ Hz, H₆ benzothiazine), 7.12_{syn} (t, $J = 7.5$ Hz, H₆ benzothiazine), 7.33_{syn} (d, $J = 8.5$, H₅ chromene), 7.44_{anti} (d, $J = 8.5$, H₅ chromene), 7.52_{anti} (d, $J = 8.5$, H_{3,5} benzoyl), 7.63_{syn} (d, $J = 8.0$ Hz, H_{3,5} benzoyl), 7.64-7.67_{anti} (m, H_{5,8} benzothiazine), 7.68_{syn} (m, H_{5,8} benzothiazine), 7.72_{syn} (s, H₄ chromene), 7.82-7.85_{anti} (m, H_{2,6} benzoyl and H_{2,6} phenyl-2-oxoethoxy), 7.85-7.89_{syn} (m, H_{2,6} benzoyl and H_{2,6} phenyl-2-oxoethoxy), 8.06_{anti} (s, H₄ chromene); ¹³C NMR (*syn*-isomer, 125 MHz, CDCl₃) δ 24.5, 37.6, 57.4, 70.6, 101.1, 111.8, 112.9, 116.9, 118.7, 119.3, 126.7, 128.3, 128.4, 129.5, 129.6, 129.9, 130.0, 131.9, 132.0, 132.3, 132.8, 135.5, 139.6, 140.9, 154.7, 160.6, 161.3,

191.6, 192.4; Anal. calcd for C₃₃H₂₃Br₂NO₅S: C, 56.19; H, 3.29; N, 1.99. Found: C, 56.21; H, 4.31; N, 2.09.

3-(3-Methyl-2-(4-methylbenzoyl)-3,4-dihydro-2H-benzo[b][1,4]thiazin-3-yl)-7-(2-oxo-2-p-tolyloethoxy)-2H-chromen-2-one (2t)

Yellow solid (397 mg, 70%); as mixture of diastereomers (*anti/syn*: 30/70); IR (KBr, cm⁻¹) 3397 (NH), 1702 (C=O); ¹H NMR (500 MHz, CDCl₃) δ 1.74_{syn} (s, CH₃ benzothiazine), 1.85_{anti} (s, CH₃ benzothiazine), 2.36_{anti} (s, CH₃ phenyl-2-oxoethoxy), 2.42_{syn} (s, CH₃ phenyl-2-oxoethoxy), 2.44_{anti} (s, CH₃ benzoyl), 2.45_{syn} (s, CH₃ benzoyl), 4.50_{syn} (s, NH), 4.61_{anti} (s, NH), 5.31_{anti} (s, O-CH₂), 5.35_{syn} (s, O-CH₂), 5.61_{anti} (s, C-H benzothiazine), 5.96_{syn} (s, C-H benzothiazine), 6.69_{anti} (d, $J = 2.1$ Hz, H₈ chromene), 6.72_{syn} (t, $J = 7.3$ Hz, H₇ benzothiazine), 6.75_{anti} (t, $J = 7.3$ Hz, H₇ benzothiazine), 6.81_{syn} (d, $J = 2.1$ Hz, H₈ chromene), 6.87_{syn} (dd, $J = 8.0$ and 2.1 Hz, H₆ chromene), 6.90-6.93_{syn/anti} (m, H_{5,8} benzothiazine), 7.05_{anti} (t, $J = 7.3$ Hz, H₆ benzothiazine), 7.11_{syn} (t, $J = 7.3$ Hz, H₆ benzothiazine), 7.17_{anti} (d, $J = 8.0$ Hz, H_{3,5} phenyl-2-oxoethoxy), 7.28-7.33 (m, H₅ chromene (*syn/anti*), H_{3,5} benzoyl (*syn/anti*) and H_{3,5} phenyl-2-oxoethoxy (*syn*)), 7.34_{anti} (d, $J = 8.0$ Hz, H_{2,6} phenyl-2-oxoethoxy), 7.73_{syn} (s, H₄ chromene), 7.79_{anti} (d, $J = 8.0$ Hz, H_{2,6} benzoyl), 7.87_{syn} (d, $J = 8.0$ Hz, H_{2,6} phenyl-2-oxoethoxy), 7.94_{syn} (d, $J = 8.0$ Hz, H_{2,6} benzoyl), 8.06_{anti} (s, H₄ chromene); ¹³C NMR (*syn*-isomer, 125 MHz, CDCl₃) δ 24.0, 27.7, 32.6, 43.3, 56.0, 80.2, 110.4, 110.6, 113.6, 116.1, 118.7, 118.9, 119.9, 122.4, 124.5, 126.2, 127.2, 128.4, 129.2, 131.6, 140.1, 154.2, 161.2, 161.4, 192.1, 192.6; MS, m/z (%) 575 (M⁺, 8%), 557 (64), 542 (43), 410 (35), 264 (44), 239 (29), 119 (100); Anal. calcd for C₃₅H₂₉NO₅S: C, 73.02; H, 5.08; N, 2.43. Found: C, 73.21; H, 5.12; N, 2.54.

3-(3-Methyl-2-(4-(methylsulfonyl)benzoyl)-3,4-dihydro-2H-benzo[b][1,4]thiazin-3-yl)-7-(2-(4-(methylsulfonyl)phenyl)-2-oxoethoxy)-2H-chromen-2-one (2u)

Yellow solid (576 mg, 82%); as mixture of diastereomers (*anti/syn*: 28/72); IR (KBr, cm⁻¹) 3394 (NH), 1688 (C=O), 1320 (SO₂), 1153 (SO₂); ¹H NMR (500 MHz, CDCl₃) δ 1.78_{syn} (s, CH₃ benzothiazine), 1.89_{anti} (s, CH₃ benzothiazine), 3.02_{anti} (s, SO₂-CH₃ phenyl-2-oxoethoxy), 3.04_{anti} (s, SO₂-CH₃ benzoyl), 3.09_{syn} (s, SO₂-CH₃ phenyl-2-oxoethoxy), 3.12_{syn} (s, SO₂-CH₃ benzoyl), 4.50_{anti} (s, NH), 4.69_{syn} (s, NH), 5.37_{syn} (s, O-CH₂), 5.41_{anti} (s, O-CH₂), 5.61_{anti} (s, C-H benzothiazine), 5.95_{syn} (s, C-H benzothiazine), 6.69_{anti} (s, H₈ chromene), 6.74_{syn} (t, $J = 7.5$, H₇ benzothiazine), 6.80_{anti} (t, $J = 7.5$, H₇ benzothiazine), 6.83_{syn} (s, H₈ chromene), 6.88-6.95_{syn} (m, H₆ chromene and H_{5,8} benzothiazine), 7.08_{anti} (d, $J = 8.0$, H₆ chromene), 7.11_{anti} (t, $J = 7.5$, H₆ benzothiazine), 7.13_{syn} (t, $J = 7.5$ Hz, H₆ benzothiazine), 7.36_{syn} (d, $J = 8.0$, H₅ chromene), 7.50_{anti} (d, $J = 8.0$, H₅ chromene), 7.59_{anti} (d, $J = 7.5$ Hz, H₈

benzothiazine), 7.74_{syn} (s, H₄ chromene), 7.82_{anti} (d, *J* = 7.5 Hz, H₅ benzothiazine), 7.94_{anti} (s, H₄ chromene), 8.04-8.19_{syn/anti} (m, H_{2,3,5,6} phenyl-2-oxoethoxy and H_{2,3,5,6} benzoyl); ¹³C NMR (*syn*-isomer, 125 MHz, CDCl₃) δ 24.5, 44.2, 44.3, 57.5, 65.5, 70.9, 101.0, 110.9, 113.0, 116.9, 119.4, 126.9, 127.4, 127.9, 128.1, 128.5, 129.1, 129.3, 129.8, 138.0, 139.6, 140.8, 141.0, 143.9, 145.1, 154.7, 160.4, 161.2, 190.6, 192.6; Anal. calcd for C₃₅H₂₉NO₅S₃: C, 59.73 ; H, 4.15; N, 1.99. Found: C, 59.59; H, 4.31; N, 2.30.

Pharmacology

Animals

Male NMRI mice weighing 20–30 g were used for studying in vivo antinociceptive activities of target compounds. Animals were maintained under standard conditions (24 ± 2°C, 60–70% humidity) and allowed food and water *ad libitum*. They were housed in appropriate cages with 12 h light/dark cycle. Before each experiment animals randomly

selected and allocated into groups. The whole protocol was approved by the Ethics Committee of the Faculty of Pharmacy at Tehran University of Medical Sciences.

Formalin-induced pain test

All target compounds **2a–u** were subjected for testing their analgesic activity using formalin paw test [25]. The compounds or standard drug mefenamic acid were administered i.p. (30 mg/kg, 0.2 mL/20 g body weight) as a suspension in saline and tween 80 (4% w/v). Each group of mice (n = 6 animals per group) were pretreated by test compounds, mefenamic acid or vehicle, 30 minutes before injection of formalin (20 µL, 0.5%, s.c.) into the planar surface of the right hind paw. The amount of time that the animal spent licking injected paw was measured during the first 10 minutes (phase 1, neurogenic) and 10–30 minutes (phase 2, inflammatory) after formalin injection.

Table 1 Antinociception activity of target compounds 2a–u and mefenamic acid (30 mg/kg, i.p.) assessed by formalin test in mice

Compounds	Phase 1			Phase 2		
	Licking time ^a	Inhibition ^b (%)	Relative activity ^c	Licking time	Inhibition (%)	Relative activity
2a	58 ± 3.46	48.10**	0.54	37.33 ± 3.93	44.28**	0.52
2b	51.33 ± 2.96	54.06***	0.61	50.33 ± 4.91	24.88	0.29
2c	68.33 ± 4.05	38.85**	0.44	38 ± 1.73	43.28**	0.51
2d	55 ± 2.74	50.78***	0.57	57.33 ± 6.35	14.43	0.17
2e	44 ± 2.89	60.63***	0.68	54 ± 4.93	19.40	0.23
2f	60.33 ± 3.76	46.01**	0.52	38.33 ± 4.63	42.79**	0.50
2g	56.66 ± 8.74	49.29**	0.55	55.66 ± 3.92	16.92	0.20
2h	70.25 ± 2.95	37.14**	0.42	38 ± 1	43.28**	0.51
2i	51.33 ± 2.40	54.06***	0.61	37 ± 1.15	44.78**	0.53
2j	46.25 ± 2.56	58.61***	0.66	50.33 ± 0.33	24.88	0.29
2k	51.66 ± 2.18	53.77***	0.60	51.66 ± 5.54	22.89	0.27
2l	70 ± 11.13	37.36**	0.42	37 ± 4.35	44.78**	0.53
2m	63.33 ± 8.21	43.33**	0.49	18.33 ± 0.33	72.64***	0.85
2n	69 ± 9.16	38.26**	0.43	40.66 ± 1.20	39.30**	0.46
2o	53.8 ± 3.21	51.85**	0.58	65 ± 6.41	2.98	0.03
2p	53.9 ± 3.18	51.76**	0.58	64.8 ± 4.19	3.28	0.04
2q	61.33 ± 5.78	45.12**	0.51	49.5 ± 2.02	26.12	0.31
2r	94.4 ± 4.89	25.86**	0.29	11 ± 1.7	93.64***	1.1
2s	50 ± 3.22	33.33**	0.37	5.2 ± 2.78	96.99***	1.14
2t	46 ± 2.4	38.86**	0.43	3 ± 1.04	98.26***	1.15
2u	34.8 ± 2.65	53.6***	0.61	14.8 ± 1.92	91.44***	1.07
Control	111.75 ± 6.94	-	-	67 ± 3.14	-	-
Mefenamic acid	12.33 ± 3.93	88.96***	1	10 ± 2.52	85.07***	1

^aData are expressed as mean ± S.E.M (number of animals in each group, n = 6).

^bThe percentage inhibition was determined by using the following formula: Inhibition % = 100 × (control – experiment)/control. The asterisks denote the levels of significance in comparison with control groups (**P* < 0.05, ***P* < 0.01 and ****P* < 0.001).

^cActivity relative to mefenamic acid was determined by using the following formula: Relative Activity = Inhibition % of compound/Inhibition % of mefenamic acid.

Acetic acid-induced writhing test

The analgesic activity was also determined *in vivo* by the abdominal constriction test induced by acetic acid (0.6%; 0.1 mL/10 g) in mice [21]. An acetic acid solution was administered i.p. 30 minutes after administration of compounds or mefenamic acid. After the treatment, pairs of mice were placed in separate boxes and the numbers of constrictions of the abdominal muscles, together with stretching, were counted cumulatively over a period of 60 minutes. Antinociceptive activity was expressed as the percentage of inhibition of constrictions when compared with the vehicle control group.

Statistical analysis

The nociception data are expressed as means \pm SEM. Variance analysis (ANOVA) followed by Bonferroni's test was used to compare means. *P*-values less than 0.05 were considered to be statistically significant.

Results and discussion

Chemistry

The dihydrobenzothiazole derivatives **1** were quantitatively obtained by reaction of 3-acetylcoumarins with 2-aminothiophenol derivatives in the presence of acetic acid under reflux condition or microwave irradiation [19,20]. The intramolecular Mannich-type reaction of compounds **1** with different phenacyl halides in the presence of KF/Al₂O₃ and catalyzing by quinine hydrochloride in ethanol afforded 3,4-dihydro-2*H*-benzothiazine derivatives **2a-r** via a ring expansion. When 7-hydroxy-3-(benzothiazol-2-

yl) coumarin derivative **1e** was treated with 2.5 equivalents of phenacyl halides, without protection of hydroxyl group, *O*-phenacyl derivatives **2s-u** was obtained in excellent yields (Scheme 1). The physicochemical and spectral data of new compounds **2e**, **2j-o**, and **2s-u** are described in experimental section.

Biological activity

Formalin-induced nociception test

All target compounds **2a-u** were tested using formalin-induced pain test in mice [25]. The obtained results were reported as mean \pm SEM of licking time and as percent of inhibition in Table 1. In general, the results showed that most of compounds were significantly able to reduce the licking time with percent of inhibition in the range of 25% to 60% at the first phase. The standard drug mefenamic acid showed 89% reduction of the licking time during the first phase. Amongst the tested compounds, **2a**, **2c**, **2f**, **2h**, **2i**, **2l-n** and **2r-u** significantly reduced the formalin induced licking time in the range of 39-98% as compared to mefenamic acid with 85% of inhibition during the second phase. Compounds **2m** and **2r-u** showed more effective antinociceptive activity in the second phase rather than first phase, indicating their ability to inhibit nociception associated with inflammatory response. Indeed, 7-hydroxy- and 7-phenacyloxy-coumarin derivatives (**2r** and **2s-u**, respectively) were more effective than mefenamic acid. Compounds **2s** and **2t** were the most effective compounds at the dose of 30 mg/kg.

Table 2 Antinociception activity of selected compounds in comparison with mefenamic acid (30 mg/kg, i.p.) assessed by acetic acid-induced writhing test in mice

Compound	Nociception (Mean \pm SEM)	Inhibition (%) ^a	Relative activity ^b
2b	0.6 \pm 0.24***	99	1.4
2c	38 \pm 4.04***	49	0.7
2d	9.6 \pm 2.54***	87	1.3
2g	3.5 \pm 1.09***	96	1.37
2h	3 \pm 1.84***	97	1.38
2i	4.6 \pm 2***	94	1.34
2k	20 \pm 2.48***	73	1.04
2o	6 \pm 3.2***	92	1.31
2r	29 \pm 2.12***	63	0.9
2s	14 \pm 2.28***	80	1.14
2t	30 \pm 7.6***	60	0.85
2u	2 \pm 1.3***	98	1.4
Control ^c	75 \pm 3.2		
Mefenamic acid	23 \pm 1.3***	70	

^aThe percentage inhibition was determined by using the following formula: Inhibition% = 100 \times (control - experiment)/control.

^bActivity relative to mefenamic acid was determined by using the following formula: Relative Activity = Inhibition % of compound/Inhibition % of mefenamic acid.

^cTween 80 in saline (4% w/v).

****P* < 0.001 vs. control.

Acetic acid-induced writhing test

The analgesic activity of compounds **2b-d**, **2g-i**, **2k**, **2o** and **2r-s** was also evaluated in vivo by using abdominal constriction test induced by acetic acid in mice [21]. The abdominal constriction response induced by acetic acid is sensitive procedure to establish efficacy of peripherally acting analgesics. The analgesic activity was expressed as the percentage of inhibition of constrictions when compared with the control group. The results are summarized in Table 2.

Significant protection against writhing was observed in animals treated with all test compounds where the mean numbers of writhes after 1 h were less than 38 compared to 75 in the control group. The percent of inhibition was in the range of 49-99%. All tested compounds were more effective than standard drug mefenamic acid with the exception of **2c**, **2r** and **2t**. Compounds **2b** and **2u** with percent of inhibition $\geq 98\%$ were the most effective compounds in acetic acid-induced writhing test. Moreover, compounds **2g-i** and **2o** exhibited high protection against writhing (percent of inhibition $> 90\%$).

Structure-activity relationships

From the structure-activity relationships of unsubstituted coumarin series (compound **2a-m**) based on the late stage of formalin-induced test, it was inferred that 3-thienylcarbonyl group is more favorable for activity. By comparing the activity of 7-substituted coumarin compounds **2r-u** with those of other compounds it is appeared that the 7-hydroxy or 7-phenacyloxy groups dramatically increase the effectiveness of compounds and their ability to inhibit nociception associated with inflammatory response. On contrary, compounds **2r-u** showed low level of inhibition at early phase of formalin test.

By comparing the percent of inhibition of 4-(methylsulfonyl)benzoyl derivatives **2d**, **2r** and **2u**, it is revealed that the introduction of hydroxyl group on 7-position of coumarin ring diminished the antinociception activity, while the introduction of 4-(methylsulfonyl)phenacyloxy- group increased the activity as resulted from writhing test. In the 7-phenacyloxy-coumarin derivatives **2s-u**, methylsulfonyl substituent was more favorable than bromo and methyl groups. The observed results of unsubstituted coumarin derivatives in Table 2 demonstrate that electron donating or bulky groups (for example, methoxy or phenyl, respectively) can increase antinociceptive activity in writhing test.

Conclusion

In summary, a series of 3-(3-methyl-3,4-dihydro-2H-benzo[*b*][1,4]thiazin-3-yl)-2H-chromen-2-one derivatives **2a-u** bearing different aroyl group on the 2-position of benzothiazine ring were described as potential analgesic agents. The antinociceptive properties of target compounds were determined by formalin-induced test and

acetic acid-induced writhing test in mice. The effect of substituent on aroyl moiety was explored by introduction of various electron withdrawing, electron donating or bulky groups. Surprisingly, compound **2u** bearing 2-[4-(methylsulfonyl)benzoyl]- moiety on benzothiazine ring and 4-(methylsulfonyl)phenacyloxy- group on the 7 position of coumarin nucleus showed better profile of antinociception in both models. It was more effective than mefenamic acid during the late phase of formalin-induced test as well as in the acetic acid-induced writhing test. However, unsubstituted coumarin derivative **2b** containing 4-methylbenzoyl moiety on benzothiazine ring, fully protected animals against writhing and was moderately able to inhibit the both phases of the formalin test. Considering the significant antinociceptive action of phenacyloxy-coumarin derivatives, compound **2u** prototype might be further used as model to obtain new more potent analgesic drugs.

Additional file

Additional file 1: Table S1. Chemical structure of coumarin compounds **2a-u**.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MA: Synthesis of target compounds. MK: Synthesis of target compounds. SE: Collaboration in identifying of the structures of target compounds, manuscript preparation. SF: Collaboration in determination of antinociceptive properties. SFG: Collaboration in determination of antinociceptive properties. MA: Supervision of the pharmacological part, AF: Collaboration in identifying of the structures of target compounds. AS: Design of target compounds and supervision of the synthetic and pharmacological parts. All authors read and approved the final manuscript.

Acknowledgments

This work was financially supported by grants from Research Council of Tehran University of Medical Sciences and INSF (Iran National Science Foundation).

Author details

¹School of Chemistry, University College of Science, University of Tehran, P.O. Box 14155-6455, Tehran, Iran. ²Department of Medicinal Chemistry, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medical Sciences, Tehran 14176, Iran. ³Department of Medicinal Chemistry and Pharmaceutical Sciences Research Center, Faculty of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran. ⁴Department of Toxicology and Pharmacology, Pharmaceutical Sciences Research Center, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran 14176, Iran.

Received: 19 May 2013 Accepted: 28 August 2013

Published: 7 January 2014

References

1. Ruoff G, Lema M: Strategies in pain management: new and potential indications for COX-2 specific inhibitors. *J Pain Symptom Manage* 2003, **25**:S21-S31.
2. Giovannoni MP, Cesari N, Graziano A, Vergelli C, Biancalani C, Biagini P, Dal Piaz V: Synthesis of pyrrolo[2,3-*d*]pyridazinones as potent, subtype selective PDE4 inhibitors. *J Enzyme Inhib Med Chem* 2007, **22**:309-318.

3. Cesari N, Biancalani C, Vergelli C, Dal Piaz V, Graziano A, Biagini P, Ghelardini C, Galeotti N, Giovannoni MP: **Arylpiperazinylalkylpyridazinones and analogues as potent and orally active antinociceptive agents: synthesis and studies on mechanism of action.** *J Med Chem* 2006, **49**:7826–7835.
4. Magiatis P, Melliou E, Skaltsounis AL, Mitaku S, Léonce S, Renard P, Pierré A, Atassi G: **Synthesis and cytotoxic activity of pyranocoumarins of the seselin and xanthyletin series.** *J Nat Prod* 1998, **61**:982–986.
5. Beillerot A, Dominguez JCR, Kirsch G, Bagrel D: **Synthesis and protective effects of coumarin derivatives against oxidative stress induced by doxorubicin.** *Bioorg Med Chem Lett* 2008, **18**:1102–1105.
6. Zhou X, Wang XB, Wang T, Kong LY: **Design, synthesis, and acetylcholinesterase inhibitory activity of novel coumarin analogues.** *Bioorg Med Chem* 2008, **16**:8011–8021.
7. Sashidhara KV, Kumar A, Kumar M, Sarkar J, Sinha S: **Synthesis and in vitro evaluation of novel coumarin–chalcone hybrids as potential anticancer agents.** *Bioorg Med Chem Lett* 2010, **20**:7205–7211.
8. Sashidhara KV, Kumar A, Kumar M, Srivastava A, Puri A: **Synthesis and antihyperlipidemic activity of novel coumarin bisindole derivatives.** *Bioorg Med Chem Lett* 2010, **20**:6504–6507.
9. Lee S, Sivakumar K, Shin WS, Xie F, Wang Q: **Synthesis and anti-angiogenesis activity of coumarin derivatives.** *Bioorg Med Chem Lett* 2006, **16**:4596–4599.
10. Leal LKAM, Ferreira AAG, Bezerra GA, Matos FJA, Viana GSB: **Antinociceptive, anti-inflammatory and bronchodilator activities of Brazilian medicinal plants containing coumarin: a comparative study.** *J Ethnopharmacol* 2000, **70**:151–159.
11. Keri RS, Hosamani KM, Shingalapuri RV, Hugar MH: **Analgesic, anti-pyretic and DNA cleavage studies of novel pyrimidine derivatives of coumarin moiety.** *Eur J Med Chem* 2010, **45**:2597–2605.
12. Kalkhambkar RG, Kulkarni GM, Kamanavalli CM, Premkumar N, Asdaq SMB, Sun CM: **Synthesis and biological activities of some new fluorinated coumarins and 1-aza coumarins.** *Eur J Med Chem* 2008, **43**:2178–2188.
13. Ghate M, Kusanur RA, Kulkarni MV: **Synthesis and in vivo analgesic and anti-inflammatory activity of some bi heterocyclic coumarin derivatives.** *Eur J Med Chem* 2005, **40**:882–887.
14. Bolakatti GS, Maddi VS, Mamledesai SN, Ronad PM, Palkar MB, Swamy S: **Synthesis and evaluation of anti-inflammatory and analgesic activities of a novel series of coumarin mannich bases.** *Arzneim-Forsch/Drug Res* 2008, **58**:515–520.
15. Khode S, Maddi V, Aragade P, Palkar M, Ronad PK, Mamledesai S, Thippeswamy AHM, Satyanarayana D: **Synthesis and pharmacological evaluation of a novel series of 5-(substituted)aryl-3-(3-coumarinyl)-1-phenyl-2-pyrazolines as novel anti-inflammatory and analgesic agents.** *Eur J Med Chem* 2009, **44**:1682–1688.
16. Melagraki G, Afantitis A, Igglessi-Markopoulou O, Detsi A, Koufaki M, Kontogiorgis C, Hadjipavlou-Litina DJ: **Synthesis and evaluation of the antioxidant and anti-inflammatory activity of novel coumarin-3-aminoamides and their alpha-lipoic acid adducts.** *Eur J Med Chem* 2009, **44**:3020–3026.
17. Rathore BS, Kumar M: **Synthesis of 7-chloro-5-trifluoromethyl-7-fluoro/7-trifluoromethyl-4H-1,4-benzothiazines as antimicrobial agents.** *Bioorg Med Chem* 2006, **14**:5678–5682.
18. Trapani G, Reho A, Morlacchi F, Latrofa A, Marchini P, Venturi F, Cantalamessa F: **Synthesis and antiinflammatory activity of various 1,4-benzothiazine derivatives.** *Farmaco Sci* 1985, **40**:369–376.
19. Khoobi M, Emami S, Dehghan G, Foroumadi A, Ramazani A, Shafiee A: **Synthesis and free radical scavenging activity of coumarin derivatives containing a 2-methylbenzothiazoline motif.** *Arch Pharm* 2011, **344**:588–594.
20. Khoobi M, Ramazani A, Foroumadi A, Hamadi H, Hojjati Z, Shafiee A: **Efficient microwave-assisted synthesis of 3-benzothiazolo and 3-benzothiazolino coumarin derivatives catalyzed by heteropoly acids.** *J Iran Chem Soc* 2011, **8**:1036–1042.
21. Collier HDJ, Dinnin LC, Johnson CA, Schneider C: **The abdominal constriction response and its suppression by analgesic drugs in the mouse.** *Br J Pharmacol Chemother* 1968, **32**:295–310.
22. Tjolsen A, Berge OG, Hunskaar S, Rosland JH, Hole K: **The formalin test: an evaluation of the method.** *Pain* 1992, **51**:5–17.
23. Khoobi M, Ramazani A, Foroumadi A, Emami S, Jafarpour F, Mahyari A, Slepokura K, Lis T, Shafiee A: **Highly cis-diastereoselective synthesis of coumarin-based 2,3-disubstituted dihydrobenzothiazines by organocatalysis.** *Helv Chim Acta* 2012, **95**:660–671.
24. Victoria FN, Radatz CS, Sachini M, Jacob RG, Perin G, da Silva WP, Lenard EJ: **KF/Al₂O₃ and PEG-400 as a recyclable medium for the selective α-selenation of aldehydes and ketones. Preparation of potential antimicrobial agents.** *Tetrahedron Lett* 2009, **50**:6761–6763.
25. Hunskaar S, Hole K: **The formalin test in mice: dissociation between inflammatory and non-inflammatory pain.** *Pain* 1987, **30**:103–114.

doi:10.1186/2008-2231-22-9

Cite this article as: Alipour et al.: Antinociceptive properties of new coumarin derivatives bearing substituted 3,4-dihydro-2H-benzothiazines. *DARU Journal of Pharmaceutical Sciences* 2014 **22**:9.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

 **BioMed Central**
www.SID.ir