

A Study on Knowledge, Attitude and Practice of Secondary School Girls in Qazvin on Iron Deficiency Anemia

***D Shojaeizadeh**

*Dept. of Public Health Administration, School of Public Health and Institute of Public Health Research,
Tehran University of Medical Sciences, P.O.Box 14155-6446, Tehran, Iran.*

Key Words: Knowledge, attitude, practice, iron deficiency anemia

ABSTRACT

A research was carried out to determine the factors affecting knowledge, attitude and practice of secondary school girls concerning Iron deficiency anemia, in Qazvin city. The study population consists of 218 Students who were randomly selected from ten secondary schools in a selected area. A questionnaire was prepared to collect the data and statistical tests of χ^2 and regression were employed to analyse the data. The results of the study indicated that 57.3 percent of students had poor knowledge, 54.1 percent unfavorable attitude and 44.5 percent weak practice on iron deficiency anemia. The results also showed that the field of education, the level of education, age, fathers' job and mothers' job had significant relation with knowledge. There is significant relationship between knowledge and attitudes. It was found that knowledge had significant relationship with practice.

INTRODUCTION

Iron deficiency anemia is a serious problem for the health of individuals. It has an effect on mental and physical health of the people.

It reduced the physical ability of children and youngsters (3,10). World wide prevalence rate of anemia among pregnant women and preschool children is 40 to 60 percent (13). In developing countries the prevalence rate is higher (44%) whereas it is lower in developed countries (12%) (16).

The results of many researches showed that the prevalence of anemia in West Africa is 31%, in China 26.61%, in South Asia 46.7% and in South America 20.9% (1). According to WHO data, half a million pregnant women die because of abortion, severe anemia and infection (14). Anemia not only has influence in physical development but has an effect on mental development too (2). Some studies showed that nutritional patterns have influence in anemia (8).

The results of the study showed that 10-50% of Iranian pregnant women have iron deficiency anemia (6). Many studies showed that there is significant relationship between incomes, Menarche age, social and economical situations, jobs, education of parents and iron deficiency anemia (7,12).

The results of a research which was carried out by the Ministry of Health in Iran, demonstrated that the prevalence rate of Iron deficiency and anemia among women aged 15 to 49 is 34% and 33.34%, respectively (11). According to the FAO, iron deficiency anemia is one of the most important factors affecting and causing malnutrition in Iran (4). Therefore, it is essential to find out what factors contribute to iron deficiency anemia.

In Iran, more than 20 percent of women aged 15-45 suffer from iron deficiency anemia (15). With respect to this issue it is very important to run health education programmes for women in order to prevent iron deficiency anemia. It is also important to conduct health education programs for women before marriage. Health education in schools plays a vital role in increasing knowledge, of the students. The objective of this

research is to study the knowledge, attitude and practice of the students on iron deficiency anemia and factors affecting their KAP.

MATERIALS AND METHODS

This is a cross-sectional study. The study population consists of 5059 secondary school girls in Qazvin city. The preparatory study showed that the knowledge of study population is 50 percent, with respect to this issue, it was planned to take a sample of 250 students with confidence of 95% 218, students for study.

The method of sampling is stratified and the stratified sampling was all secondary schools in Qazvin city. There is 10 secondary girl schools in Qazvin city and from each school, 25 students randomly selected for study.

A questionnaire was developed to gather data. The questionnaire consists of 53 questions: 11 questions on demographic characteristics, 25 questions on knowledge, 11 questions on attitude and 6 questions on practice. Chi-square (χ^2) and Regression have been used to analyse the data.

RESULTS

The results of the study indicated that 57.3 percent of students had poor knowledge on iron deficiency anemia, 54.1 percent unfavorable attitude towards iron deficiency anemia and 44.5 percent did not perform appropriate behaviour to prevent iron deficiency anemia (Table 1).

The results of the study showed that the knowledge of students who were 15-17 years old is higher than those who were less than 15 and more than 17 years old. There was statistically significant relationship between knowledge and the age of student (Table 2).

The findings of the study indicated that the knowledge of students whose field of education was natural sciences, was higher than those whose field of education was non-natural

*Corresponding author, Tel: + 98 -21-8951319; Fax: + 98-21-6462267; E-mail: shojaes@yahoo.com

sciences. There was statistically significant relationship between knowledge and the field of education of students (Table 3).

The results also indicated that the knowledge of those who their mothers were employed was higher than those who their mothers were. There is significant relationship between mothers' job and students' knowledge on iron deficiency anemia (Table 4).

The results of the study showed that students whose level of education was high (year 3-4), had more knowledge about Iron deficiency anemia in comparison with other students. There was significant relationship between level of education and knowledge (Table 5).

The findings showed that students who had poor knowledge on iron deficiency anemia, had unfavorable attitude, the students who had good knowledge had favorable attitude. There was statistically significant relationship between knowledge and attitude (Table 6).

The results of the study indicated that students who had poor knowledge on iron deficiency anemia, had weak practice and those who had good knowledge had appropriate behaviour. There was significant relationship between knowledge and practice (Table 7).

No significant relationship was found between students' attitude and independent variables of the study. The findings showed that the more mothers were illiterate the less students performed preventive behaviour on iron deficiency anemia.

DISCUSSION

The results of the study showed that there was a need for education of individuals concerning iron deficiency anemia. The findings indicated that there were poor knowledge, unfavorable attitude and poor practice about the iron deficiency anemia, among the students. It was found that the main source of information for the students were family members and school teachers. The teachers and family members training will likely increase students' knowledge, influence their attitudes and motivate their practice. Health educators, therefore, should be involved in education of parents and teachers.

The results indicated that students with inappropriate beliefs and habits were more likely to have unfavorable attitude. The students who were in need of education on iron deficiency anemia, are ranked as follows:

1. The students whose educational subject was experimental.
2. The students of first and second grade in secondary school.
3. The students whose mothers are house wives or unemployed.
4. The students whose fathers were workers or unemployed.

Table 1. Relationship between KAP and mothers' Job

KAP Score		Mothers' Job						
		Housewife		Employed		Total		Test
		N0	%	N0	%	N0	%	
Knowledge	0-50	116	60.4	9	34.6	125	57.3	X ² = 6.23
	50-75	66	43.3	15	57.6	81	37.2	df=1
	75 +	10	5.3	2	7.8	12	5.5	P= 0.0125
	Total	192	100	26	100	218	100	
Attitude	0-50	106	55.2	12	46.2	118	54.1	
	50-75	80	41.8	14	53.8	94	43.1	X ² = 0.76
	75 +	6	3.3	0	0	6	2.8	df=1
	Total	192	100	26	100	218	100	P=0.3
Practice	0-50	87	45.3	10	38.5	97	44.5	
	50-75	87	45.3	16	45.3	103	47.2	X ² = 0.4
	75 +	18	9.4	0	0	18	18.3	df=1
	Total	192	100	26	100	218	100	P=0.5

Score : 0 – 50 : Poor
50 – 70 : Intermediate
70 + : good

Table 2. Relationship between knowledge and age of students

Knowledge	Age								Test
	Less then 15		15 – 17		17+		Total		
	No.	%	No.	%	No.	%	No.	%	
0 – 50	39	70.9	79	51.6	7	70	125	57.3	X ² = 5.54 df=1 P=0.02
50 – 75	11	20	67	43.8	3	30	81	37.2	
75 +	5	9.1	7	4.6	0	0	12	5.5	
Total	55	100	153	100	10	100	218	100	

Table 3. Relationship between knowledge and field of education of students

Knowledge	Field of education							
	Natural		Non – natural		Not clear		Total	
	Sciences		Sciences					
	No.	%	No.	%	No.	%	No.	%
0 – 50	25	39.6	41	54.4	59	72.8	125	57.3
50 – 75	34	54	30	40.5	17	20.9	81	57.3
75 +	4	6.4	3	4.1	5	6.3	12	5.5
Total	63	100	74	100	81	100	218	100

X² = 16.1 ; df=2 ; P=0.0001

Table 4. Relationship between knowledge and mothers' job

Knowledge	Mothers' Job						Test
	Housewife		Employed		Total		
	N0	%	N0	%	N0	%	
0-50	116	60.4	9	34.6	125	57.3	X ² = 6.23 df=1 P= 0.0125
50 -75	66	43.3	15	57.6	81	37.2	
75 +	10	5.3	2	7.8	12	5.5	
Total	192	100	26	100	218	100	

Table 5. Relationship between knowledge and students' level of education

Knowledge	Level of education						Test
	Year		Year		Total		
	1-2		3- 4				
	No.	%	No.	%	No.	%	
0 – 50	89	64.4	36	42.8	125	57.3	X ² = 11.72 df=1 P=0.001
50 – 75	38	28.3	43	51.2	81	31.2	
75 +	7	5.3	5	6	12	5.5	
Total	13	100	89	100	218	100	

Table 6. Relationship between knowledge and attitude toward iron deficiency anemia

Knowledge	Attitude						Test
	0 – 50		50 +		Total		
	No.	%	No.	%	No.	%	
0 – 50 *	86	72.8	39	39	125	57.3	X ² = 25.4
50 +	32	27.2	61	61	93	42.7	df=1
Total	118	100	100	100	218	100	P = 0.001

0 – 50 : Poor knowledge ; 50 + : Knowledge

Table 7. Relationship between knowledge and practice on iron deficiency anemia

Knowledge	Practice						Test
	0 – 50		50 +		Total		
	No.	%	No.	%	No.	%	
0 – 50	49	50.5	76	62.8	125	57.3	X ² = 3.3
50 +	48	49.5	45	37.2	193	42.7	df=1
Total	97	100	121	100	218	100	P = 0.05

REFERENCES

1. ACC/SCN Second Report on the World Nutritional Situation (1993): Methods and Statistics, Vol. II, Chap.2, PP: 116, March.
2. Bararom BA, Kovitz RH and Meytes D (1992): Iron State in Female Adolescents, *Am J Dis Child*, **146**(83): 3.
3. Enwonwu G (1993): Functional Significance of Iron Deficiency, annual Nutrition Workshop series, Volume II, Center for Nutrition Meharry Medical College.
4. ESN-Nutrition Country Profile (1991): Iran, FAO, PP: 9.
5. Fairbanks VF (1994): Iron in Medicine, and Nutrition, Modern Nutrition in Health and Disease, Shils, ME Olson, J A and Shine, M. & ed. Lea and Febige, London, PP: 196.
6. Food and Nutrition Situation in the Islamic Republic of Iran (1991): ICN Reparatory Meeting, 18-21 August.
7. Hagshenas M, Mahloutgi RJ and Mohammadi N: Iron deficiency Anemia in Iranian Population Associated with high intake of Iron, *Am J Clin Nutr*, **25**: 143-6.
8. Jumesje J (1989): Preventing Iron Deficiency in Preschool Children by Implementing Educational and Screening Programme in an Inner City Practice, *BMJ*, 209.
9. Karani, Chanad, Combatting Anemia in Adolescent Girls (1994): Report from India, Mothersa Children.
10. Major Issues for Nutrition Strategies (1992): International Conferences on Nutrition, PP: 21.
11. Multi-Centre Study on Iron Deficiency Anemia 15 to 40 Years old (1995): Ministry of Health and Medical Education and Unicef. Women in the Islamic Republic of Iran Characteristics of for Selected Village and Tribal Communication Fars, Iran, PP: 12-16.
12. Petrosina A, Shayan K, Bash K and B (1969): Report of the Health
13. Preventing Specific Micronutrient Deficiency and Related' Wajessup (1992): International conference on Nutrition, PP: 21
14. Smyke P (1991): Women and Health, 2nd Book Company, London, PP: 61.
15. WHO/Geneva (1989): The Reproductive Health of Adolescent girls, PP: 5-15.
16. WHO (1992): Women's health. A Cross Age and frontier, WHO, Geneva, PP: 1-6.
17. WHO (1994): Women's Health: Age and Frontier, WHO Geneva.