

## Seroepidemiological Survey of Brucellosis Among Animal Farmers of Yazd Province

\*MH Salari

Dept. of Pathobiology, School of Public Health and Institute of Public Health Research, Tehran University of Medical Sciences, P.O.Box 14155-6446, Tehran, Iran.

**Key Words :** Seroepidemiology, brucellosis, animal farmers, Yazd province

### ABSTRACT

Brucellosis has remained a great problem of health in most of countries, which have failed in control of zoonosis infections. This disease is caused by species of *Brucella* and usually is transferred from animals to humans. The purpose of this study was to investigate the seroprevalence of brucellosis among animal farmers of Yazd province. In this descriptive study, 933 animal farmers were investigated by serological tests and blood culture method. Then the data were analyzed by SPSS statistical program. The results showed that the frequency distribution of seropositive cases by MAT and STAT were 35 (3.2%); 25 (2.7%) males and 10 (1.1%) females. The highest and lowest incidences of seropositive cases were among age group of 21-30 (1%) and more than 60 (0.3%) years old, respectively. Of the 35 seropositive cases, 2MET positive were 5 (0.5%), while all blood cultures were negative.

### INTRODUCTION

Brucellosis is a major cause of Zoonoses and an important public health problem in many parts of the world (27). The incidence of the disease has decreased markedly in industrialized countries (9,15). However, it remains a major public health problem in many developing countries (1,24).

Diagnosis of human brucellosis relies on blood culture and serological tests, such as the standard tube agglutination test (STAT), Coombs' test, and enzyme-linked immuno – sorbent assay (ELISA) (6,7).

Transmission of brucellosis can be the result of ingestion, direct contact via skin abrasion and mucous membranes (including the conjunctiva), and inhalation (in abattoirs and laboratories). Risk factors for infection include the handling of infected animals, ingestion of contaminated animal products such as unpasteurized milk and milk products (including cow, goat, and camel milk), meat, and handling of cultures of *Brucella* spp. in laboratories. This has significantly decreased the risk of this infection in individuals, who historically have been at increased risk, including veterinarians, farmers, meat inspectors and abattoir workers (8,10,20).

The purpose of this study was to investigate the seroprevalence of brucellosis among animal farmers of Yazd province.

### MATERIALS AND METHODS

Yazd province, central Iran, covers an area of about 74214 square kilometers and has an estimated population of about 800000 with almost 75% living in urban areas.

In this descriptive study 933 peripheral blood samples were collected from animal farmers of Yazd province from 1997 to 1999. For isolation and identification of *Brucella* species, biphasic blood culture medium (Hemolin, Biomerieux, France) which was incubated in an atmosphere of 5% to 10% carbon

dioxide for 30 days has been used (16). Serum specimens were analyzed in three phases, using suspension of *B. abortus* and *B. melitensis* (Wellcome Laboratories, UK). In the first phase, all specimens were screened by the microplate agglutination test (MAT). A titer of 1:80 or greater was considered to represent the presence of specific agglutination *Brucella* antibodies (seropositive). In the second phase, seropositive specimens were analyzed by the standard tube agglutination test (STAT). A titer of 1:160 or greater was taken as an index of seropositivity (6,7). In the third phase, *Brucella* antibody of animal farmers were investigated by 2-mercapthoethanol test (2 MET) (13,26).

The collected data and the results of laboratory tests were analyzed by statistical package for social science (SPSS), to determine those variable that were significantly associated with seropositivity to *Brucella*.

### RESULTS

Table 1 shows some demographic characteristic of animal farmers who provided blood samples for this study. Distribution of 35 seropositive cases by MAT and STAT were 25 (2.7%) males and 10 (1.1%) females. The highest and lowest incidence of brucellosis were observed among age group of 21-30 (1%) and more than 60 (0.3%) years old, respectively (Table 2 and Fig1). Statistical analysis showed significant differences between sexes ( $\chi^2 = 9.32$ ,  $P < 0.001$ ). There were no significant differences between age groups ( $\chi^2 = 3.22$ ,  $P < 0.5$ ). findings show that 5 (0.5%) of seropositive cases were 2 MET positive, and all blood cultures were negative.

The most common presenting clinical symptoms with active brucellosis were back pain, headache, fever, chill, night sweats weakness, myalgia, arthralgia and bone pain (Table 3).

\* Corresponding author, Tel:+98-352622-6557; Fax:+98-352622-6559; E-mail: msalari2000@yahoo.com

Table 1. Distribution of animal farmers on the basis of age and sex

Age ( year)	Sex				Total	
	Male		Female			
	( n = 811 )		( n = 122 )		( n = 933 )	
	No	%	No	%	No	%
< 10	4	( 0.4 )	( 0 )	0	4	( 0.4 )
10 – 20	194	( 20.8 )	( 1.9 )	18	212	( 22.7 )
21 – 30	142	( 15.2 )	( 1.2 )	11	153	( 16.4 )
31 – 40	117	( 12.5 )	( 2.7 )	25	142	( 15.2 )
41 – 50	116	( 12.4 )	( 4.2 )	39	155	( 16.6 )
51 – 60	123	( 13.2 )	( 2.5 )	23	146	( 15.6 )
> 60	115	( 12.3 )	( 0.6 )	6	121	( 13 )


Table 2. The results of serological tests and blood culture method

Tests	Serological tests								Blood culture	
	MAT		STAT				2 MET		Positive	Negative
	Positive	Negative	Positive		Negative		Positive	Negative		
Antibody titers			1/160	1/320	1/640	1/1280	≤1/80	≥1/160	<1/160	
Number	35	898	17	9	4	5	2	5	30	35
Percentage	3.8	96.2	1.8	1	0.4	0.5	0.2	0.5	3.2	3.8

Table 3. Distribution of seropositive animal farmers on the basis of clinical signs and symptoms

Clinical Signs and symptoms	Back pain	Headache	Fever	Chill	Night sweats	Weakness	Myalgia	Arthralgia	Bone pain
Number	3	14	27	21	23	18	11	17	13
Percentage	0.3	1.5	2.9	2.3	2.5	1.9	1.2	1.8	1.4

Fig.1. Distribution of seropositive animal farmers on the basis of age



## DISCUSSION

Brucellosis is diagnosed either by isolation of *Brucella* organism in culture, or by a combination of serological tests and clinical findings consistent with brucellosis. Isolation of the *Brucella* organism is the definitive means of diagnosis, but in practice it is difficult due to the early tissue localization and the exacting culture requirements of the organism. In practice, blood cultures are positive in 10-30% of brucellosis and the remainder is diagnosed serologically (28). Although no single test provides 100% sensitivity and specificity, STAT still remains the test of choice in diagnosis. In the presence of appropriate signs and symptoms, a presumptive diagnosis of brucellosis is usually defined serologically as a standard tube agglutination titer of 1:160 or greater (7).

Recent serological data on brucellosis in developed countries are not available. Indeed, brucellosis has been brought under control in these countries through rigorous diagnostic and control procedures at the animal production level, as well as through elimination of *Brucella* in livestock and proper pasteurization of milk (2). Therefore, there are very few reports of indigenously acquired human cases of brucellosis, while acute imported human infections continue to occur, often associated with the consumption of raw cheese or milk (5,25). Data from developing countries in the Mediterranean basin, particularly the Middle East, report seroprevalence rates ranging from 8% in Jordan (12) to 12% in Lebanon and Kuwait (4,17). Even higher seroprevalence rates have been reported in sub-Saharan countries, with percentages of 18% in Uganda (21) and 13% in Nigeria (24). According to the our findings most of patients had a history of infected cheese and milk consumption or direct contact with domestic animal. However, rate of seropositive cases among Yazd animal farmers population was very lower than above mentioned reports; 3.8% or 4.3 cases per 100000. The prevalence rate of brucellosis increases with age, a result that is consistent with observations made in Iran, Jordan, Lebanon, and Kuwait (4,11,17,18,19,22,29). In contrast to other studies, no significant difference was found in prevalence rate between age groups.

In an analysis of 104 cases of brucellosis in Saudi Arabia, 1288 cases in United States, and 60 cases in Iran, the most common symptoms and physical findings were reported fever, chills, weakness, malaise, sweating, backache, headache, lymphadenopathy, splenomegaly and arthritis (12,18,23). These results are nearly the same as our findings.

Brucellosis in animals remains a major public health hazard due to its transmissibility to man (2). The only effective way to control of the disease in man is by elimination of the infected animals and vaccination of healthy ones in order to reduce the risk of those in regular contact with animals, and to produce brucellosis free animal products (3).

The effectiveness of vaccination programs can be evaluated by investigating incidence rate in humans, especially people at high risk, before and after vaccination. It is recommended that surveillance of brucellosis be strengthened. Cooperation and joint supervision between the Ministries of Health and Veterinary Organization, and cooperation with neighboring countries, should also be encouraged. Furthermore, health education programs should be adopted that aim at stopping the spread of infection among animals and then to humans, and also adoption of hygienic measures among high-risk population.

## ACKNOWLEDGEMENT

It is with sincere appreciation that we acknowledge the following people who assisted the authors in this work: Dr. Abbas Mirshafie, from School of Public Health and Institute of Public Health Research, Tehran University of Medical Sciences; Dr. Mohammad Reza Sharefi from Shahid Sadoughi University of Medical Sciences; Gholam Reza Hassanpour, Ali Asghar Sadrabadi, Mohammad Mehdi Hakimian, Reza Beick, Ahmad Ali Hanafi-Bojd and Fatemah Falah, Yazd Training and Health Research Center; Dr. Yadolah Amrolahi and Dr. Mahmoud Bahri, Yazd Veterinary Organization.

## REFERENCES

1. AL-Freih HM, AL-Mohaya SA, Al Mohsen MF, Ibrahim EM, Al-Sohaibani M and et al. (1986): Brucellosis in Saudi Arabia: diverse manifestations of an important cause of pyrexial illness. *Ann Saudi Med*, **6**: 95-7.
2. Al-Khalaf SA, Mohammed BT and Nicoletti P (1992): Control of brucellosis in Kuwait. *Trop Anim Health Prod*, **24**: 45-9.
3. Al-Mezaini S, Sonoussi Y and Chang S (1984): A study on brucellosis among farm animals in rural Saudi Arabia Processing of the seventh conference on Biological Aspect of Saudi Arabia. *Saudi Biological Society*. PP: 3517.
4. Araj GF and Azzam RA (1996): Seroprevalence of antibodies among persons in high-risk occupation in Lebanon. *Epidemiol Infect*, **117**: 281-8.
5. Arnov PM, Smaron M and Ormiste V (1984): Brucellosis of a group of travelers to Spain. *JAMA*, **251**: 505-7.
6. Bettelheim KA, Maskill WJ, Metcalfe RV and Pearce JL (1984): The use of micro-agglutination technique to determine the antibody status of healthy New Zealanders to *Brucella abortus*. *J Hygiene*, **92**: 401-10.
7. Bettelheim KA, Maskill WJ and Pearce JL (1983): Comparison of standard tube and microagglutination technique for determining *Brucella* antibodies. *J Hygiene*, **90**: 33-9.
8. Buchanan TM, Faber LC and Feldman RA (1974): Brucellosis in the United states, an abattoir associated disease, Part 1, **53**: 403-13.
9. Chomel BB, De Bass EE, Manginamele DM, Rely KF, Faver TB, Sun RK and et al. (1994): Changing tenders in the epidemiology of human brucellosis in california from 1973 to 1992. *J Infect Dis*, **170**: 1216-30.
10. Corbel MJ (1997): Brucellosis: an overview. *Emerging infectious diseases*, **3**(2): 213-21.
11. Dajani YH, Masoud AA and Barakat HF (1989): Epidemiology and diagnosis of human brucellosis in Jordan. *J Trop Med Hyg*, **92**: 209-14.
12. Fox MD and Kaufman AF (1977): Brucellosis in the united states, 1965-1976. *J Infect Dis*, **153**: 122-5.
13. Landry ML, Rosa NR, De Macario EC, Folds JD, Lane HC and Nakamhira RM (1997): Manual of Clinical Laboratory Immunology. 5<sup>th</sup> edition. Washington DC: ASM Press, PP: 608-17.
14. Lulu AR, Arja GF, Khateeb MY, Yusuf AR, Ferech FF (1998): Human brucellosis in Kuwait. *Q J Med*, **66**: 39-44.
15. Madkour M and Gargani G (1985): Epidemiology aspects. In: Madkour M, editor. *Brucellosis*. London: Butterworths. PP: 11-28.
16. Mahon CR and Manuselis G (2000): Textbook of diagnostic Microbiology. 2<sup>nd</sup> edition. WB Saunders Company, PP: 1070-2.
17. Makarem E, Karjoo R and Omid A (1982): Frequency of *Brucella melitensis* in southern Iran. *J Trop Pediatr*, **28**: 97-100.
18. Malik GM (1997): A clinical study of brucellosis in adults in the Asir region of southern Saudi Arabia. *Am J Trop Med Hyg*, **56**(4): 375-7.
19. Mousa A, Elhag K, Khogali M and Marie A (1998): The nature of human brucellosis in Kuwait. *Rev infect Dis*, **10**: 211-7.

20. Murray PR, Baron EJ, Pfaller MA and et al. ( 1999 ): *Manual of clinical Microbiology*. 7<sup>th</sup> edition. Washington DC: ASM Press. PP: 625-31.
21. Ndyabahinduka DG and Chu IH (1984): Brucellosis in Uganda, *Int J Zoonoses*. **11**: 59-64.
22. Sabbaghian H and Nadim A (1974): Epidemiology of human brucellosis in Isfahan, *Iran J Hyg Camb*, **73**: 221.
23. Samar G, Aminia N and Zowghi E (1996): A survey on human brucellosis in Iran (Diagnosis and treatment of 60 cases). *Iran J Inf Trop Med*, **1**(3):31-6.
24. Six W, Rosegger H, Schneeweiss H and Withalam H( 1987 ): Serological investigation in Nigeria for anthroponozoonosis in human sera. *J Hyg Epidemiol Microbiol Immunol*, **31**: 493-5.
25. Taylor JP and Perdue JN (1989): The changing epidemiology of human brucellosis in Texas, 1977 – 1986, *Am Med J Epidemiol*, **130**:160-5.
26. Thomas MB and Luke CF (1980): 2 -mercaptoethanol *Brucella* agglutination test usefulness for predictiny Recovery from brucellosis. *J Clin Microbiol*, **11**: 6.
27. Volk WA, Gebhardt BM, Hammerskjold ML and Kadner RJ (1996): *Brucella Yersinia, Francisella and Pasteurella*. In: *Essentials of Medical Microbiology*. 5<sup>th</sup> edition. Philadelphia: Lippincott – Raven. PP:383-95.
28. Young EJ (1995): An overview of human brucellosis. *Clin Infect Dis*, **21**:278.
29. Zowghi E and Ehadi A (1986): A survey on human brucellosis "Malt Fever" in Iran (serological and bacteriological investigations). *Arch Inst Razi*, **36**(37): 69-73.

Archive of SID