

Frequencies of Mutations in the Connexin 26 Gene (GJB2) in Two Populations of Iran (Tehran and Tabriz)

***M Hashemzadeh Chaleshtori^{1,2}, L Hoghooghi Rad³, M Dolati³, R Sasanfar⁴, A Hoseinipour⁴
M Montazer Zohour¹, H Pourjafari⁵, A Tolooi⁴, M Ghadami⁴, DD Farhud¹, MA Patton⁶**

¹*Dept. of Human Genetics, School of Public Health, Tehran University of Medical Sciences, Iran*

²*Dept. of Biochemistry and Genetics, Medical School, Shahrekord University of Medical Sciences, Iran*

³*Dept. of Biology, School of Basic Sciences, Science and Research Campus, Islamic Azad University, Tehran, Iran*

⁴*Dept. of Exceptional Children, Ministry of Education and Training of the Islamic Republic of Iran*

⁵*Genetic Division, Medical School, Hamadan University of Medical Sciences, Iran*

⁶*Medical Genetics Unit, St George's Hospital Medical School, University of London, UK*

Abstract

While hearing loss has been considered to be a very heterogeneous disorder, mutations in Gap junction beta 2 (GJB2) gene encoding Connexin 26 (Cx26) protein are the major cause of autosomal recessive and sporadic non-syndromic deafness in many populations. In this study, we have investigated the prevalence of the GJB2 gene mutations using nested PCR pre screening strategy and direct sequencing method. Two hundred and seventy two hearing impaired subjects were studied from 210 families obtained from two large cities of Iran (Tehran and Tabriz). Twenty four different genetic variants were identified. Cx26 mutations were found in 53 of the 210 families (25.2%) including T8M, 35delG, W24X, R32H, V37I, E47X, 167delT, delE120, Y136X, R143W, R184P, 235delC and V27I+E114G. Homozygosity and compound heterozygosity for the Cx26 mutations were found in 39 of 210 (18.5%) families. Homozygosity for the 35delG mutation was the most common that causes hearing loss in 28 (13.3%) patients. Six novel variants H16R, E101E, K102Q, G200R, 327delG and G130A were detected in this study. As a conclusion, the present survey revealed that the rate of mutation in Cx26 gene in our area is lower than in Europe; nevertheless, this rate is regarded as a considerable cause of deafness in the cited provinces in Iran.

Keywords: *Connexin 26, GJB2, Deafness, Autosomal recessive non syndromic hearing loss, Iran*

Introduction

Hearing loss is the most frequent sensory disorder that affects 1 in 1000 neonates with more than 60% hereditary form (1). About 80% of the hereditary cases are non-syndromic and the major mode of inheritance is autosomal recessive (2). Mutations in the Connexin 26 (GJB2) gene (GJB2; MIM# 121011) at the DFNB1 locus (DFNB1; MIM#220290) on chromosome 13q12 have been shown to be involved in the

development of autosomal recessive and sporadic non-syndromic deafness in many populations (3-8). Connexin 26 is a member of the family proteins encodes the gap junction protein that is expressed in a variety of tissues including cochlea. Gap junctions are being served as a major communication system allowing the rapid exchange of electrolytes, second messengers and metabolites between adjacent cells (4). It is also thought that this gap junction plays an

important role in auditory transduction, by recycling endolymphatic potassium ions (9). Although, a single mutation known as 35delG accounts for the majority of Cx26 mutations in some populations (10-12) but is less frequent or even absent in other ethnic groups. They care other common mutations such as 235delC in the Japanese and Korean (13-15), 167delT in the Ashkenazi Jews (16) and R143W mutation in an African village (17).

In a previous study, 43 autosomal recessive non-syndromic hearing loss subjects from 34 families in Iran were analysed to identify GJB2 gene mutations and 11 different genetic variants were observed. GJB2-related deafness mutations were found in 9 of 34 ARNSHL families (26.5%) and homozygosity in 7 (20.6%) (18). In the present study we have identified the spectrum and prevalence of Connexin 26 mutations using nested PCR pre screening strategy and direct sequencing technique in coding region of the gene.

Materials and Methods

Patients We have analysed 272 patients from 210 unrelated families, consisting of 139 families from Tehran and 71 families from Tabriz, diagnosed with non-syndromic and autosomal recessive hearing loss (ARNSHL). Their ages were between 2 and 35 years old (mean: 13.3), attending hearing-impaired schools in Tehran and Tabriz. Medical history and pedigree information were obtained by interviewing and filling questionnaires. Consanguinity was observed as 74.8% in both populations under study. There were no syndromic features and the hearing of all parents was normal. All patients had mild to profound sensorineural hearing loss.

DNA analysis DNA was extracted from 10 ml peripheral whole blood using standard salting out procedures. In the first step, all samples were screened for the 35delG mutation using nested PCR strategy. Using a nested PCR procedure, the one base pair deletion of 35delG

was screened in 210 families. The entire coding sequence of Cx26 gene (Genbank accession# M86849) was amplified using primers CX26F5 5'CCTGTGTTGTGTGCATTCGTC3'/CX26R 5'5'CTCATCCCTCTCATGCTGTC3' (782 bp) at an annealing temperature of 59°C. The amplified product was then diluted and used as a template for a second round of PCR using primers CX26F4 5'CACGCTGCAGACGATCC3'/CX26R4 5'GGTGGAGTGTTTGTTCAC3'(43 bp) at an annealing temperature of 56° C. The amplified products were then separated by electrophoresis on a 15% polyacrylamide gel (40% 19:1 acrylamide: bisacrylamide) at 35 mA for 5:30 hours and the products were then visualised by silver staining. The 35delG mutations were detected by identification of two separated bands of 43 bp for the wild type and 42 bp for the mutant allele. Then, in order to confirm the results of this study and also for finding other mutations in the GJB2 gene, direct sequencing was used. One set of primers was used to produce the template for sequencing in coding (Genbank accession# U43932) regions. The primers CX26F5/ CX26R5 (as described above) were used to amplify the entire coding region of the gene. Sequencing was carried out in Sequence Laboratories (SeqLab), Gottingen, Germany, using ABI 3100 and ABI 3700 sequencer.

Results

In this study we analyzed the coding region of connexin 26 in 210 ARNSHL families from two large cities of Iran (139 families in Tehran and 71 families in Tabriz) using nested PCR and consequent direct sequencing. Thirteen Cx26 mutations including 35delG, T8M, 167delT, R143W, W24X, R32H, V37I, E47X, delE120, Y136X, R184P, 235delC and V27I+E114G were found in 53 of 210 (25.2%) families (Table 1). We identified 39 of 210 families (18.5%) with homozygous and compound heterozygous for Cx26 mutations, whereas, 14 other families (6.7%) revealed mutation only in

one allele. Twenty eight families (13.3%) were homozygous for the 35delG mutation in two cities. Also, 4 polymorphisms V27I, S86T, R127H and V153I and one unknown mutation M163V were found. S86T polymorphism was observed in all families (100%). Six new variants (H16R, E101E, K102Q, G130A, G200R and 327delG) were also detected (Fig. 1).

In Tehran, with a mixed population, 31(22.3%) families expressed 9 mutations among 139 families. Four novel variants including E101E, K102Q, G200R and 327delG were found in Tehran. Three variants E101E, K102Q and

327delG were heterozygous, which occurred in the second intracellular domain (IC2) of the Cx26 protein. The G200R, containing one homozygous change, occurred in the fourth transmembrane domain (TM4).

Twenty two (31%) families revealed 8 mutations among 71 deaf families from Tabriz. Also two new variants were found in this city including H16R and G130A. Both of these variants were heterozygous, that occurred in the first intracellular domain (IC1) and second intracellular domain (IC2) of the Cx26 protein, respectively.

Table 1: Genetic variation in the GJB2 gene identified in Iranian autosomal recessive non-syndromic hearing loss families

Genotype	Tehran Families n (%)	Tabriz Families n (%)	Total
T8M/V153I	2(1.4)	-	2
35delG/35delG	17(12.2)	11(15.5)	28
35delG/wt	2(1.4)	3(4.2)	5
35delG/167delT	1(0.7)	1(1.4)	2
35delG/R143W	1(0.7)	-	1
H16R/R143W	-	1(1.4)	1
W24X/wt	1(0.7)	-	1
V27I/wt	-	1(1.4)	1
V27I+E114G/wt	-	1(1.4)	1
R32H/R32H	1(0.7)	-	1
V37I/wt	-	1(1.4)	1
E47X/wt	1(0.7)	-	1
S86T	139(100)	71(100)	210
E101E/wt	1(0.7)	-	1
K102Q/wt	1(0.7)	-	1
delE120/delE120	-	1(1.4)	1
R127H/wt	-	2(2.8)	2
G130A/wt	-	1(1.4)	1
Y136X/Y136X	-	1(1.4)	1
R143W/R143W	1(0.7)	-	1
V153I/wt	5(3.6)	2(2.8)	7
M163V/wt	-	1(1.4)	1




Fig. 1: Nucleotide sequences of the novel variants have been identified in Cx26 gene (A:H16R, B:G130A, C:G200R, D:K102Q, E:E101E, F:327delG) in comparison with control group from Iran.

Discussion

Mutations in connexin 26 gene have been detected in many ethnic populations and are the most common cause of ARNSHL in many populations (4, 14). Mutations in Cx26 are responsible for 50% of autosomal recessive deafness. Thirty-five delG mutation is more frequently observed in Caucasian families from Tunisia, France, New Zealand and UK (10). This mutation is responsible for 10% of all childhood hearing loss and for 20% of all childhood hereditary hearing loss in American Caucasians originated from north and south Europeans (19). Our finding showed a lower rate of mutations in GJB2 (25.2%). This is also lower than those reported from Italy, Spain (49%) and Lebanon (33%) (11, 20). However, the high frequency of the 35delG mutation in white populations possibly is the results of a founder effect rather than a mutational hot spot (5,10,19).

Rate of 35delG mutation in Tehran (22.3%) was lower than that in Tabriz (31%) is the most common in both cities (58.5% of 53 families). Genetic variations in the cited locus may be due to migration during recent decades in Tehran and Tabriz. Since the rate of migration to Tehran is high, we consider it with a mixed population. Tabriz is alike Tehran, where migration occurs frequently.

The genetic variation, 327delG, was found in one of our patients, which observed in Gilan province (21). Upon reviewing this patient questionnaire, we noted that her mother was from Gilan. Therefore, such variation is probably originated from Gilaki nation. However more investigations are warranted to confirm such hypothesis.

In the previous study (18), using a method similar to the present survey (nested PCR pre screening strategy and direct sequencing method), in coding and non-coding regions of the gene, the GJB2-related deafness mutations were found in 7 from 34 families (20.6%) with homozygosity in this gene. In the present in-

vestigation we identified 39 from 210 families (18.5%) with homozygous and compound heterozygous mutations. The difference in observed mutation rate might be happened due to ignoring non-coding region analysis in this study.

Najmabadi and coworkers (22) studied 168 autosomal recessive nonsyndromic deafness from 83 families in Iran, using allele-specific PCR for screening the 35delG and pre screening method of single strand conformational polymorphism (SSCP) followed by sequencing to screen the other variants. GJB2-related deafness was diagnosed in 9 families (11%) with homozygous and compound heterozygous. Moreover, they found 9 different genetic variants. However, we detected 24 different genetic variants and found 39 (18.5%) of 210 families with homozygous and compound heterozygous for Cx26 mutations. This difference might be due to lower SSCP analysis efficiency and/or selected areas where the rate of mutation was lower than our.

In our survey, S86T polymorphism was observed in all cases. Similar observation was previously reported in Gilan and Khorasan provinces (18). Thus, it seems that such polymorphism exists in all populations of Iran.

In conclusion, the role of GJB2 gene mutations in autosomal recessive and sporadic non-syndromic hearing loss in Iran may be lower than that in other populations, studied so far (10, 11, 19). The rate of consanguinity in this study was relatively high (74.8%), compared to the rate of 37.3% consanguinity reported before (23). Such a high rate may be due to the consanguinity or ascertainment among our deaf population. Since populations in Iran are composed of several ethnic groups, more works on the ethnic populations are warranted.

Acknowledgments

We would like to thank the Ministry of Education and Training of the Islamic Republic of Iran, Department of Exceptional Children and

all the individuals, families, students and schools for hearing-impaired in Tehran and Tabriz for their contribution to this study. The Department of Exceptional Children, Zafar Avenue, Tehran, Iran, supported this study.

References

1. Marazita ML, Ploughman LM, Rawlings B, Remington E, Arnos KS, Nance WE (1993). Genetic epidemiology studies of early-onset deafness in the U.S. school-age population. *Am J Med Genet*, 46: 486-91.
2. Skvorak GAB, Morton CC (1999). Genetic causes of non-syndromic hearing loss. *Curr Opin Pediatr*, 11: 551-57. Review
3. Guilford P, Ben Arab S, Blanchard S, Le-villiers J, Weissenbach J, Belkahia A, et al (1994). A non-syndrome form of neurosensory, recessive deafness maps to the pericentromeric region of chromosome 13q. *Nat Genet*, 6:24-8.
4. Kelsell DP, Dunlop J, Stevens HP, Lench NJ, Liang JN, Parry G, et al (1997). Connexin 26 mutations in hereditary non-syndromic sensorineural deafness. *Nature*, 387:80-83.
5. Morell RJ, Kim HJ, Hood LJ, Goforth L, Frederici K, Fisher R, et al (1998). Mutations in the connexin 26 gene (GJB2) among Ashkenazi Jews with non-syndromic recessive deafness. *N Engl J Med*, 339: 1500-505.
6. Cohn ES, Kelley PM (1999). Clinical phenotype and mutations in connexin 26 (DFNB1/GJB2), the most common cause of childhood hearing loss. *Am J Med Genet*, 89: 130-36. Review
7. Gasparini P, Rabinonet R, Barbu-jani G, Melchionda S, Peterson M, Brondum-Nielsen K, et al (2000). High carrier frequency of the 35delG mutation in European populations. *Eur J Hum Genet*, 8: 19-23.
8. Van Camp G, Smith RJH (2003). Hereditary hearing loss homepage. Available from: www.uia.ac.be/dnalab/hhh
9. Holt JR and Corey DP (1999). Ion channel defects in hereditary hearing loss. *Neuron*, 22: 217-19.
10. Denoyelle F, Weil D, Maw MA, Wilcox SA, Lench NJ, et al (1997). Prelingual deafness: high prevalence of 30delG mutation in the Connexin 26 gene. *Human Mol Genet*, 6: 2173-77.
11. Estivill X, Fortina P, Surrey S, Rabionet R, Melchionda S, D'Agruma L, Mansfield E, et al (1998). Connexin-26 mutations in sporadic and inherited sensorineural deafness. *Lancet*, 351: 394-398.
12. Green GE, Scott DA, McDonald JM, Woodworth GG, Sheffield VC, et al (1999). Carrier rates in the Midwestern United States for GJB2 mutations causing inherited deafness. *JAMA*, 281: 2211-16.
13. Abe S, Usami S, Shinkawa H, Kelley PM, Kimberling WJ (2000). Prevalent connexin 26 gene (GJB2) mutations in Japanese. *J Med Genet*, 37:41-43.
14. Park HJ, Hahn SH, Chun YM, Park K, Kim HN (2000). Connexin 26 mutations associated with nonsyndromic hearing loss. *Laryngoscope*, 110: 1535-1538.
15. Kudo T, Ikeda K, Kure S, Matsubara Y, Oshima T, Watanabe KL, et al (2000). Novel mutations in the connexin 26 gene (GJB2) responsible for childhood deafness in the Japanese population. *Am J Med Genet*, 90: 141-145.
16. Sobe T, Ehrlich P, Berry A, Korostichevsky M, Vreugede S, Avraham KB, et al (1999). High frequency of the deafness-associated 167delT mutation in the connexin 26 (GJB2) gene in Israeli Ashkenazim. *Am J Med Genet*, 86: 499-500.
17. Brobby GW, Muller-Myhsok B, Horstmann RD (1998). Connexin 26 R143W mutation associated with recessive nonsyn-

- dromic sensorineural deafness in Africa. *N Engl J Med*, 338:548-49.
18. Hashemzadeh Chaleshtori M, Farhud DD, Taylor T, Hadavi V, Pattpn MA, Afzal AR (2002). Deafness-associated connexin 26 gene (GJB2) mutation in Iranian population. *Iranian J Publ Health*, 31(3-4): 75-9.
 19. Kelley PM, Harris DJ, Comer BC, Askew JW, Fowler T, Smith SD, et al (1998). Novel mutations in the connexin26 gene (GJB2) that cause autosomal recessive (DFNB1) hearing loss. *Am J Hum Genet*, 62:792-99.
 20. Mustapha M, Salem N, Delague V, Chouery E, Ghassibeh M, Rai M, Loiselet J, et al (2001). Autosomal recessive nonsyndromic hearing loss in the Lebanese population: revalence of the 30delG mutation and report of two novel mutations in the connexin 26 (GJB2) gene *J Med Genet*, 38: 36e ([electronic](#)).
 21. Hashemzadeh Chaleshtori M, Dowlati M, Farhud DD, Hoghooghi Rad L, Sasanfar R, Hoseinipour A, et al (2004). Two novel mutations and predominant 35 delG mutation in the connexin 26 gene (GJB2) in Iranian populatins. *Iranian J Publ Health*, 33:14-19.
 22. Najmabadi H, Cucci RA, Sahebjam S, Kouchakian N, Farhadi M, Kahrizi K, et al (2002). JGB2 mutations in Iranian with autosomal recessive non-syndromic sensorineural hearing loss. *Human Mut*, Mutation in Brief#504 online.
 23. Farhud DD, Kamali MS, Marzban M, Andonian L, Saffari R (1991). Consanguinity in Iran. *Iranian J Publ Health*, 20: 1-14.

Archive of SID