

Glucose-6-phosphate dehydrogenase (G6PD) Deficiency

DD Farhud¹, *L Yazdanpanah²

¹Genetic Clinic, Vallie Asr Sq, 16 Keshavarz Blvd. Tehran, Iran.

²Dept.of Nutrition, School of Public Health, Iran University of Medical Sciences, Tehran, Iran

(Received 28 Apr 2008; accepted 3 Dec 2008)

Abstract

Glucose-6-phosphate dehydrogenase (G6PD) Deficiency is the most prevalent enzymopathy in mankind. It has sex-linked inheritance. This enzyme exists in all cells. G6PD deficiency increases the sensitivity of red blood cells to oxidative damage. G6PD deficiency was discovered in 1950 when some people suffered hemolytic anemia as a result of taking antimalarial drugs (primaquin). Most people with G6PD deficiency do not have any symptoms, till they are exposed to certain medications, *Fava beans* and infections; and then their red blood cells are hemolyzed. The degree of hemolysis changes according to the degree of enzyme deficiency and the oxidant agent exposure. G6PD deficiency has many different variants and Mediterranean variant is the most common mutation in the world. G6PD deficiency is considered a health problem worldwide, especially in Asia, Middle East and Mediterranean countries. In this article, we have reviewed the importance and function of G6PD enzyme, incidence rate of G6PD deficiency in the world and Iran, genetic and variants of this enzyme, clinical manifestation, diagnosis and treatment of the enzyme deficiency.

Keywords: G6PD, Oxidative damage, Sex-linked inheritance, Hemolytic anemia, Mediterranean variant

Introduction

Glucose-6-phosphatase dehydrogenase (G-6-PD) deficiency is the most common enzymopathy in humans (1). G6PD deficiency was discovered for the first time when hemolytic anemia occurred in some persons who consumed anti-malarial drug named primaquine (1, 2). It is an X-linked disorder and is a highly polymorphic enzyme. This enzymopathy affects 400 million people worldwide (3). The enzyme gene is located on the long arm of the X- chromosome (Xq28) (1).

G6PD catalyzes the first step of pentose phosphate pathway, during this reaction NADPH is produced and protects red blood cells from oxidative damage (4-6).

Most deficient people do not show any symptoms until following exposure to oxidative drugs, some infections and ingestion of *Fava bean* (1). Mediterranean mutation is the most common variant of enzyme deficiency and often associated with favism (2, 3).

Avoiding of the oxidative agents that induces hemolytic anemia is the most important treatment

in the enzyme deficiency. Neonatal screening and health education can reduce the incidence rate of G6PD deficiency clinical manifestations (1). The goal of this article is to review the importance of G6PD enzyme in antioxidant defense enzyme system, function of the enzyme, prevalence of G6PD deficiency in the worldwide, structure and different mutations of this enzyme, The major clinical manifestations like acute hemolytic anemia induced by oxidative drugs and some infections, neonatal jaundice and chronic non-spherocytic hemolytic anemia, diagnosis and treatment of the enzyme deficiency.

Importance of G6PD Enzyme

Super oxide, hydrogen peroxide, hydroxyl radical is called ROS (Reactive Oxygen Species). They cause oxidative stress and damage cell's Membranes (7, 8).

Increased production of ROS or decreased antioxidant defense enzymes play a major role in oxidative injuries in different organs, tissues and cells including brain, heart, vascular cells (8) and causes brain diseases like Alzheimer and Parkin-

son diseases and also considered to contribute to the aging process (9-11).

Enzymes like superoxide dismutases, catalase, glutathione peroxidases, glutathione reductase and glucose-6-phosphate dehydrogenase are antioxidant defense enzymes. In body defense anti-oxidant system, G6PD is considered as an essential modulator enzyme that has a very important role in all cells especially in red blood cells (8).

To know the crucial role of G6PD, at first the role of other anti-oxidant enzymes must be understood. Superoxide dismutase converts two superoxide radicals into one hydrogen peroxide and one oxygen (8). Catalase converts hydrogen peroxide into water and oxygen (8, 12). Glutathione peroxidase is necessary for reducing hydrogen peroxide and lipid peroxides to water and lipid alcohols (8). Glutathione reductase: main role of this enzyme is regenerating reduced glutathione (GSH) from oxidized form (GSSG) (8, 12). G6PD is a key enzyme for maintenance of redox potential in cells. G6PD produces NADPH in pentose phosphate pathway. NADPH is important as a central reductant and regulates of redox potential (13). It also acts as a cofactor for other anti-oxidant enzymes like glutathione reductase (8, 12).

Reduced glutathione is required as a cofactor for the glutathione peroxidases and, thus reduced glutathione and glutathione cycles is crucial for neutralization of hydrogen peroxide and lipid peroxides and also protects protein sulfhydryl groups against oxidation (2).

Catalase is found in two forms, active and inactive, NADPH is critical for conversion of inactive form into active form (2).

Different studies point out the importance of G6PD enzyme in other cell functions like its control of cell death. G6PD is a principal enzyme in cell death and intracellular redox potential needs to be regulated to control cell death (13). As the production of ROS increases, so does cell death.

In another study, role of anti-oxidant enzyme and oxidative stress in Parkinson disease was surveyed. One hundred and fifteen patients with Parkinson and 37 healthy people were selected.

In these samples, activities of superoxide dismutase, Catalase, Glutathione peroxidase and G6PD in red blood cells were measured. They showed that activities of this enzyme in patients were significantly lower than normal person. These findings demonstrated that oxidative stress is the main cause of the start and progress of neurodegenerative in the patients and can also be related with intensity of the disease (14).

Function of G6PD

G6PD catalyzes the first step in the pentose phosphate pathway, converts Glucose-6-phosphate into 6-phosphogluconolactone and during this conversion, the important reductant metabolite named NADPH is provided. The pathway is the only source for producing of NADPH in red blood cells (2), because they lack mitochondria, nucleus and ribosomes and other pathways that produce NADPH(4). NADPH is necessary for generating of GSH from its oxidized form, GSSG, and subsequent maintenance of intracellular GSH pools. GSH maintains normal structure, elasticity and integrity of red blood cells, and sustain hemoglobin in ferrous state that is essential for carrying oxygen (4). G6PD and NADPH are key factors for protection of red blood cells from oxidative damage and peroxides (2). Peroxides are usually removed from red blood cells by glutathione peroxidase that uses reduced glutathione. Reduced glutathione reacts with harmful peroxides and neutralize them. In this reaction reduced glutathione is oxidized and glutathione reductase regenerates reduced glutathione by using of NADPH. NADPH is oxidized and G6PD is required for producing reduced NADPH again (1, 2).




Fig. 1: Function of G6PD (5)

Catalase is another anti-oxidant enzyme that is abundant in red blood cells and helps to the removal of peroxides from red blood cells through activation by NADPH (2). Unless peroxides are neutralized, they will cause oxidative injuries. Hemoglobin and red blood cell membrane molecules that contain SH groups are destroyed (1, 2).

Hemoglobin is denatured irreversibly, precipitates and forms Heinz bodies. Heinz bodies destroy membranes of red blood cells then leads to haemolysis and acute anemia.

Regarding to the roles of NADPH in redox state of cells, G6PD is the principal enzyme in chain reaction that is necessary for protecting all cells especially red blood cells against oxidant agents (2). Then G6PD-deficient cells especially red blood cells are susceptible to damage by reactive oxygen species and oxidative stress (2).

Malaria and G6PD deficiency

The geographical distribution of malaria is similar to the world distribution of deficient G6PD vari-

ants. It is postulated that the high frequency of G6PD deficiency has arisen because G6PD deficient variants confer some resistance against severe malaria caused by *Plasmodium falciparum* (4, 15, 16). The exact mechanism of this protection is still unknown. Red blood cells are the host cells for *Plasmodium falciparum*. *Plasmodium* parasites oxidize NADPH and diminish the level of reduced glutathione (GSH) in red blood cells. This effect in G6PD deficiency is more severe, leading to oxidative-induced damage to the RBC (4). Also *Plasmodium* parasites break down hemoglobin, and release toxic substances like Fe, which is a source of oxidative stress and it will cause hemolysis (17). Therefore the growth of *Plasmodium* parasites is decreased. In addition, damaged red blood cells are removed by phagocytosis at an early ring-stage of parasites' maturation, and that decreases the growth of parasites even more (1, 18).

Farhud et al. performed a study in the north of Iran. They measured serum proteins and immunoglobulins in favic patients and healthy controls, both school boys. The results showed higher amounts in healthy individuals and normal range in patients. The normal range in the patients is suggested to be due to positive selection that induces a developed immune response and produces better chances not to get affected by some other endemic infectious diseases such as malaria (19).

Genetics, structure

The gene for the G6PD enzyme is one of the important genes located on the telomeric region of the long arm of X chromosome (Xq28) (1, 5, 20), are more likely to affect males than females. Males are more likely than females to suffer all X-linked genetic conditions, such as G6PD deficiency (21).

Clinical and biochemical analyses have characterized more than 400 (2, 22) variants most emerging from point mutations in the coding region of gene (2, 23), base changes that result in amino acid substitution, displaying a field of phenotypes with different abnormal enzyme activity and wide ranging degree of clinical symptoms (23). Monomer of G6PD has 515 amino acids with a molecular weight 59 kDa (2, 24).

G6PD in the active form consists of the same subunits of either dimer or tetramer and includes tightly bound NADP (2, 24). It is supposed that Lys and Arg, amino acids 386 and 387 in en-

zyme, bind one of the phosphates in NADP (2). The shift between two active forms depends on PH. Aggregation of inactive monomers and conversion into an active form requires presence of NADPH (2, 24).

Variants of G6PD

According to the level of enzyme activity, World Health Organization classified variants of G6PD to five groups (15, 25).

Class 1: Severe deficiency of the enzyme with Chronic nonspherocytic hemolytic anemia.

Class 2: Severe deficiency of the enzyme, enzyme activity is less than 10% of normal.

Class 3: Moderate deficiency of the enzyme, enzyme activity is %10-60 of normal.

Class 4: Very mild to none deficiency of the enzyme, enzyme activity is %60-100 of normal.

Class 5: There is increased enzyme activity.

Mutations that are responsible for class 1 variants are confined to the NADP binding site of the enzyme or glucose-6-phosphate binding site (2, 24). They are located near the carboxy end of the enzyme (25). They form severe clinical symptoms because they are clustered at the position involved in the dimer formation of the active form of G6PD (26, 27) and influence stability and activity of enzyme (27). But mutations that cause mild clinical symptoms are located at the amino end of the molecule (25). G6PD variants that have been discussed in this article are shown in Table 1. G6PD variants that have been characterized at the DNA level are shown in Table 2.

Table 1: G6PD variants that have been discussed in this article (2)

Biochemical variants	Nucleotide substitution		Amino acid substitution		WHO class
A ⁻	202G	A	68Val	Met	3
	376A	G	126Asn	Asp	
Mediterranean	563C	T	188Ser	Phe	2
Mahidol	487G	A	163Gly	Ser	3
Viangchan	871G	A	291Val	Met	2
Cosenza	1376G	C	459Arg	Pro	2
Chatham	1003G	A	335Ala	Thr	3

Table 2: G6PD variants that have been characterized at the DNA level (2)

Variant	Nucleotide Substitution		WHO Class	Amino Acid Substitution	
Gaohe	95 A	G	2	32 His	Arg
Gaozhou					
Sunderland	105-107 del		1	35 Ile	del
Aures	143 T	C	2	48 Ile	Thr
Metaponto	172 G	A	3	58 Asp	Asn
A ⁻					
Distrito Federal					
Metera					
Castilla	202 G	A	3	68 Val	Met
Alabama	376 A	G		126 Asn	Asp
Betica					
Tepic					
Ferrara					
Ube	241 C	T	3	81 Arg	Cys
Konan					
Lagosanto	242 G	A	3	81 Arg	His
Vancouver	317 C	G	1	106 Ser	Cys
	544 C	T		182 Arg	Trp
	592 C	T		198 Arg	Cys
Sao Borga	337 G	A	4	113 Asp	Asn
A	376 A	G	4	126 Asn	Asp
Chinese- 4	392 G	T	?	131 Gly	Val
Ilesha	466 G	A	3	156 Glu	Lys
Mahidol	487 G	A	3	163 Gly	Ser
Plymouth	488 G	A	1	163 Gly	Asp
Chinese-3	493 A	G	2	165 Asn	Asp
Shinshu	527 A	G	1	176 Asp	Gly
Santamaria	542 A	T	2	181 Asp	Val
	376 A	G		126 Asn	Asp
Mediterranean					
Dallas					
Birmingham	563 C	T	2	188 Ser	Phe
Sassari					
Cagliari					
Panama					
Coimbra	592 C	T	2	198 Arg	Cys
Santiago	593 G	C	1	198 Arg	Pro
Sibari	634 A	G	3	212 Met	Val
Minnesota					
Marion	637 G	T	1	212 Val	Leu
Gastonia					
Harilaou	648 T	G	1	216 Phe	Leu
Mexico City	680 G	A	3	227 Arg	Gln
A-	680 G	T	3	227 Arg	Leu
	376 A	G		126 Asn	Asp
Stonybrook	724-729		1	242-243	
	GGC del			Gly & Thr	
Wayne	769 G	C	1	257 Arg	Gly
Cleveland	820 G	A	1	274 Glu	Lys
Chinese-1	835 A	T	2	279 Thr	Ser

Table 2: Continued...

Variant	Nucleotide Substitution		WHO Class	Amino Acid Substitution	
Seattle					
Lodi	844 G	C	2	282 Asp	His
Modena					
Lodi	844 G	C	2	282 Asp	His
Modena					
Montalbano	854 G	A	3	285 Arg	His
Viangchan	871 G	A	2	291 Val	Met
Jammu					
West Virginia	910 G	T	1	303 Val	Phe
Kalyam	949 G	A	3	317 Glu	Lys
Kerala					
A-	968 T	C	3	323 Leu	Pro
Betica	376 A	G		126 Asn	Asp
Seima					
Nara	953-976 del		1	319-326 del	
Chatham	1003 G	A	3	335 Ala	Thr
Fushan	1004 C	A	2	335 Ala	Asp
Chinese-5	1024 C	T	?	342 Leu	Phe
Irepetra	1057 C	T	2	353 Pro	Ser
Loma Linda	1089 C	A	1	363 Asn	Lys
Olomouc	1141 T	C	1	381 Phe	Leu
Tomah	1153 T	C	1	385 Cys	Arg
Iowa					
Walter Reed	1156 A	G	1	386 Lys	Glu
Iowa City					
Springfield					
Guadalajara	1159 C	T	1	387 Arg	Cys
Mt. Sinai	1159 C	T	1	387 Arg	Cys
	376 A	G		126 Asn	Asp
Beverly Hills					
Genova	1160 G	A	1	387 Arg	His
Worcester					
Praba	1166 A	G	1	389 Glu	Gly
Nashvill					
Anaheim	1178 G	A	1	393 Arg	His
Calgary					
Portici					
Alhambra	1180 G	C	1	394 Val	Leu
Puetro Limon	1192 G	A	1	398 Glu	Lys
Riverside	1228 G	T	1	410 Gly	Cys
Japan	1229 G	A	1	410 Gly	Asp
Shinagava					
Alhambra	1180 G	C	1	394 Val	Leu
Puetro Limon	1192 G	A	1	398 Glu	Lys
Riverside	1228 G	T	1	410 Gly	Cys
Japan	1229 G	A	1	410 Gly	Asp
Shinagava					
Alhambra	1180 G	C	1	394 Val	Leu
Puetro Limon	1192 G	A	1	398 Glu	Lys
Riverside	1228 G	T	1	410 Gly	Cys

Table 2: Continued...

Variant	Nucleotide Substitution		WHO Class	Amino Acid Substitution	
Alhambra	1180 G	C	1	394 Val	Leu
Puetro Limon	1192 G	A	1	398 Glu	Lys
Riverside	1228 G	T	1	410 Gly	Cys
Japan Shinagava	1229 G	A	1	410 Gly	Asp
Tokyo	1246 G	A	1	416 Glu	Lys
Georgio	1284 C	A	1	428 Tyr	End
Varnsdorf	3' interon 10 splice site del		1	N/A	
Pawnee	1316 G	C	2	439 Arg	Pro
Telti Kobe	1318 C	T	1	440 Leu	Phe
Santiago de Cuba	1339 G	A	1	447 Gly	Arg
Cassano	1347 G	C	2	449 Gln	His
Union Maewo	1360 C	T	2	454 Arg	Cys
Andalus	1361 G	A	1	454 Arg	His
Cosenza	1376 G	C	2	459 Arg	Pro
Taiwan-Hakka Gifu-like	1376 G	T	2	459 Arg	Leu
Kaiping Anant Dhon Petrich Sapporo	1388 G	A	2	463 Arg	His
Campinas	1463 G	T	1	488 Gly	Val

G6PD deficiency in the World

Based on the findings of WHO, 7.5% of the world populations have one or two genes for G6PD deficiency and 2.9% are G6PD deficient (28).

The best known G6PD-deficient variants that occur at a high frequency are G6PD A⁻ and the Mediterranean variants. Mediterranean mutations are identified by very low activity in red blood cells (29). Mediterranean variant is the most common variant in Southern Europe, Middle East, and India (2). Rate of incidence of Mediterranean mutation in Turkey is (77%), Iran (69%), India (60.4%), and also in Pakistan (30) and Saudi Arabia the Mediterranean mutation is the prevalent variant (31).

In Kuwaiti population, the most common variants are G6PD A⁻ and the Mediterranean variants and also a lower rate of Chatham variant and Aures

are detected in Kuwaiti population. Postulating that gene flow from the Indian Sub-continent, sub-Saharan Africa and other parts of Mediterranean is responsible for molecular heterogeneity of G6PD variants in the population (30, 32). The most G6PD variant in Italy, Sardinia and Greece is G6PD Mediterranean.

In Egypt and Libya, North of Africa, G6PD Mediterranean is the most common variant (30). G6PD A⁻ and G6PD Mediterranean have respectively the highest rate of incidence in Algeria and both of them are associated with Favism. Molecular heterogenous G6PD deficiency in Algeria suggests gene flow from Sub-Saharan Africa and other parts of Mediterranean (33).

G6PD A⁻ is found in Africa, Southern Europe and all areas African people were taken to (2). G6PD deficiency in Spain is heterogenous. The most

prevalent variant in Spain is G6PD A⁻. Two other important variants are Seattle and Union. The existence of G6PD Aures and Santamaria, that are polymorphic in Algeria, suggests significant gene migration from Africa to Europe through Spain (30, 34).

G6PD deficiency in Mexico is heterogenous and G6PD A⁻ is relatively prevalent. Prevalence of G6PD A⁻ and Seattle mutations in Mexico were likely introduced by African slaves and Spanish immigrants (35).

Viangchan, Mediterranean and Mahidol are common G6PD mutations in the Malays (36). Vi-

angchan, Mahidol and canton were the most prevalent G6PD variants in Thais. G6PD Viangchan in Laotians and non-Chinese Southeast Asian population is probably the most common variant (37).

G6PD deficiency in IRAN

The prevalence of G6PD deficiency in Iranian population is 10-14.9% reported by WHO (28). The most prevalent variant of G6PD deficiency in Iran is the Mediterranean (38-40), and the next most prevalent variant is Chatham, and Chatham incidence rate is 13-27% (39).

The incidence rates of G6PD mutations in different provinces in Iran are shown in Table 3.

Table 3: The Prevalence of G6PD mutations in provinces of Iran (38, 40)

	Khorasan	Mazandaran	Golestan	Hormozgan	Sistan and Balochestan	Gilan	kermanshah
Percent of Medit mutation	66	69	62.2	79.45	80.42	86.4	91.2
Percent of Chatham mutation	12	27	26.8	8.21	2.17	9.7	7.3

The incidence rate of G6PD deficiency in Tehran neonates was 2.1% (41). In a study in Kermanshah Province 5.3% of samples were severely deficient in G6PD enzyme. Polymorphic mutations in the region were Mediterranean, Chatham, Cosenza (40). There are higher prevalence of G6PD deficiency in northern and southeastern provinces of Iran (8.6- 16.4% in Northern provinces, 12% in southern part [Shiraz] and 19.3% in southeastern of Iran) (42, 43-45).

The three mutations found in the three northern provinces of Iran were Mediterranean, Chatham, Cosenza (42). Khalili et al reported the prevalence of G6PD deficiency in Gilan, a northern province of Iran 6.4% (46). In Khorasan, North-eastern province of Iran, 22% of samples did not display one of the known mutations in Iran. Different ethnic groups who are living in this province include Persian, Turkmen, Afghan, Turk, Kurd and Arab. The unknown variants of enzyme deficiency observed in the region may bear simi-

larities to those in Khorasan neighboring such as Afghanistan and Turkmenistan (30).

Clinical Manifestations of G6PD deficiency

Acute hemolytic anemia, neonatal jaundice and chronic non spherocytic hemolytic anemia are the major clinical manifestations associated with G6PD deficiency that discussed in this article:

1. Acute hemolytic anemia
 - a. induced by oxidative drugs
 - b. Favism
 - c. Infection-induced hemolysis
2. Chronic non spherocytic hemolytic anemia
3. Neonatal jaundice

1. Acute hemolytic anemia

Unless exposed to oxidative agents like oxidative drugs, infections and ingestion of fava beans, most patients with G6PD deficiency show no sign of acute hemolytic anemia (1, 4). Hemolytic anemia in G6PD A⁻ is self-limited because in this

mutation the younger red blood cells contain normal G6PD enzyme activity. But in Mediterranean G6PD with severe enzyme deficiency, hemolytic anemia is not self-limited. All red blood cells have enzyme deficiency and don't resist in hemolysis (2, 15, 47).

a. Induced by oxidative drugs

Each person's response to specific drugs is affected by genetic variations in an enzyme or enzyme system (48). Adverse drug reactions (ADRs) are a significant cause of morbidity and mortality. Acute hemolytic anemia induced by anti-malarial drugs is an example of adverse drug reactions that had been recognized for long time in G6PD deficient people. In these people, due to a defect in the gene coding for G6PD, the activity of the enzyme is reduced and that caused this adverse effect (49). Pharmacogenetics was introduced by Vogel in 1959 for study the relationship between genetic variations in genes involved in drug metabolism and drug response. More recently, the term pharmacogenomics has also been introduced, that is the broader application of genomic technologies and considers the genetic variants, patterns of gene expression and the way drugs influence gene function. It is postulated that pharmacogenomics is the study of entire genome, meanwhile pharmacogenetics is the study of a single gene. The term pharmacogenomics can be used to cover both Pharmacogenetics and Pharmacogenomics. Pharmacogenomics offers major potential benefits by improving drug response, reducing ADRs and understanding disease susceptibility (50).

Some drugs cause oxidative stress and induce hemolysis in G6PD-deficient red blood cells. They form hydrogen peroxide when they come in contact with hemoglobin (47).

During this reaction reduced glutathione is oxidized rapidly, glutathione pools are exhausted, hemoglobin is denatured and Heinz bodies are formed (2, 47). Red blood cells with enzyme deficiency are not capable of reducing NADP to NADPH for regenerating reduced glutathione from oxidized form that is necessary for inactivating peroxides and protecting cells against oxidative

injuries that cause hemolysis (47, 51). Drugs, substances and herbs can induce hemolytic anemia in G6PD deficient people are shown in tables 4, 5.

Primaquine

People with G6PD deficiency are about 20 to 30 times more sensitive to the hemolytic activity of primaquine than the people with normal G6PD (52). Mechanism for induction of hemolytic anemia was studied in rats (52, 53). Toxic metabolites of primaquin cause reduction of GSH in red blood cells (53), formation of methemoglobin (51) and Heinz bodies that induce hemolytic anemia (52, 53). The primaquine metabolite, 6-methoxy-8-hydroxylaminoquinoline after N-hydroxylation from 6-methoxy-8-aminoquinoline (52), by peroxidation of lipids in red blood cells with significant GSH and oxidation of protein in GSH-depleted red blood cells can induce hemolytic response (52).

Henna

Henna is a cosmetic dye that is used for dying hair, nails (54, 55) and also for treatment dermatitis (56). Some studies identified that henna can induce hemolytic anemia (55, 57, 58). Lawson (2-hydroxy-1,4-naphthoquinone) is a chemical substance in henna, and its structure and redox potential is similar to naphthalene metabolites that induce oxidative damage in red blood cells especially in G6PD-deficient person (55).

Tea and Polyphenols

In a study in China effects of extracts of black tea, green tea and decaffeinated green tea and their polyphenols on G6PD-deficient red blood cell in vitro were examined. The results showed that extracts of tea and their two polyphenol (epigallocatechin-3-gallate and epigallocatechin) changed oxidative condition in G6PD-deficient red blood cells in vitro. Level of GSH is lowered; methemoglobin, hemoglobin and GSSG are raised. The observed changes depend to doses of tea extracts and polyphenols that are used. These changes are not observed in normal red blood cells (59).

Table 4: Drugs and substances should be avoided by G6PD deficient individuals (1, 2, 60).

Drugs and substances (group)	Examples
Anti malarial drugs	Primaquin
Sulphonamides	Sulphacetamid, Sulphametoazole, Sulphanilamid, Sulphapyridin
Sulphones	Thiazolesulfone, Dapsone
Other sulphur-containing drugs	Glibenclamide
Nitrofurans	Nitrofurantoin(Furadantin)
Other drugs	Toluidin blue, Trinitrotoluene(TNT), Urate oxidase, Phenylhydrazine, Furazolidone (Furoxone), Methylene Blue, Nalidixic acid, Niridazole, Phenazopyridine, Isobutyl Nitrite, Acetanilide, Aspirin
Cosmetic substance	Henna
Other substances	Naphthalene, Moth balls, High-dose of vitamin K or ascorbic acid
Food substance	Fava beans

Table 5: Drugs and Substances (with their molecular formulation) should be avoided by G6PD deficient individuals according to the G6PD Deficiency Association (61)

Name	Molecular Formula	Risk Levels	Population at risk
Acetanilide (acetanilid)	C ₈ H ₉ N O	High	Medit., Asian
Acetylphenylhydrazine (2-Phynylacetohydrazide)	C ₈ H ₁₀ N ₂ O	High	All
Aldesulfone sodium (sulfoxone)	C ₁₄ H ₁₄ N ₂ Na ₂ O ₆ S ₃	High	All
Aminophenazone (aminopyrine)	C ₁₃ H ₁₇ N ₃ O	Low	All
Antazoline (antistine)	C ₁₇ H ₁₉ N ₃	Low	All
Arsine	As-H ₃	High	All
Ascorbic Acid	C ₆ H ₈ O ₆	Low	All
Beta-Naphthol (2-Naphthol)	C ₁₀ H ₈ O	High	All
Chloramphenicol	C ₁₁ H ₁₂ C ₁₂ N ₂ O ₅	High	Medit., Asian
Chloroquine	C ₁₈ H ₂₆ Cl N ₃	High	Medit., Asian
Ciprofloxacin	C ₁₇ H ₁₈ F N ₃ O ₃	High	Medit., Asian
Colchicine	C ₂₂ H ₂₅ N O ₆	Low	All
Dapsone (diaphenylsulfone)	C ₁₂ H ₁₂ N ₂ O ₂ S	High	All
Dimercaprol	C ₃ H ₈ O S ₂	High	All
Diphenhydramine (difenilhydramine)	C ₁₇ H ₂₁ N O	Low	All
Dopamine (L-dopa)	C ₈ H ₁₁ N O ₂	Low	All
Doxorubicin	C ₂₇ H ₂₉ N O ₁₁	High	Medit., Asian
Furazolidone	C ₈ H ₇ N ₃ O ₅	High	All
Glibenclamide	C ₃₂ H ₂₈ Cl N ₃ O ₅ S	High	Medit., Asian
Glucosulfone (glucosulfone sodium)	C ₂₄ H ₃₄ N ₂ Na ₂ O ₁₈ S ₃	High	All
Isobutyl Nitrite	C ₄ H ₉ N O ₂	High	Medit., Asian
Isoniazid	C ₆ H ₇ N ₃ O	Low	All
Menadiol Sodium Sulfate (Vitamin k4 sodium sulfate)	C ₁₁ H ₈ Na ₂ O ₈ S ₂	High	All
Menadione (menaphtone)	C ₁₁ H ₈ O ₂	High	All
Menadione sodium Bisulfite (Vitamin K3 sodium bisulfite)	C ₁₁ H ₈ O ₂ NaHSO ₃	High	All
Mepacrine (Quinacrine)	C ₂₃ H ₃ O Cl N ₃ O	High	Medit., Asian
Mesalazine-5-Aminosalicylic Acid(paraminosalicylic acid)	C ₇ H ₇ N O ₃	High	Medit., Asian
Methyltinionium Chloride (methylene blue)	C ₁₆ H ₁₈ Cl N ₃ S	High	All
Nalidixic Acid	C ₁₂ H ₁₂ N ₂ O ₃	High	Medit., Asian
Naphtalene, Pure (naphtalin)	C ₁₀ H ₈	High	All
Niridazole	C ₆ H ₆ N ₄ O ₃ S	High	All
Nitrofural (nitrofurazone)	C ₆ H ₆ N ₄ O ₄	High	All
Nitrofurantoin	C ₈ H ₆ N ₄ O ₅	High	All
Norfloxacine	C ₁₆ H ₁₈ F N ₃ O ₃	Low	All
O-Acetylsalicylic Acid (acetylsalicylic acid)	C ₉ H ₈ O ₄	High	Medit., Asian

Table 5: Continued...

Name	Molecular Formula	Risk Levels	Population at Risk
O-Acetylsalicylic Acid (acetylsalicylic acid)	C ₉ H ₈ O ₄	High	Medit., Asian
Oxidase, Urate (urate oxidase)		High	Medit., Asian
Pamaquine	C ₄₂ H ₄₅ N ₃ O ₇	High	All
Para-Aminobenzoic Acid (4-Aminobenzoic Acid)	C ₇ H ₇ N O ₂	Low	All
Paracetamol (acetaminophen)	C ₈ H ₉ N O ₂	Low	All
Pentaquine	C ₁₈ H ₂₇ N ₃ O	High	All
Phenacetin (acetophenetidin)	C ₁₀ H ₁₃ N O ₂	High	Medit., Asian
Phenazone (antipyrine)	C ₁₁ H ₁₂ N ₂ O	Low	All
Phenazopyridine	C ₁₁ H ₁₁ N ₅	High	Medit., Asian
Phenylbutazone	C ₁₉ H ₂₀ N ₂ O ₂	Low	All
Phenytoin	C ₁₉ H ₂₀ N ₂ O ₂	Low	All
Phynylhydrazine	C ₆ H ₈ N ₂	High	All
Phytomenadione (Vitamin K1)	C ₃₁ H ₄₆ O ₂	Low	All
Primaquine	C ₁₅ H ₂₁ N ₃ O	High	All
Probenecid	C ₁₃ H ₁₉ NO ₄ S	High	All
Procainamide	C ₁₃ H ₂₁ N ₃ O	Low	All
Proguanil (chlorguanidine)	C ₁₁ H ₁₆ Cl N ₅	Low	All
Pyrimethamine	C ₁₂ H ₁₃ Cl N ₄	Low	All
Quinidine	C ₂₀ H ₂₄ N ₂ O ₂	Low	All
Quinine	C ₂₀ H ₂₄ N ₂ O ₂	Low	All
Stibophen (2-(2-Oxido-3,5-Disulphonatophenoxy)-1,3,2-Benzodioxastibole-4-6-Disulphonate)	C ₁₂ H ₄ Na ₅ O ₁₆ S ₄ Sb	High	All
Streptomycin	C ₂₁ H ₃₉ N ₇ O ₁₂	Low	All
Sulfacetamide	C ₈ H ₁₀ N ₂ O ₃ S	High	All
Sulfacytine	C ₁₂ H ₁₄ N ₄ O ₃ S	Low	All
Sulfadiazine	C ₁₀ H ₁₀ N ₄ O ₂ S	Low	All
Sulfadimidine	C ₁₂ H ₁₄ N ₄ O ₂ S	High	All
Sulfafurazole (sulfafurazone, sulfisoxazole)	C ₁₁ H ₁₃ N ₃ O ₃ S	High	Medit., Asian
Sulfaguanidine	C ₇ H ₁₀ N ₄ O ₂ S	Low	All
Sulfamerazine	C ₁₁ H ₁₂ N ₄ O ₂ S	Low	All
Sulfamethoxazole	C ₁₀ H ₁₁ N ₃ O ₃ S	High	All
Sulfanilamide (Sulphanilamide)	C ₆ H ₈ N ₂ O ₂ S	High	All
Sulfapyridine	C ₁₁ H ₁₁ N ₃ O ₂ S	High	All
Sulfasalazine, Salazosulfapyridine (salazopyrin)	C ₁₈ H ₁₄ N ₄ O ₅ S	High	All
Thiazosulfone (thiazolesulfone)	C ₉ H ₉ N ₃ O ₂ S ₂	High	Medit., Asian
Tiaprofenic Acid	C ₁₄ H ₁₂ O ₃ S	Low	All
Tolonium Chloride, Tolonium Chloride (toluidine blue)	C ₁₅ H ₁₆ Cl N ₃ S	High	All
Trihexyphynidyl (benzhexol)	C ₂₀ H ₃₁ N O	Low	All
Trimethoprim	C ₁₄ H ₁₈ N ₄ O ₃	Low	All
Trinitrotoluene (2,4,6-Trinitrotoluene)	C ₇ H ₅ N ₃ O ₆	High	Medit., Asian
Tripelennamine	C ₁₆ H ₂₁ N ₃	Low	Medit., Asian

Food substances and herbs that should be avoided by G6PD deficient people

Fava Beans

Some prefer also to avoid red wine, all legumes, blueberries (also with yogurt), soya products, tonic water.

Chinese Herbs to Avoid: Cattle Gallstone Bezoar (Bos Taurus Domesticus), Honeysuckle (Lonicera Japonica),

Chimonanthus Flower(Chimonanthus Praecox), Huang Lian (黄连), 100% Pearl Powder

b. Favism

Favism is hemolytic anemia which occurs after the ingestion of fava beans especially fresh fava beans in G6PD-deficient individual and that's why the highest incidence rate of favism is at the time of harvest (1, 2, 15, 62).

Favism is always observed in people with G6PD deficiency, but all G6PD-deficient people do not develop hemolysis. Therefore, G6PD deficiency is an essential factor for the occurrence of favism but it is not enough (2, 15). Most cases of favism are observed in Mediterranean type of G6PD-deficient because the level of enzyme activity is very low and deficiency is severe, and occasionally in individuals with G6PD A⁻ variant (1, 2).

Clinical symptoms of favism are pallor, jaundice, hematuria and acute hemolytic anemia occurs 24-48 h after consumption of fava beans, suddenly (1, 15, 63). Ahaptoglobinemia was found in favism patients (64, 65). There is a low haptoglobin serum in G6PD deficient individuals. Haptoglobin is a serum protein of the alpha-2 fraction, binds to free hemoglobin whenever intravascular hemolysis occurs, the complex cleared by phagocytes and then haptoglobin disappeared from plasma (65, 66).

Vicine and convicine are pyrimidine glucoside that are abundant in fava beans and make up 6.7 g/100 g of dry weight of fava beans (67). These substances are metabolized by B-glucosidase in body to divicine and isoramnil that are unstable aglycons, and are oxidized rapidly and form hydrogen peroxide and superoxide anion. These metabolites oxidize reduced glutathione in normal and G6PD-deficient red blood cells (67-69). Depletion of reduced glutathione and impairment of some important enzyme induce oxidative stress in G6PD-deficient red blood cells and lead to acute hemolytic anemia called favism (2, 68, 69).

c. Infection-induced hemolysis

The most common cause of hemolysis is possibly the infection in G6PD-deficient individuals (1, 2). Viral, bacterial and rickettsial infections, especially hepatitis, pneumonia and typhoid fever induce hemolysis in G6PD-deficient people (1). During phagocytosis, leukocytes may release ac-

tive oxygen species that induce oxidative stress to the erythrocytes, and will damage them and will cause hemolysis (1, 2).

2. Chronic non spherocytic hemolytic anemia

Some of the rare variants of G6PD deficiency which are designated as class 1 variants, according to the classification of WHO, are associated with Chronic non spherocytic hemolytic anemia (1, 2, 26).

Individuals in this class have very low enzyme activity (26) and suffer from hemolytic anemia even when oxidative agents are not present (2). They are variably anemic in steady state condition (1, 2). Splenomegaly is commonly present. The disorder is usually identified during infancy and childhood (15).

3. Neonatal jaundice

Serum bilirubin levels are determined by production of bilirubin, bilirubin conjugation and elimination. Hyperbilirubinemia is caused by the imbalance between the rate of production of bilirubin, the end product of heme metabolism, and restricted excretion of bilirubin in newborns (70, 71). The average total serum bilirubin level is usually 5-6 mg per dl (86-103 μmol per l) in full term newborns (71). Hyperbilirubinemia occurs due to either physiologic or pathologic causes (71). Physiologic jaundice occurs when the serum total bilirubin is in a range of 7 to 17 mg per dl (104-291 μmol per l). A total serum bilirubin level higher than 17 mg per dL is considered pathologic hyperbilirubinemia. One of the common risk factors for pathologic hyperbilirubinemia in newborn infants is deficiency of G6PD enzyme (71). Deficiency of this enzyme is the most prevalent enzymopathy in red blood cells that causes hemolysis and hyperbilirubinemia (72). Hyperbilirubinemia can be very severe in G6PD-deficiency and induces permanent damage to the brain and causes kernicterus and death (72). Overproduction of unconjugated bilirubin and lack of proper management of hyperbilirubinemia cause changes in the mitochondria of the basal ganglia and leads to impaired mitochondrial respiration, and also induce apoptosis, and cause bilirubin encephalopathy (73).

The pathogenesis of hyperbilirubinemia in G6PD-deficient newborn babies is different from that in G6PD-normal ones. Meanwhile hemolysis is considered to be a principal cause of bilirubinemia in G6PD-normal neonates; but diminished bilirubin conjugation would be the main cause of hyperbilirubinemia in G6PD-deficient newborn infants (70). Profile of serum bilirubin in newborns with G6PD-deficiency showed that the amount of total and unconjugated bilirubin is high and conjugated bilirubin is low. These levels of different forms of bilirubin that are seen in the neonates are similar in conditions of partial deficiency of the bilirubin conjugating enzyme UDP glucuronosyltransferase, such as Gilberts syndrome. High levels of unconjugated bilirubin in G6PD-deficient neonates are the result of an interaction between G-6-PD deficiency and variant promoter for the gene encoding this enzyme, UDP glucuronosyltransferase (74).

Although there is a natural immaturity of bilirubin conjugation in neonates, the bilirubin conjugation ability of G6PD-deficient neonates who are also hyperbilirubinemic is even less efficient. Bilirubin conjugation ability in G6PD-deficient neonates may become worse due to increased hemolysis and more bilirubin production (70, 74).

Diagnosis

Beutler fluorescent spot test, dichlorophenol indophenol decolorization and quantitative spectrophotometric assay are methods used for the diagnosis of G6PD deficiency (15, 75). Beutler fluorescent spot test is the most acceptable method for screening of G6PD deficiency (76). During this method, the rate of NADPH production from NADP by G6PD is measured under ultraviolet light (15, 66). G6PD non-deficient blood samples fluoresce brightly, but deficient samples show little or no fluorescence (75-77).

When individuals are actively hemolyzing, the test can show normal results wrongly, because sustaining younger erythrocytes show normal enzyme activity or near normal. Thus the individuals should be screened several weeks after a hemolytic episode (15, 75).

By using the Quantitative spectrophotometric method, activity of G6PD Enzyme was determined by measuring the rate of the reduction of NADP to NADPH in the presence of G6P and hemolysate (15, 76). Another Screening method for G6PD deficiency is dichlorophenol-indophenol (DPIP) dye decolorization. This method is used for determining of G6PD in red blood cells. By this method heterozygotes are diagnosed easily and is used when a large number of population are screened for determination of G6PD deficiency (15).

Laboratory tests and clinical features in patients with G6PD deficiency and hemolysis are: Hemoglobinuria, Increased indirect bilirubin, Elevated serum lactate dehydrogenase,

Low serum haptoglobin (up to ahaptoglobinemia), Complete blood count (CBC), elevated reticulocyte count, Heinz bodies are presented on peripheral blood smear, Comb^s test (the test in hemolysis phase is negative) (77, 78).

Treatment

The main treatment for G6PD deficiency is avoidance of oxidative agents like infection, fava beans and oxidative drugs that induce hemolysis (1, 4). Hemolysis may be so severe that it may even require blood transfusion (2). Screening of newborns for early diagnosis of G6PD deficiency and proper education can reduce the incidence of clinical symptoms (1). These methods have been successfully applied in northern Sardinia and the result was that the incidence of Favism has become less and also in Singapore kernicterus in the neonates is now very rare. Proper control and early treatment in neonatal jaundice are very important (1). Phototherapy is used to decrease bilirubin concentration and when the bilirubin levels exceeds 20 mg/dl, exchange transfusion is necessary (2). In regions that G6PD deficiency is prevalent, there is a grave danger of giving G6PD-deficient blood to such newborns and this has to be prevented (1). Using desferrioxamine could make the attacks of Favism less severe. Use of Sn-mesoporphyrin (SnMP) that inhibits heme oxygenase and decreases bilirubin production in G6PD-deficient neonates has decreased the need

for phototherapy and exchange transfusion (1, 79, 80). Because of some adverse effects of Sn-mesoporphyrin, SnMP should only be used for neonates who are in a clear danger of developing bilirubin-induced neurologic dysfunction or those who are taking part in clinical trials (79). In clinical experiments, use of antioxidants like vitamin E and selenium have not shown any benefit for the treatment of G6PD deficiency (1).

Conclusion

G6PD deficiency is one of the most common X-link inherited hemolytic disorders reported, affecting around 400 million people worldwide. The main function of G6PD is to protect the RBC against oxidative damage. The most important way for prevention and reduction in the incidence rate of clinical symptoms of G6PD deficiency is to avoid oxidative agents like infection, fava beans and oxidative drugs that induce hemolysis, also screening of newborns for early diagnosis of G6PD deficiency and proper education is recommended.

Acknowledgements

The authors are both thankful to Prof A Jazayeri, School of Public Health, and Dr H Sadighi, Genetic Clinic, for reading the manuscript and giving some suggestions.

This study was supported partly by Iran TWAS chapter based at ISMO in Tehran.

The authors declare that they have no conflict of interests.

References

1. Mehta A, Mason PJ, Vulliamy TJ (2000). Glucose-6-phosphate dehydrogenase deficiency. *Best Prac Res Clin aematol*, 13: 21-38.
2. Beutler E (1994). G6PD deficiency. *Blood*, 84(11): 3613-36.
3. Noori-Dalooi MR, Najafi L, Ganji SM, Hajebrahimi Z, Sanati M H (2004). Molecular identification of mutations in G6PD gene in patients with favism in Iran. *J Physiol Biochem*, 60: 273-77.
4. Prchal JT, Gregg XT (2005). Red Cell Enzymes. *Hematol*, 1: 19-23.
5. Greene LS (1993). Deficiency as Protection Against falciparum Malaria: An Epidemiological Critique of Population and Experimental Studies. *Yearbook of Phys Anthropol*, 36: 153-78.
6. Jain M, Brenner DA, Cui L, Lim CC, Wang B, Pimentel DR, et al. (2003). Glucose-6-Phosphate Dehydrogenase Modulator Cytosolic Redox Status and Contractile Phenotype in Adult Cardiomyocytes. *Circ Res*, 93: e9.
7. Shihabi A, Li WG, Miller FJ, Weintraub NL (2002). Antioxidant therapy for atherosclerotic vascular disease: the promise and the pitfalls. *Am J Physiol Heart Circ Physiol*, 282: H797-H802.
8. Leopold JA, Loscalzo J (2005). Oxidative Enzymopathies and Vascular Disease. *Arterioscler Thromb Vasc Biol*, 25:1332.
9. Tsun-Yee Chiu D, Liu TZ (1997). Free radicals and oxidative damage in human blood cells. *J Biochem Sci*, 4: 256-59.
10. Halliwell B (1992). Reactive Oxygen Species and the Central Nervous System. *J Neurochem*, 59 (5), 1609-23.
11. Savitha S, Tamilselvan J, Anusuyadevi M, Panneerselvam C (2005). Oxidative stress on mitochondrial antioxidant defense system in the aging process: role of DL-alpha-lipoic acid and L-carnitine. *Clin Chim Acta*, 355: 173-80.
12. Fico A, Paglialonga F, Cigliano L, Abrescia P, Verde P, Martini G, et al. (2004). Glucose-6-phosphate dehydrogenase plays a crucial role in protection from redox-stress-induced apoptosis. *Cell Death Differ*, 11: 823-831.
13. Tian WN, Braunstein LD, Apse K, Pang J, Rose M, Tian X, et al. (1999). Importance of glucose-6-phosphate dehydrogenase activity in cell death. *Am J Physiol Cell Physiol*, 276: 1121-31.
14. Abraham S, Soundararajan CC, Vivekanandhan S, Behari M (2005). Erythrocyte

- antioxidant enzymes in Parkinson's disease. *Indian J Med Res*, 121: 111-15.
15. Mohanty D, Mukherjee MB, Colah RB (2004). Glucose-6-phosphate dehydrogenase deficiency in India. *Symposium*, 71(6): 525-29.
 16. Cappellini MD, Fiorelli G (2008). Glucose-6-phosphate dehydrogenase deficiency. *Lancet*, 5; 64-74.
 17. Kwiatkowski DP (2005). How Malaria Has Affected the Human Genome and What Human Genetics Can Teach Us about Malaria. *Am J Hum Genet*, 77(2): 171-92.
 18. Fortin A, Stevenson MM, Gros P (2002). Susceptibility to malaria as a complex trait: big pressure from a tiny creature. *Hum Mol Genet*, 11: 2469-78.
 19. Farhud DD, Sadighi H, Amirshahi P, Tavakkoli F Sh (1993). Serum level measurements of Gc, Cp, IgG, IgA and IgM in patients with favism in Iran. *Iranian J Publ Health*, 22:1-4.
 20. Tripathy V, Reddy BM (2007). Present status of understanding on the G6PD deficiency and natural selection. *Rev Article*, 53: 193-202.
 21. Anonymous (2006). Glucose-6-phosphate dehydrogenase deficiency. Genetics Home Reference. Available from: ghr.nlm.nih.gov/condition=glucose6phosphatedehydrogenasedeficiency
 22. Usanga EA, Ameen R (2000). Glucose-6-Phosphate Dehydrogenase Deficiency in Kuwait, Syria, Egypt, Iran, Jordan and Lebanon. *Hum Hered*, 50: 158-61.
 23. Vulliamy TJ, Urso MD, Battistuzzi G, Estrada M, Foulkes N S, Martini G, et al. (1988). Diverse point mutations in the human glucose-6-phosphate dehydrogenase gene cause enzyme deficiency and mild or severe hemolytic anemia. *Proc Natl Acad Sci U S A*, 85: 5171-75.
 24. Au SW, Gover S, Lam VM, Adams MJ (2000). Human glucose-6-phosphate dehydrogenase: the crystal structure reveals a structural NADP⁺ molecule and provides insights into enzyme deficiency. *Struc*, 8(3): 293-303.
 25. Anonymous. Glucose-6-phosphate dehydrogenase; G6PD. Johns Hopkins University. Available from: www.ncbi.nlm.nih.gov/entrez/dispomim.cgi?id=305900.
 26. Wijk RV, Huizinga EG, Prins I, Kors A, Rijksen G, Bierings M, et al. (2004). Distinct phenotypic expression of two de novo missense mutations affecting the dimer interface of glucose-6-phosphate dehydrogenase. *Blood Cells Mol Dis*, 32: 112-17.
 27. Hundsdoerfer P, Vetter B, Kulozik AE (2002). Chronic Haemolytic Anaemia and Glucose-6-Phosphate Dehydrogenase Deficiency. *Acta Haematol*, 108:102-5.
 28. WHO Working Group. Glucose-6-phosphate dehydrogenase deficiency. *Bull WHO* 1989; 67: 601-11.
 29. Corons JV, Kuhl W, Pujades MA, Beutler E (1990). Molecular Genetics of the Glucose-6-Phosphate Dehydrogenase (G6PD) Mediterranean Variant and Description of a New G6PD Mutant, G6PD Andalus 1361A. *Am J Hum Genet*, 47:575-79.
 30. Noori-Dalooi MR, Soltanian S, Mohammad Gangi SH, Yousefi A, Hejazi S, Banihashem A, et al. (2006). Molecular Identification of the Most Prevalent Mutations of Glucose-6-Phosphate Dehydrogenase (G6PD) Gene in deficient Patients in Khorasan Province of Iran. *J Sci I R Iran*, 17(2): 103-106.
 31. Warsy AS, El-Hazmi MAF (2001). G6PD deficiency, distribution and variants in Saudi Arabia. *Ann Saudi Med*, 21:174-77.
 32. AlFadhli S, Kaaba S, Elshafey A, Salim Matra, Alawadi A, Bastaki L (2005). Molecular Characterization of Glucose-6-Phosphate Dehydrogenase Gene Defect in the Kuwaiti Population. *Arch Pathol Lab Med*, 129:1144-47.
 33. Nafa Kh, Reghis A, Osmani N, Baghli L, Aït-Abbes H, Benabadjji M, et al. (1994). At least five polymorphic mutants account

- for the prevalence of glucose-6-phosphate dehydrogenase deficiency in Algeria. *Hum Genet*, 94.
34. Vives Corrons JL, Zarza R, Aymerich JM, Boixadera J, Carrera A, Colomer D, et al. (1997). Molecular analysis of glucose-6-dehydrogenase deficiency in Spain. *Sangre*, 42(5):391-8.
 35. Medina MD, Vaca G, Lopez-Guido B, Westwood B, Beutler E (1997). Molecular Genetics of Glucose-6-Phosphate Dehydrogenase Deficiency in Mexico. *Blood Cells Mol Dis*, 23: 88-94.
 36. Ainoon O, Yu YH, Amir Muhriz AL, Boo NY, Cheong SK, Hamidah NH (2003). Glucose-6-phosphate dehydrogenase (G6PD) variants in Malaysian Malays. *Hum Mutat*, 21(1):101.
 37. Nuchprayoon I, Sanpavat S, Nuchprayoon S (2002). Glucose-6-phosphate dehydrogenase (G6PD) mutations in Thailand: G6PD Viangchan (871G> A) is the most common deficiency variant in the Thai population. *Hum Mutat*, 19: 185- 85.
 38. Noori-Dalooi MR, Yousefi A, Mohammad Ganji S, Hejazi SH, Soltanian S, Sanei Moghadam E, et al. (2005). Molecular Identification of the Most Prevalent Mutation of Glucose-6-Phosphate Dehydrogenase Gene in Deficient Patients in Sistan and Balochestan Province of Iran. *J Sci I R Iran*, 16.
 39. Karadsheh NS, Gelbart T, Schulten HJ, Efferth T, Awidi A(2005). Relationship between Molecular Variants and Clinical Manifestations in Twelve Glucose-6-Phosphate Dehydrogenase-Deficient Patients in Jordan. *Acta Haematol*, 114:125-26.
 40. Rahimi Z, Vaisi-Raygani A, Ronald L (2006). Molecular characterization of glucose-6-phosphate dehydrogenase deficiency in the Kurdish population of Western Iran. *Blood Cells Mol Dis*, 37: 91-4.
 41. Abolghasemi H, Mehrani H, Amid A (2004). An update on the prevalence of glucose-6-phosphate dehydrogenase and neonatal jaundice in Tehran neonates. *Clin Biochem*, 37:241-44.
 42. Mesbah-Namin SA, Sanati MH, Mowjood A, Mason PJ, Vulliamy TJ, Noori-Dalooi MR (2002). Three major glucose-6-phosphate dehydrogenase-deficient polymorphic variants identified in Mazandaran state of Iran. *British J Haematol*, 117: 763-4.
 43. Mortazavi Y, Mirimoghaddamm E, Pourfathollah AA (2003). 8th Annual Congress of the European Hematology Association. *Hematol J*, 4: 155.
 44. Ohkura K, Miyashita T, Nakajama H, Matsumoto H, Matsutomo K, Rahabar S, et al. (1984). Distribution of polymorphic traits in Mazandaranian and Guilanian in Iran. *Hum Hered*, 34: 27-39.
 45. Pishva N, Amoozgar H (2001). Hyperbilirubinemia following exchange transfusion with G6PD deficient donor blood. *Ir J Med Sci*, 26:143-45.
 46. Khalili D, Jafroodi M, Sajedi SA, Shameli Rad M, Abdollahi F, Alipour Kanafi K, et al. (2007). Survey of the prevalence of glucose-6-phosphate dehydrogenase deficiency in Rasht-Iran. *J Gilan Univ Med Sci*, 63: 51-6.
 47. Beutler E. Erythrocyte disorders: Anemias due to increased destruction of erythrocytes with enzyme deficiencies, Glucose-6-phosphate dehydrogenase deficiency. G6PD Deficiency Association is affiliated with Italian Federation for Rare Diseases. Available at: www.g6pd.org/favism/english/print.mv?pgid=beutler
 48. Pronsky ZM, Crowe SJP (2008). Assessment: Food-Drug interactions. In: Krause's Food and Nutrition Therapy. eds, Mahan and Escott- Stump. 12th ed, Saunders Elsevier Inc. St. Louis, Missouri, P. 434.
 49. Lowitt MH, Shear NH (2001). Pharmacogenomics and dermatological therapeutics. *Arch Dermatol*, 137.
 50. WHO (2007). The Ethical, Legal and Social implications of pharmacogenomics in de-

- veloping countries. *Rep Int Group experts*, pp. 3-13.
51. Nicol CJ, Zielenski J, Tsui LC, Wells PG (2000). An embryoprotective role for glucose-6-phosphate dehydrogenase in developmental oxidative stress and chemical tetraetogenesis. *Faseb J*, 14: 111-27.
 52. Bolchoz LJC, Morrow JD, Jollow DJ, McMillan DC (2002). Primaquine-Induced Hemolytic Anemia: Effect of 6-Methoxy-8-hydroxylaminoquinoline on Rat Erythrocyte Sulfhydryl Status, Membrane Lipids, Cytoskeletal Proteins, and Morphology. *Pharmacol Exp Ther*, 303: 141-48.
 53. Bowman ZS, Oatis JE, Whelan JL, Jollow DJ, McMillan DC (2004). Primaquine-Induced Hemolytic Anemia: Susceptibility of Normal versus Glutathione-Depleted Rat Erythrocytes to 5-Hydroxyprimaquine. *J Pharmacol Exp Ther*, 309: 79-85.
 54. Zinkham WH, Oski FA (1996). Henna: A Potential Cause of Oxidative Hemolysis and Neonatal Hyperbilirubinemia. *Pediatr*, 97: 707-9.
 55. McMillan DC, Sarvate SD, Oatis JE, Jollow DJ (2004). Role of Oxidant Stress in Lawsone-Induced Hemolytic Anemia. *Toxicol Sci*, 82: 647-55.
 56. Seyedzadeh A, Hemmati M, Gheiny S (2007). Henna- Induced Severe Hemolysis: In Glucose 6- Phosphate Dehydrogenase Deficiency. *Pak J Med Sci*, 23: 119-21.
 57. Raupp P, Hassan JA, Varughese M, Kristiansson B (2001). Henna causes life threatening haemolysis in glucose-6-phosphate dehydrogenase deficiency. *Arch Dis Child*, 85: 411-12.
 58. Zinkham WH, Oski FA (1996). Henna: A Potential Cause of Oxidative Hemolysis and Neonatal Hyperbilirubinemia. *Pediatr*, 97: 707-9.
 59. Ko CH, Li K, Ng PC, Fung KP, Li CL, Wong RP, et al. (2006). Pro-oxidative effects of tea and polyphenols, epigallocatechin-3-gallate and epigallocatechin, on G6PD-deficient erythrocytes in vitro. *Int J Mol Med*, 18:987-994.
 60. Pronsky ZM, Crowe JP (2008). Assessment: Food- Drug Interactions. In: Krause's Food and Nutrition Therapy. Eds, Mahan and Escott- Stump. 12th ed, Saunders Elsevier Inc. St. Louis, Missouri, P.435.
 61. Anonymous. G6PD Deficiency Association is affiliated with Italian Federation for Rare Diseases. Available at: www.g6pd.org/favism/english/index.mv?pgid=intro.
 62. McMillan DC, Jollow DJ (1999). Favism: divicine hemotoxicity in the rat. *Toxicol Sci*, 51: 310-16.
 63. Hedayat SH, Farhud DD, Montazami K, Ghadiryan P (1981). The Pattern of Bean Consumption, Laboratory Findings in Patients with Favism, G-6-P-D Deficient, and a Control Group. *J Trop Pediatr*, 27: 110-13.
 64. Farhud DD, Amirshahi P (1985). Ahaptoglobinemia in favism patients from Iran. *Acta Anthropogenet*, 9: 206-13.
 65. Farhud DD, Amirshahi P, Hedayat SH (1979). The distribution of haptoglobin type in Bandarabas. *Iranian J Publ Health*, 7: 181-82.
 66. Mobasheri MB, Farhud DD (2003). Investigation of ahaptoglobinemia and its association with malaria endemicity in south of Iran. *Iranian J Publ Health*, 32: 19-22.
 67. Arese P, Bosia A, Naitana A, Gaetani S, D'Aquino M, Gaetani GF (1981). Effect of divicine and isouramil on red cell metabolism in normal and G6PD-deficient (Mediterranean variant) subjects. Possible role in the genesis of favism. *Prog Clin Biol Res*, 55: 725-46.
 68. Pedersen JZ, Musci G, Rotilio G (1988). Electron Spin Resonance Characterization of the Radicals Produced by Enzymatic or Chemical Cleavage of Vicine. *Biochem*, 27: 8534-36.
 69. McMillan DC, Schey KL, Meier GP, Jollow DJ (1993). Chemical Analysis and Hemo-

- lytic Activity of the Fava Bean Aglycon Divicine. *Chem Res Toxicol*, 6:439-44.
70. Kaplan M, Muraca M, Vreman HJ, Hammerman C, Vilei MT, Rubaltelli FF, et al. (2005). Neonatal bilirubin production-conjugation imbalance: effect of glucose-6-phosphate dehydrogenase deficiency and borderline prematurity. *Arch Dis Child Fetal Neonat Ed*, 90: F123-F127.
71. Dennery PA, Seidman DS, Stevenson DK (2001). Neonatal Hyperbilirubinemia. *N Engl J Med*, 344: 581-90.
72. Kaplan M, Hammerman C (2000). Glucose-6-Phosphate Dehydrogenase Deficiency: A Worldwide Potential Cause of Severe Neonatal Hyperbilirubinemia. *Pediatr Rev Neo Rev*, 1: e32-e39.
73. Groenendaal F, der Grond JV, De Vries LS (2004). Cerebral Metabolism in Severe Neonatal Hyperbilirubinemia. *Pediatr*, 114: 291-94.
74. Kaplan M, Muraca M, Hammerman C, Vilei MT, Leiter C, Rudensky B, et al. (1998). Bilirubin Conjugation, Reflected by Conjugated Bilirubin Fractions, in Glucose-6-Phosphate Dehydrogenase-deficient Neonates: A Determining Factor in the Pathogenesis of Hyperbilirubinemia. *Pediatr*, 102 (3): e37.
75. Zahed Pasha A, Ahmadpoor M, Zahed Pasha Y (2006). Glucose-6-phosphate dehydrogenase Enzyme Deficiency. *J Babol Univ Med Sci*, 8(1).
76. Markić J, Krželj V, Markotić A, Marušić E, Stričević L, Zanchi J, et al. (2006). High Incidence of Glucose-6-phosphate Dehydrogenase Deficiency in Croatian Island Isolate: Example from Vis Island, Croatia. *Croat Med J*, 47: 566-70.
77. en.wikipedia.org/wiki/Glucose-6-phosphate_dehydrogenase_deficiency.
78. Frank EJ (2005). Diagnosis and Management of G6PD Deficiency. *Am family phys*, 72: 1277-82.
79. Wong RJ, Bhutani VK, Vreman HJ, Stevenson DK (2007). Tin Mesoporphyrin for the Prevention of Severe Neonatal Hyperbilirubinemia. *Neo Rev*, 8.
80. Martinez JC, Garcia HO, Otheguy LE, Drummond GS, Kappas A (1999). Control in Severe Hyperbilirubinemia in Full-term Newborns with the Inhibitor in Bilirubin Production. *Pediatr*, 103: 1-5.