



## **The Efficient Estimation of Motor Unit Excitability Parameters in Needle Electromyography Experiments**

**Nammam Ali AZADI<sup>1,2</sup>, Keramat Nouri JELYANI<sup>3</sup>, \*Daem ROSHANI<sup>1,2</sup>**

1. Dept. of Epidemiology and Biostatistics, Medical School, Kurdistan University of Medical Sciences, Sanandaj, Iran
2. Kurdistan Research Center for Social Determinants of Health, Medical School, Kurdistan University of Medical Sciences, Sanandaj, Iran
3. Dept. of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

**\*Corresponding Author:** Email: d.roshany@muk.ac.ir

(Received 23 Dec 2013; accepted 11 Jan 2014)

### **Dear Editor-in-Chief**

Moving our body is the outcome of our muscles' contraction. Muscles themselves are composed of groups of muscle fibres each controlled by a single motor neuron. The motor neuron and its associated muscle fibres form the elementary functional unit of the neuromuscular system, the so-called motor unit (MU). By activating more or less MUs, the amount of generated muscle force can be increased or decreased. This is the key concept in testing muscle activities in patients with muscular disorders (1).

Motor unit activities (or equivalently, nerve excitability) can be studied by means of the so-called threshold tracking techniques, widely available by use of two widespread electrophysiological techniques; surface electromyography (SEMG) and needle electromyography (NEMG). Threshold tracking assesses excitability property of an MU by recording its response to a range of stimulus intensities required to elicit a target response (2). This experiment yields to a stimulus-response curve from which one can estimate MU's threshold (the stimulus intensity that elicits a response to 50% of the stimuli) and the range of stimulus intensities over which the MU shows a stochastic behavior. One demand, therefore, is to obtain efficient estimates of these parameters. For estima-

tion of MU's threshold, various strategies are available (3). However, the judgments underlying these approaches are mainly subjective and statistical methods are seldom used (4).

Two main interests in threshold tracking studies are; i) to develop a novel model-based approach in order to obtain reliable estimation of the MU's excitability parameters, and ii) to obtain an efficient experimental design for both NEMG and SEMG experiments. Azadi, 2012 has addressed both issues; using statistical models for parameter estimation in NEMG experiment and a Bayesian experimental design for SEMG technique (5).

### **Acknowledgement**

The authors declare that there is no conflict of interests.

### **References**

1. Ignacio RC, Luis GU, Armando MT (2012). Motor Unit Action Potential Duration: Measurement and Significance. In: *Adv in Clin Neurophysiol*. Eds, Ajeena. Rijeka, InTech. pp. 133-160,
2. Bostock KC, Burke D (1998). Threshold tracking techniques in the study of human

- peripheral nerve. *Muscle Nerve*, 21(2): 137-58.
3. Stein RB, Yang JF (1990). Methods for estimating the number of motor units in human muscles. *Annals of Neurology*, 28(4): 487-95
  4. Ridall, PG, Pettitt AN, McCombe PA, Henderson RD (2006). Motor unit number estimation, a Bayesian approach. *Biometrics*, 62(4): 1235-50
  5. Azadi, NA. Modeling the excitability behaviour of a single motor unit and experimental design for the surface EMG test [PhD thesis]. Mathematics and Statistics Department, Lancaster University, UK; 2011.

Archive of SID