



Identification of *Malassezia* Species Isolated from Patients with Pityriasis Versicolor Using PCR-RFLP Method in Markazi Province, Central Iran

***Mojtaba DIDEHDAR¹, Amir Sayed Ali MEHBOD², Zahra ESLAMIRAD³, Mahdi MOSAYEBI³, Reza HAJIHOSSEIN³, Behzad GHORBANZADE³, Mahmoud Reza KHAZAET⁴**

1. Molecular and Medicine Research Center, Arak University of Medical Sciences, Arak, Iran
2. Dept. of Medical Parasitology and Mycology, Army University of Medical Sciences, Tehran, Iran
3. Dept. of Medical Parasitology and Mycology, Arak University of Medical Sciences, Arak, Iran
4. Dept. of Laboratory Science, Islamic Azad University of Arak, Arak, Iran

***Corresponding Author:** Email: didehdar_m@arakmu.ac.ir

(Received 12 Oct 2013; accepted 15 Feb 2014)

Abstract

Background: The lipophilic yeasts of *Malassezia* species are members of the normal skin microbial that are cause of pityriasis versicolor. Pityriasis versicolor is a common superficial fungal infection with world-wide distribution. The phenotypic methods for identification of *Malassezia* species usually are time consuming and unreliable to differentiate newly identified species. But DNA-based techniques rapidly and accurately identified *Malassezia* species. The purpose of this study was isolation and identification of *Malassezia* Species from patients with pityriasis versicolor by molecular methods in Markazi Province, Central Iran in 2012.

Methods: Mycologic examinations including direct microscopy and culture were performed on clinical samples. DNA extraction was performed from colonies. The ITS1 region of rDNA from isolates of *Malassezia* species were amplified by PCR reaction. The PCR were digested by *Cfo I* enzyme.

Results: From 70 skin samples, were microscopically positive for *Malassezia* elements, 60 samples were grown on culture medium (85.7%). Using PCR-RFLP method, that was performed on 60 isolates, 37(61.6%) *M. globosa*, 14(23.3%) *M. furfur*, 5(8.4%) *M. sympodialis* and 4(6.7%) *M. restricta* were identified. In one case was isolated *M. globosa* along with *M. restricta*.

Conclusion: The PCR-RFLP method is a useful and reliable technique for identification of differentiation of *Malassezia* species.

Keywords: *Malassezia* spp., Pityriasis versicolor, Identification, PCR-RFLP, Iran

Introduction

Malassezia species are lipophilic yeasts which are as part of the normal flora of human skin, and are found in mostly of healthy adults. These yeasts are the cause of pityriasis versicolor and *Malassezia* folliculitis, and also implicated in the pathogenesis of common skin disorders such as seborrheic

dermatitis, psoriasis, and atopic dermatitis (1). Pityriasis versicolor is a superficial fungal infection that usual clinical feature is slightly scaly patches of variable color (either pink, brown, or white) with irregular margin, most commonly found on the trunk and shoulders. These lesions different in

color, and can be hypo- or hyperpigmented. The relationship between *Malassezia* species and their pathogenetic relationship was determined.

In 1996, Guého et al. classified *Malassezia* genus into seven species, *M. furfur*, *M. obtusa*, *M. globosa*, *M. slooffiae*, *M. sympodialis*, *M. pachydermatis*, and *M. restricta*, based on morphological and biochemical characteristics (2). However, with the development of molecular methods such as polymerase chain reaction (PCR), additional five species have been described which are *M. yamatoensis*, *M. nana*, *M. japonica*, *M. equi*, and *M. dermatitis*(3-8).

However, these phenotypic methods are usually time consuming, lack sufficient discriminatory power, and are unable to unambiguously differentiate newly identified species. Various DNA-based molecular methods have been described to overcome this problem (9).

The aim of this study was to identify *Malassezia* species on their colonies, which were grown in culture media using 26S rDNA polymerase chain reaction restriction fragment length polymorphism (PCR-RFLP).

Materials and Methods

Subjects

Patients with clinically suspected with pityriasis versicolor, referred to the Mycology Laboratory of the Arak University of Medical Sciences were recruited. Seventy patients (85.2%) were confirmed with pityriasis versicolor, based on the presence of both hyphae and yeasts in direct microscopy.

Collection and culture of sample

The samples were collected by scraping. The sampled skin was inoculated in modified Leeming and Notman culture medium (1% w/v peptone, 1% w/v glucose, 0.2% w/v yeast extract, 0.8% desiccated ox bile, 0.1% v/v glycerol, 0.05% w/v glycerol monostearate, 0.5% v/v Tween 60, and 2% v/v oleic acid, 1% w/v agar in distilled water) supplemented with cyclohexamide (0.5%) and chloramphenicol (0.05%), and was incubated at 32°C for two weeks. Isolated yeasts in 30% glycerol solution at -70 °C were stored in the freezer.

DNA extraction and 26S rDNA PCR

Genomic DNA was extracted using glass bead method (10). Primers were selected to allow the amplification of target DNA in all species. Their sequences were: forward, 5'-TAACAAGGAT-TCCCCTAGTA-3' and reverse, 5'-ATTAC-GCCAGCATCCTAAG-3' (11).

PCR amplification was performed in a final volume of 50 µl. Each reaction contained 1 µl of template DNA, 0.5 µM of each primer, 0.20 mM of each deoxynucleoside triphosphate (dNTPs), 5 µl of 10x PCR buffer, and 1.25 U of Taq polymerase. An initial denaturation step at 94 °C for 5 min was followed by 30 cycles of denaturation at 94 °C for 45 s, annealing at 55 °C for 45 s, and extension at 72 °C for 1 min, with a final extension step of 72 °C for 7 min. Amplified products were visualized by 1.5% (w/v) agarose gel electrophoresis in TBE buffer, stained with ethidium bromide (0.5 µgml⁻¹), and photographed under UV trans illumination.

RFLP analysis

The *Cfo* I enzyme (Roche Diagnostics, Mannheim, Germany) was used in this study (7). The restriction enzyme reaction was performed by incubating a 21.5 µl aliquot of PCR product with 10 U of the enzyme in a final reaction volume of 25 µl at 37 °C for 3 h. After the reaction, RFLP pattern was analyzed with DNA fragments in 2% agarose gel electrophoresis and staining with ethidium bromide.

Results

From 70 skin samples, which were microscopically positive for *Malassezia* elements, 60 samples were grown on culture medium (85.7%).

Using PCR reaction to amplify the 26s rDNA region of all *Malassezia* species, 580 bp PCR bands were confirmed in all of isolates (Fig.1). RFLP analysis of PCR products of the ITS1 region with restriction enzyme *Cfo* I was undertaken on 60 isolates.



Fig.1: 26S rDNA PCR products before digestion with *CfoI* enzyme. Lanes 1-2: *M. globosa*, Lane 3:*M. globosa*&*M. restricta*, Lane 4: *M. furfur*, Lane 5: *M. sympodialis*, Lane 6: *M. restricta*, M: 50 bp ladder

Using enzyme *CfoI* (Roche Diagnostics, Mannheim, Germany), we could distinguish four different species, including 37 (61.6%) *M. globosa*, 14 (23.3%) *M. furfur*, 5 (8.4%) *M. sympodialis* and 4 (6.7%) *M. restricta*. In one case was isolated *M. globosa* along with *M. restricta*. In this study *M. pachydermatis*, *M. obtusa*, *M. slooffiae* and other *Malassezia* species were not isolated (Fig. 2).




Fig. 2: 26S rDNA PCR products after digestion with *CfoI* enzyme. Lanes 1-2:*M. globosa*, Lane 3: *M. globosa*&*M. restricta*, Lane 4: *M. furfur*, Lane 5: *M. sympodialis*, Lane 6: *M. restricta*, M: 50 bp ladder

Discussion

The *Malassezia* genus has undergone several taxonomic revisions since 1996 (2). Recently, on the basis of DNA relatedness, five new species have been included: *M. dermatis*, *M. nana*, *M. japonica*, *M. yamatoensis*, and *M. aequi* (3-8).

Accurate and reproducible methods of species identification are essential for epidemiological purposes.

Various molecular methods of characterizing species of *Malassezia* including Pulsed-field gel electrophoresis (PFGE) of *Malassezia* species, internal transcribed spacer 1 (ITS1) ribosomal DNA sequences, AFLP genotyping and random amplified polymorphic DNA (RAPD) have been used (12-14). But this methods are time consuming and too expensive for use as a routine diagnostic method.

Recently, PCR-RFLP method used for identification of *Malassezia* species from colony and patient skin scales (11, 13, 15-17). Thus in this study, we used PCR-RFLP for identification of *Malassezia* species in Arak City, Iran.

In our study, 70 skin samples were microscopically positive for *Malassezia* elements and 60 samples were grown on culture medium (85.7%). The PCR-RFLP performed on 60 colonies and identified common species of *Malassezia*. The commonest of species was *M. globosa*, that was Similar to other studies (11,17-24). However, Makimura et al., isolated *M. furfur* as the commonest of species (9). In this study, in one case *M. globosa* long with *M. restricta* was isolated.

Among *Malassezia* species of human skin flora (25-27), *M. sympodialis* is the dominant species. Predominance of *M. globosa* from patients with pityriasis versicolor can be due to pathogenic characteristics of this species.

Conclusion

M. globosa was the commonest of *Malassezia* species isolated from patients with pityriasis versicolor in Arak City. This study provides important data for epidemiological status of *Malassezia* species in center of Iran.

Ethical considerations

Ethical issues (Including plagiarism, Informed Consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Acknowledgements

This project was supported by Arak University of Medical Sciences, Arak, Iran. The authors declare that there is no conflict of interests.

References

- Janik MP, Heffernan MP (2008). *Yeast infections: candidiasis, pityriasis versicolor*. In: *Fitzpatrick's dermatology in general medicine*. Eds, Wolff K, Goldsmith LA, Katz SI, Gilchrist BA, Paller AS, Leffell DJ. 7th ed. McGraw-Hill. New York, pp. 1822-1830.
- Guého E, Midgley G, Guillot J (1996). The genus *Malassezia* with description of four new species. *Antonie Van Leeuwenhoek*, 69 (6): 337-355.
- Nell A, James SA, Bond CJ, Hunt B, Herrtage ME (2002). Identification and distribution of a novel *Malassezia* species yeast on normal equine skin. *Vet Rec*, 150: 395-98.
- Sugita T, Takashima M, Shinoda T, Suto H, Uno T, Tsuboi R, Ogawa H, Nishikawa A (2002). New yeast species, *Malassezia dermatis*, Isolated from patients with atopic dermatitis. *J Clin Microbiol*, 40 (4): 1363-67.
- Sugita T, Takashima M, Kodama M, Tsuboi R, Nishikawa A (2003). Description of a new yeast species, *Malassezia ponica*, and its detection in patients with atopic dermatitis and healthy subjects. *Clin Microbiol*, 41(10): 4695-4699.
- Sugita T, Tajima M, Takashima M, Amaya M, Saito M, Tsuboi R, Nishikawa A (2004). A new yeast, *Malassezia yamatoensis*, isolated from a patient with seborrheic dermatitis, and its distribution in patients and healthy subjects. *Microbiol Immunol*, 48 (8): 579-83.
- Hirai A, Kano R, Makimura K, Duarte ER, Hamdan JS, Lachance MA, Yamaguchi H, Hasegawa A (2004). *Malassezianana* sp. nov., a novel lipid-dependent yeast species isolated from animals. *Int J Syst Evol Microbiol*, 54 (Pt2): 623-27.
- Sugita T, Tajima M, Ito T, Saito M, Tsuboi R, Nishikawa A (2005). Antifungal activities of tacrolimus and azole agents against the eleven currently accepted *Malassezia* species. *J Clin Microbiol*, 43: 2824-2829.
- Makimura K, Tamura Y, Kudo M, Uchida K, Saito H, Yamaguchi H (2000). Species identification and strain typing of *Malassezia* species stock strains and clinical isolates based on the DNA sequences of nuclear ribosomal internal transcribed spacer 1 regions. *J Med Microbiol*, 49 (1): 29-35.
- Yamada Y, Makimura K, Mirhendi H, Ueda K, Nishivama Y, Yamaguchi H, Osumi M (2002). Comparison of different methods for extraction of mitochondrial DNA from human pathogenic yeasts. *Jpn J Infect Dis*, 55 (4): 122-125.
- Mirhendi H, Makimura K, Zomorodian K, Yamada T, Sugita T, Yamaguchi H (2004). A simple PCR-RFLP method for identification and differentiation of 11 *Malassezia* species. *Microbiol Meth*, 61: 281-4.
- Senczek D, Siesenop U, Bohm K H (1999). Characterization of *Malassezia* species by means of phenotypic characteristics and detection of electrophoretic karyotypes by pulsed-field gelelectrophoresis (PFGE). *Mycoses*, 42 (5-6): 409-414.
- Makimura K, Tamura Y, Kudo M, Uchida K, Saito H, Yamaguchi H (2000). Species identification and strain typing of *Malassezia* species stock strains and clinical isolates based on the DNA sequences of nuclear ribosomal internal transcribed spacer 1 regions. *J Med Microbiol*, 49 (1): 29-35.
- Theelen B, Silvestri M, Gueho E, Van Belkum A, Boekhout T (2001). Identification and typing of *Malassezia* yeasts using amplified fragment length polymorphism (AFLP), random amplified polymorphic DNA (RAPD) and denaturing gradient gel electrophoresis (DGGE). *FEMS Yeast Res*, 1 (2): 79-86.
- Gupta AK, Kohli Y, Summerbell RC (2000). Molecular differentiation of seven *Malassezia* species. *J Clin Microbiol*, 38: 1869-1875.

16. Gaitanis G, Velegraki A, Frangoulis E, Mitrousia A, Tsigonia A, Tzimogianni A, Katsambas A, Legakis NJ (2002). Identification of *Malassezia* species from patient skin scales by PCR-RFLP. *Clin Microbiol Infect*, 8 (3): 162-173.
17. Shokohi T, Afshar P, Barzgar A (2009). Distribution of *Malassezia* species in patients with pityriasis versicolor in Northern Iran. *Indian J Med Microbiol*, 27: 321-24.
18. Tarazooie B, Kordbacheh P, Zaini F and et al (2004). Study of the distribution of *Malassezia* species in patients with pityriasis versicolor and healthy individual in Tehran, Iran. *BMC Dermatol*, 4(1):5.
19. Gupta AK, Kohli Y, Faergemann J, Summerbell RC (2001). Epidemiology of *Malassezia* yeasts associated with pityriasisversicolor in Ontario. *Canada Med Mycol*, 39 (2):199-206.
20. CrespoErchiga V, Ojeda Martos A, Vera Casano A (1999). Isolation and identification of *Malassezia* spp. In pityriasis versicolor, Seborrheic dermatitis and healthy skin. *Rev IberoamMicol*, 16: S16-S21.
21. Saleh SB, Makni F, Marrakchi S, Sellami H, Cheikhrouhou F, Bouassida S, Zahaf A, Ayadi A (2005). Identification of *Malassezia* species from Tunisian patients with pityriasisversicolor and normal subjects. *Mycoses*, 48 (4): 242-45.
22. Crespo Erchiga V, Ojeda MartosA, Vera CasanoA, Crespo Erchiga A, Sanchez FF (2000). *Malassezia globosa* as the causative agent of pityriasis versicolor. *Br J Dermatol*, 143: 799-803.
23. Nakabayashi A, Sei Y, Guillot J (2000). Identification of *Malassezia* species isolated from patients with seborrheic dermatitis, atopic dermatitis, pityriasis versicolor and normal subjects. *Med Mycol*, 38: 337-41.
24. Gaitanis G, Velegraki A, Alexopoulos EC, Chaspai V, Tsigonia A, Katsahbas A (2006). Distribution of *Malassezias* species in pityriasis versicolor and seborrheic dermatitis in Greece. Typing of the major pityriasisversicolor isolate *M. globosa*. *Br J Dermatol*, 154: 854-59.
25. Crespo Erchiga V, Ojeda Martos A, Vera Casario A et al. (1999). Mycology of pityriasisversicolor. *J Mycol Med*, 9: 143-48.
26. Gupta AK, Kohli Y, Summerbell RC et al. (2001). Quantitative culture of *Malassezias* species from different body sites of individuals with or without dermatoses. *Med Mycol*, 39: 243-51.
27. Arzumaniyan VG (2001). The yeast *Malassezia* on the skin of healthy individuals and patients with atopic dermatitis. *Vestn Ross Akad Med Nauk*, 2: 29-31.