Original Article



Environmental Risk Assessment for Veterinary Antibiotics and

*Yu Bin HO¹, Mohamad Pauzi ZAKARIA², Puziah Abdul LATIF², Nazamid SAARI³

Hormone in Malaysian Agricultural Soil

1. Dept. of Environmental and Occupational Health, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, Selangor, Malaysia

2. Faculty of Environmental Studies, Universiti Putra Malaysia, Selangor, Malaysia

3. Faculty of Food Science and Technology, Universiti Putra Malaysia, Selangor Malaysia

*Corresponding Author: Email: yubin@upm.edu.my

(Received 20 July 2014; accepted 10 Sep 2014)

Abstract

Iranian J Publ Health, Vol. 43, Suppl. No.3, Oct 2014, pp. 67-71

Background: Repeated applications of animal manure as fertilizer are normal agricultural practices in Malaysia that may release veterinary antibiotics to environment from treated animals.

Methods: Environmental risk assessment (ERA) on 5 commonly used antibiotics and one hormone in Malaysian broiler farm was calculated using the ratio of measured environmental concentration (MEC) and predicted no effect concentration (PNEC) in the environment. PNEC was derived from the available acute and chronic toxicity data in the open peer-reviewed literature. Risk quotients (RQ) were then calculated for 5 antibiotics (erythromycin, norfloxacin, sulfadiazine, trimethoprim, and tylosin) and one hormone (progesterone).

Results: RQ for tylosin has exceeded 1, indicating that this compound has high risk of acute toxicity in Malaysian agricultural soil while trimethoprim and tylosin possessed medium risk of chronic toxicity. The rest of the compounds showed low risk or no risk in acute or chronic ecological toxicity.

Conclusion: The release of tylosin, trimethoprim, norfloxacin and progesterone from broiler manure to agricultural soil may potentially harm the environment.

Keywords: Veterinary antibiotic, Environmental risk assessment, Risk quotient, Malaysia

Introduction

Livestock producers often misused veterinary pharmaceuticals as supplement in animal feed to increase weight gain and prevent disease among their livestock. These chemicals are introduced increasingly without realising the consequences for the environment, direct and indirect effects for human health.

Antibiotics are compounds that present naturally in the environment. However, the frequent use of antibiotics either by human to treat diseases or as animal feed supplements has raised concerns about the increased adaptation of antibiotics resistant bacteria (1) and bioaccumulation of veterinary pharmaceuticals in plants (2-7).

Among the endocrine disrupting compounds (EDCs), steroid hormones are of particular concern because of their capacity to induce strong endocrine responses. Steroid hormones can exist as either natural or synthetic substances, mimicking the effects of endogenous estrogens as EDCs through binding to specific receptors common to non-target organisms (invertebrates, fish, reptiles, birds and mammals) (8).

Therefore, the investigation of pharmaceutical pollution in broiler manure has special importance because they constitute a major source of hazardous unmetabolized veterinary pharmaceutical to the environment via fertilization of agricultural soil. An environmental risk assessment is conducted based on the study reported by Ho et. al (9) in order to assess the potential risk of 6 compounds (erythromycin, norfloxacin, sulfadiazine, trimethoprim, tylosin and progesterone) in agricultural soil by evaluating the ratio between the measured environmental concentration (MEC) and the predicted no-effect concentration (PNEC).

Materials and Methods

Soil samples were collected from 10 respective agricultural fields in Selangor, Negeri Sembilan and Melaka in Malaysia according to Ho et. al (9). The target antibiotics and hormone in soil samples were ultrasonic extracted using the optimized extraction buffer (MeOH: ACN: 0.1M EDTA: McIlvaine buffer (pH 4), 30:20:25:25). The extracts were then cleaned up using Oasis HLB 3cc/60mg cartridge and analyzed in liquid chromatography tandem mass spectrometry (LC-MS/MS).

The risk of quotient (RQ) is a useful tool to characterize potential ecological risk of many contaminants in the environment (10-12). RQ was calculated as the ratio between maxiumum measured environmental concentration (MEC) and predicted no effect concentration (PNEC) (Eq. 1). Maximum MEC was obtained from the occurrence data whereas PNEC was estimated by dividing the lowest values of acute EC_{50} or LC_{50} or the chronic no observed effect concentration (NOEC) with a default assessment factor (AF). All values below the method quantification limit (MQL) were set to 0. Up until now, there are limited toxicological data of antibiotics and hormones in terrestrial organisms have been reported. Therefore, $PNEC_{soil}$ values were estimated from $PNEC_{water}$ values by applying the equilibrium partition approach as suggested by Martin et al. (13) (Eq. 2).

Risk Quotient =
$$\frac{MEC_{soil}}{PNEC_{soil}}$$
 Eq. (1)
PNEC_{soil} = PNEC_{water} × K_d Eq. (2)
PNEC_{acute} = $\frac{EC_{so}orLC_{50}}{1000}$ Eq. (3)
PNEC_{chronic} = $\frac{NOEC}{AF}$ Eq. (4)

According to the Technical Guidance Document on Risk Assessment of the European Commission 2003, when only short-term/acute toxicity data EC_{50}/LC_{50} are available, the calculation of PNECwater is obtained from EC50/LC50 divided by an assessment factor (AF) of 1000. The availability of long-term/chronic NOEC values for one, two or three trophic levels are depending on assessment factor (AF) of 100, 50 and 10 respectively (European Commission, 2003). For risk characterization, a commonly used risk-ranking criterion was applied. When the RQ equals or exceeds 1, an ecological "high risk" is suspected. "Low risk" is suspected when 0.01 < RQ < 0.1, whereas, 0.1 < RQ < 1 indicates "medium risk" (11, 14)

Results

Risk quotients (RQ) and the corresponding levels of potential ecological risks to the terrestrial environment were calculated as shown in Table 1 and Table 2 respectively. Acute and chronic environmental risks of each compound were calculated based on the availability of PNEC and K_d data from the literature.

The acute environmental risks of 5 antibiotics are shown in Table 1. The results showed that tylosin possessed high acute ecological risk after onemonth application of the broiler manure to agricultural soil, while trimethoprim is estimated to have medium acute risk in the environment. Norfloxacin has shown low acute ecological risk while erythromycin and sulfadiazine are estimated to have no acute ecological risk in the terrestrial environment.

The chronic environmental risks of 3 antibiotics and one hormone are summarized in Table 2. Two antibiotics, trimethoprim, and tylosin were estimated to have medium chronic risk in terrestrial environment. The results showed that norfloxacin and progesterone possessed low chronic risk in the terrestrial environment after manure amendment.

Table 1: Estimated	risk quotients and	corresponding lev	els of potential	acute ecological risks to	the terrestrial envi-
	1	1 0	1	0	

forment									
	Lowest EC ₅₀ or LC ₅₀ (ng/L)	AF	PNEC _{water} (ng/L)	K _d (L/kg)	PNEC _{soil}	MEC ^a (ng/kg)	RQ (MEC/PNEC)	ERA	
Erythromycin	3.1×10 ^{7b}	1000	3.1×10 ⁴	164.76	5.0×10 ⁶	0	0	No	
Norfloxacin	7.0×10 ^{7b}	1000	7.0×10^{4}	7943	5.5×10^{8}	9.6×10 ⁴	1.7×10-4	Low	
Sulfadiazine	2.2×10 ^{6c}	1000	2.2×10^{3}	2.8	6.2×10^{3}	0	0	No	
Trimethoprim	1.6×10 ^{7b}	1000	1.6×10^{4}	25.7	4.1×10 ⁵	6.0×10 ⁴	0.15	Medium	
Tylosin	1.4×10 ^{6b}	1000	1.3×10^{3}	128	1.8×10^{5}	6.8×10 ⁵	3.84	High	
^a (9), ^b (15), ^c (11)									

Table 2: Estimated risk quotients and corresponding levels of potential chronic ecological risks to the ter-

restrial environment.								
	Lowest NOEC (ng/L)	AF	PNEC _{water} (ng/L)	K _d (L/kg)	PNEC _{soil}	MECª (ng/kg)	RQ (MEC/PNEC)	ERA
Norfloxacin	1.7E×10 ^{5b}	100	1.7×10^{3}	7943	1.3×10 ⁷	9.6×10 ⁴	7.1×10-3	Low
Trime- thoprim	1.0×10 ^{6b}	100	1.0×10^{4}	25.7	2.6×10 ⁵	6.0×10^{4}	0.23	Medium
Tylosin	3.0×10 ^{5b}	50	6.0×10^{3}	128	7.7×10^{5}	6.8×10^{5}	0.88	Medium
Progester- one	1.0×10 ^{5d}	50	2.0×10^{3}	204	4.1×10 ⁵	2.4×10 ⁴	0.059	Low

^a(9), ^b(15), ^d(16)

Discussion

The results of the ERA performed in this study indicate that the release of antibiotic and hormone residues after soil amendment is posing certain ecological risks to soil environment. In this study, only the maximum MEC were considered for the calculation of RQ in the worst-case scenario. However, there are some limitations to assess the environmental risk of all target compounds reported by Ho et al. (9); because we are not able to collect all the toxicity data and the solid-water partition coefficient values in the literature. Moreover, the toxicity of the mixture of compounds was not considered. The potential environmental risk of each compound was characterized base on three risk levels: 0.01-0.1, low risk; 0.1-1, medium risk; and >1, high risk.

Previous studies are focussed in aquatic toxicity of pharmaceuticals; ecotoxicity of pharmaceuticals in soil environment was not well documented. However, RQ of trimethoprim in soil was previously reported by Martin et al. (13), the results of the study showed that trimethoprim has no risk in digested sludge amended soil and compost amended soil. Other compounds, such as erythromycin, tylosin and norfloxacin showed low risk, moderate risk and low risk respectively in aquatic environment (15).

Conclusion

Veterinary antibiotics and hormone were detected in the broiler manure and the manure was subsequently applied onto agricultural soil as fertilizer. The MEC of these compounds in manure amended agricultural soil revealed that the application of broiler manure to agricultural soil may potentially pollute the environment and exerts considerable ecological risk in the soil environment. The results of this study showed that tylosin has high acute risk in Malaysian agricultural soil while trimethoprim and tylosin exerted medium chronic risk in the soil environment. The environmental risk of pharmaceuticals in agricultural soil is very much depending on the farm management and the sorption affinity of the compounds. The MEC in manure and soil is highly depending on the amount of antibiotics administered to the animals by the farmers. Therefore, it is advised to treat or compost the animal manure before applying on agricultural soil to reduce the ecological risk.

Ethical considerations

Ethical issues (Including plagiarism, Informed Consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc) have been completely observed by the authors.

Acknowledgement

This study was financially supported by Universiti Putra Malaysia Grant (project number 9423900). The authors declare that there is no conflict of interest.

References

1. Sarmah AK, Meyer MT,Boxall ABA (2006). A global perspective on the use, sales, exposure

pathways, occurrence, fate and effects of veterinary antibiotics (VAs) in the environment. *Chemosphere*, 65:725-759.

- Kumar K, Gupta SC, Baidoo SK, Chander Y,Rosen CJ (2005). Antibiotic Uptake by Plants from Soil Fertilized with Animal Manure. J Environ Qual, 2082-2085.
- Boxall ABA, Johnson P, Smith EJ, Sinclair CJ, Stutt E,Levy LS (2006). Uptake of Veterinary Medicines from Soils into Plants. J Agr Food Chem, 54:2288-2297.
- Kong WD, Zhu YG, Liang YC, Zhang J, Smith FA,Yang M (2007). Uptake of oxytetracycline and its phytotoxicity to alfalfa (Medicago sativa L.). *Environ Pollut*, 147:187-193.
- 5. Redshaw CH, Wootton VG, Rowland SJ (2008). Uptake of the pharmaceutical Fluoxetine Hydrochloride from growth medium by Brassicaceae. *Phytochemistry*, 69:2510-2516.
- 6. Herklotz PA, Gurung P, Vanden Heuvel B,Kinney CA (2010). Uptake of human pharmaceuticals by plants grown under hydroponic conditions. *Chemosphere*, 78:1416-1421.
- Eggen T, Asp TN, Grave K, Hormazabal V (2011). Uptake and translocation of metformin, ciprofloxacin and narasin in forage- and crop plants. *Chemosphere*, 85:26-33.
- Santos LHMLM, Araújo AN, Fachini A, Pena A, Delerue-Matos C,Montenegro MCBSM (2010). Ecotoxicological aspects related to the presence of pharmaceuticals in the aquatic environment. J Hazard Mater, 175:45-95.
- 9. Ho YB, Zakaria MP, Latif PA, Saari N (2012). Simultaneous determination of veterinary antibiotics and hormone in broiler manure, soil and manure compost by liquid chromatography-tandem mass spectrometry. J Chromatogr A, 1262:160-168.
- Gonzalez-Pleiter M, Gonzalo S, Rodea-Palomares I, Leganes F, Rosal R, Boltes K, Marco E,Fernandez-Pinas F (2013). Toxicity of five antibiotics and their mixtures towards photosynthetic aquatic organisms: Implications for environmental risk assessment. *Water Res*, 47:2050-2064.
- Zhang R, Tang J, Li J, Zheng Q, Liu D, Chen Y, Zou Y, Chen X, Luo C,Zhang G (2013). Antibiotics in the offshore waters of the Bohai Sea and the Yellow Sea in China: Occurrence, distribution and ecological risks. *Environ Pollut*, 174:71-77.

www.<u>SID.ir</u>

- Xu W, Yan W, Li X, Zou Y, Chen X, Huang W, Miao L, Zhang R, Zhang G,Zou S (2013). Antibiotics in riverine runoff of the Pearl River Delta and Pearl River Estuary, China: Concentrations, mass loading and ecological risks. *Environ Pollut*, 182:402-407.
- Martin J, Camacho-Munoz MD, Santos JL, Aparicio I,Alonso E (2012). Distribution and temporal evolution of pharmaceutically active compounds alongside sewage sludge treatment. Risk assessment of sludge application onto soils. J Environ Manage, 102:18-25.
- Kosma CI, Lambropoulou DA, Albanis TA (2013). Investigation of PPCPs in wastewater treatment plants in Greece: Occurrence, removal and environmental risk assessment. *Sai Total Emiron*, 466-67:421-438.
- 15. Turkdogan FI, Yetilmezsoy K (2009). Appraisal of potential environmental risks associated with human antibiotic consumption in Turkey. *J Hazard Mater*, 166:297-308.
- Orias F, Perrodin Y (2013). Characterisation of the ecotoxicity of hospital effluents: A review. *Sci Total Emiron*, 454-455:250-276.