**Original Article** 



Iran J Public Health, Vol. 44, No.1, Jan 2015, pp.119-129

# The Social Determinants of Health in Association with Women's Health Status of Reproductive Age: A Population-Based Study

# Azam BAHEIRAEI<sup>1</sup>, \*Fatemeh BAKOUEI<sup>2</sup>, Eesa MOHAMMADI<sup>3</sup>, Ali MONTAZERI<sup>4</sup>, Mostafa HOSSENI<sup>5</sup>

1. Community Based Participatory Research Center, Iranian Institute for Reduction of High-Risk Behaviors, Tehran University of Medical Sciences, Tehran, Iran

2. School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran

3. Dept. of Nursing, Tarbiat Modares University, Tehran, Iran

4. Mental Health Research Group, Health Metrics Research Center, Iranian Institute for Health Sciences Research, ACECR,

Tehran, Iran

5. Dept. of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

\*Corresponding Author: Email: bakouei2004@yahoo.com

(Received 15 Oct 2014; accepted 06 Dec 2014)

#### Abstract

**Background:** Prioritizing women's health helps achievement of the 4<sup>th</sup> and 5<sup>th</sup> goals of Millennium Development Program. This study aimed to investigate association between social determinants of health and women's health of reproductive age.

**Methods**: This population-based cross-sectional study, using multi-stage sampling procedure was conducted on 770, 15 to 49-year-old women residing in any one of the 22 municipality zones across Tehran, Iran. Eligible women were interviewed at home with SF-36 (Short Form Health Survey) and socio-demographic questionnaires. Social determinants of health contains; ethnicity, education, job, income, and crowding index. Data were analyzed by ANOVA and Multiple Linear Regression using SPSS-16. The threshold of P.V was considered 0.05.

**Results:** Overall, 770 women with mean age  $33.9\pm9.3$  years were interviewed. Majority of them were married (72.27%), housewives (62.2%), of Persian ethnicity (64.3%), and educated to high school level (43.8%). Age with dimensions of health except role emotional, mental health, and social functioning had significant association with B from -0.65 to -0.16.educational level with dimensions of health except role emotional and social functioning had significant association with B from 3.61 to 6.43 (P<0.05). Income with dimensions of health except role physical had significant association with B from -9.97 to -4.42.

**Conclusion:** Reflection of unfavorable economic conditions and low education level on negative women' health experiences are alarming. Interaction between social determinants of health and health status must be considered in policymaking, and there is a need for policies that would enhance health of women in the low education and income brackets.

Keywords: Women's health status, Social determinants of health, Reproductive age

### Introduction

Several factors are considered as determinants of health and complex interactions among these may have a profound impact on health (1). Prioritizing women's health helps achievement of the 4<sup>th</sup> and

5<sup>th</sup> goals of Millennium Development Program (2). Health inequalities refer to differences in health status or health outcomes associated with such factors as gender, race, ethnicity, and socioeconomic status (3). In addition to association of health inequalities with genetic and biological variables, social structure is also another health-related factor (4). Men and women are exposed to determinants of health in different ways (5). In 2008, the World Health Organization's special commission emphasized the importance of research to address Social Determinants of Health (SDH) to identify and determine health inequalities according to geographical regions and groups of people (6), as progress in eliminating health inequalities in the absence of good data will be impossible (7).

SDH contain socioeconomic factors that reflect, "the conditions in which people are born, grow up, live, work, and age" (8). Evidence on the relationship between SDH and health outcomes, the prevalence of diseases and health inequalities are necessary (9, 10). Their impact on health is complicated, ranging from biology to behavior. Health inequalities usually have reverse relationship with social determinants of health, which means people with lower socioeconomic status have poor health (11, 12). Furthermore, people at lower education or income brackets, the unemployed people, and manual workers are more likely to have poor health (13, 14), yet this obvious pattern is widely varied in different populations (15). In developing countries within the Middle East like Iran, with different cultures from that of western societies, research into social health patterns, especially women's health is in its early stage (16, 17).

As a developing country, Iran is rapidly moving toward urbanization and industrialization. A higher percentage of 71% of Iran's population in 2006(71%), compared to 61.3% in 1996, and 54.3% in 1986; live in the cities (18). It is expected that socioeconomic changes at society level will affect family and household levels. Determining current condition of health status provides a basis for a series of health planning processes, including prioritizing plans, setting goals and strategy, and implementing plans (19). According to the statistics published by the Iran's Statistics Center in 2011, the highest percentage of all female population (about 60%) are women of reproductive age, and approximately 16.2% of these women live in Tehran (20). Given that women's health in this period affects long-term health of theirs and their family members(21), and that a large proportion of women's population in this country are women of reproductive age, it is necessary to study women's health status.

This study aimed to investigate the relationship between social determinants of health and women's health status of reproductive age. Understanding social determinants of health related to changes in health is important for policy making to reduce health inequalities.

# Material & Methods

### Study design and sampling method

This population-based cross-sectional survey has been conducted in Tehran during February to September 2013. This study is a part of a mixed method study using a sequential explanatory design, the protocol of which has been previously published. The sample size in the current study was 770 people as explained in the protocol (22).Participants were 770 women of reproductive age (15- to 49-year-old), residing in any one of the 22 municipality zones across Tehran.

These women were selected through a multi-stage sampling procedure. Tehran is divided into 22 zones and each zone is divided to a number of blocks with unequal population. For our trained interviewers it was only possible to question 10 eligible women from each block. Therefore, 77 blocks were surveyed in our study. The number of blocks in each zone was proportion to size of study population in the corresponding zone. In each zone the required number of blocks was randomly selected from the list of blocks. In each city block, the total number of households was divided by 10 to find the distance between household in the list. The eligible women were chosen via systematically random sampling to interview. In case a woman declined to take part, the family next door was invited to participate in the study and complete the questionnaire. Considering random sampling from all 22-city zones in this study, there is no selection bias and the internal validity is increased.

Eligibility criteria in this study were women in age group of 15-49 years and no any known diseases such as severe mental retardation and mental disorders (making them unable to respond to the questionnaire).

### Study Scales

**Dependent variable**: Health status of women of reproductive age

The SF-36 questionnaire (Short Form Health Survey) is a general scale of health status. It is a tool for assessment of health of the general population, and can be used to compare health status over time, place, and people (23). This scale was first designed by Ware et al. to evaluate health of the community, determine health policies, and assess the efficiency of the designed treatment. It can be completed as self-administered, or with help of an interviewer in person or by telephone. SF-36 questionnaire consists of 8 health-related concepts/domains: physical functioning with 10 items, role limitations due to physical problems with 4 items, bodily pain with 2 items, general health with 5 items, vitality with 4 items, social functioning with 2 items, role limitations due to emotional problems with 3 items, and mental health with 5 items. In addition, there is an item in this questionnaire that indicates perceived change in general health status by the person over last one-year period (24).

SF-36 questionnaire was translated in Iran using forward-backward translation method, and it has also been culturally adapted and its validity and reliability determined. Reliability was found using internal consistency, validity was determined using known groups comparison, and convergent validity was measured. Cronbach's coefficient of all dimensions of the questionnaire, except vitality (0.65), ranged from 0.77 to 0.90. Mean score for the 8 indicators of this test based on the protocol of the questionnaire ranged from 0 (worst status) to 100 (best status) (25). According to different Likert scale ranging from 2 to 6 items, at first all questions were weighted and changed to range from 0 to 100, with a higher score indicating higher health status. Scores for each dimension of SF-36 scale were calculated similar the instructions provided in the SF-36 manual (26). The researcher had obtained prior permission for using this questionnaire.

# Independent variables: Social Determinants of Health

Independent variables included social determinants of health (ethnicity, education level, occupation status, sufficiency of income for expenses, and crowding index) and other variables (age, marital status, and family size) that were assessed by socio-demographic questionnaire designed and approved by the researcher and experts. Crowding index was determined by dividing the number of family members by the number of rooms, not considering the bathroom and kitchen: low (less than1), average (between 1-2), and high (more than 2).

# Data collection

All participants were interviewed face-to-face in their own house by the team of interviewers. Interviewers were trained by the research team for conducting standard interviews. To ensure reliability of the collected data, interviewers were closely controlled and monitored. To this end, the following control methods were implemented; simultaneous control of interviewers, control after performing their tasks, review of the questionnaires, and statistical control of the questionnaires. The women were interviewed alone to minimize potential response bias. Study protocol was approved by the ethics committee of Tehran University of Medical Sciences. Following a detailed explanation, informed consent was obtained from all participating women.

### Data analysis

Data were analyzed using SPSS-16. The percentage of missing in variables used in the analysis was negligible and varied only from 0.2% to 1.7%; therefore ignored in the analysis. At first, the independent and dependent variables were assessed descriptively with frequency and percentage.

To investigate the relationship between dimensions of health and social determinants of health and other demographic variables, analysis of variance (ANOVA) as the univariate analysis was used and then multiple linear regression analysis as the adjusted one were used to analyze associations between variables. On the other hand, to avoid effects of the confounding factors, and to predict effects of independent variables on dependent ones (dimensions of health), variables with significant associations in the ANOVA test were entered into the Stepwise Multiple Linear Regression. Authors have conducted 8 model of linear regression as in each of model, one of the dimensions of health considered as dependent variable and all independents variables with significant association in ANOVA were entered model. The threshold of type one error was considered 0.05.

### Results

#### Descriptive variables

In total, 770, 15- to 49-year-old women of reproductive age with mean age  $33.9\pm9.3$  years were interviewed. Majority of them were married (72.27%) housewives (62.2%), of Persian ethnicity (64.3%), and educated to high school level (43.8%) (Table 1).

Table 2 presents mean score and standard deviation obtained in all dimensions of health in women of reproductive age. The highest mean score related to the physical functioning and the lowest mean score to vitality dimensions. Data on the item of perceived change in the health status by the person over one-year period revealed 40.2% and 29.5% of women had similar health status to the previous year's and slightly worse than the previous year's, respectively.

# Health in association with demographic and socioeconomic variables

The relationship between health status of women of reproductive age and independent variables through one-way analysis of variance (ANOVA) is also has been shown in Table 2. According to the results, there was a significant association between some dimensions of health and variables; age, marital status, education level, occupation status, and sufficiency of income for expenses. To avoid effects of the confounding factors of the mentioned variables, these variables were entered into the stepwise multivariable linear regression model. Table3 presents adjusted association of the mentioned independent variables with health dimensions as dependent variables. The results of linear regression analysis in the models 1 to 8 are shown in Table 3.

# **Table 1:** The characteristics of the study sample (n=770)

Variables	n (%)				
Age groups (yr)					
15-25	165 (21.5)				
26–35	266 (34.5)				
36–45	224 (29.1)				
46-49	115 (14.9)				
Marital status					
Married	556 (72.2)				
Divorced/Widowed	31 (4)				
Single	183 (23.8)				
Ethnicity					
Fars	487 (64.3)				
Azari	179 (23.2)				
Others	91 (13.5)				
Educational level (yr)					
Illiterate/ Primary (years 1-5)	50 (6.5)				
Intermediate (years 6-8)	83 (10.8)				
High school (years 9–12)	337 (43.8)				
University >12	300 (38.9)				
Occupation status					
Housewife	479 (62.2)				
Employed	190 (24.7)				
Student	101 (13.1)				
Sufficiency of income for expenses					
Absolutely not	337 (44.1)				
To some extent	240 (31.4)				
Completely	188 (24.5)				
Crowding index					
Low	215 (27.5)				
Average	385 (51.6)				
High	168 (20.9)				
Family Size					
Less than four	321 (42.3)				
Four to five	384 (50.7)				
More than five	53 (7.0)				

#### Iran J Public Health, Vol. 44, No.1, Jan 2015, pp.119-129

SF-36 subscales: mean (SD)								
	General	Physical	Role	Role	Body Pain	Vitality	Mental Health	Social
T 1 1 . T7 · 11	Health	Functioning	Physical	Emotional				Functioning
Independent Variables								
Age groups (yr)								
15–25	67.68 (19.26)	86.71 (18.62)	75.00 (33.56)	60.36 (41.21)	67.25 (23.76)	64.52 (19.58)	65.65 (21.18)	74.37 (23.99)
26-35	65.07 (20.23)	84.84 (18.52)	70.72 (36.78)	65.01 (40.55)	66.56 (23.06)	60.50 (21.28)	62.38 (20.12)	74.41 (24.77)
36–45	58.60 (19.86)	78.13 (20.17)	62.27 (39.77)	60.81 (41.95)	57.63 (25.30)	60.04 (20.53)	62.66 (22.07)	73.32 (23.27)
46-49	55.75 (18.61)	68.02 (26.37)	54.13 (41.34)	57.10 (43.20)	58.97 (26.24)	56.65 (18.65)	62.45 (19.51)	69.09 (24.28)
ANOVA p	.000	.000	.000	.342	.000	.016	.414	.239
Marital status								
Married	61.13 (19.46)	79.60 (21.195)	64.67 (38.77)	62.74 (41.33)	61.71 (24.78)	59.75 (20.17)	63.16 (20.79)	73.00 (23.94)
Divorced/ Widowed	52.74 (21.86)	74.33 (25.24)	59.67 (39.09)	55.91 (43.36)	55.00 (26.09)	54.83 (21.92)	52.27 (23.12)	65.94 (28.91)
Single	67.73 (20.69)	85.58 (20.04)	74.02 (35.97)	59.07 (41.81)	68.30 (23.39)	64.39 (20.52)	64.99 (20.15)	75.42 (23.68)
ANOVA p	.000	.001	.010	.436	.002	.009	.010	.126
Ethnicity								
Fars	62.69 (19.12)	80.37 (21.56)	65.56 (39.04)	61.74 (41.95)	63.62 (24.41)	61.28 (19.42)	63.66 (19.50)	73.93 (24.03)
Azari	60.28 (22.32)	81.29 (21.14)	67.23 (37.59)	57.86 (41.21)	61.51 (25.45)	59.24 (21.79)	61.96 (22.62)	71.53 (25.04)
Others	64.35 (20.67)	82.24 (20.16)	70.91 (36.22)	67.30 (39.68)	62.77 (24.78)	59.95 (22.56)	62.90 (23.67)	73.40 (22.97)
ANOVA p	.225	.696	.424	.182	.628	.501	.648	.534
Educational level (yr)								
Illiterate/ Primary (years 1-5)	51.45 (14.98)	67.81 (22.94)	55.10 (41.13)	58.66 (41.81)	52.82 (25.61)	57.18 (19.72)	59.02 (21.36)	69.94 (21.75)
Intermediate (years 6-8)	57.46 (20.20)	79.35 (22.13)	62.04 (41.20)	59.03 (43.34)	59.50 (26.26)	56.82 (21.03)	60.97 (21.44)	70.03 (22.84)
High school (years 9–12)	60.42 (20.87)	79.27 (22.28)	64.56 (37.94)	60.46 (41.89)	61.60 (24.89)	59.20 (21.18)	61.58 (21.83)	72.57 (25.72)
University >12	67.72 (18.55)	85.13 (18.31)	72.23 (36.79)	64.08 (40.60)	67.18 (23.15)	63.87 (19.10)	66.29 (19.05)	75.60 (22.83)
ANOVA p	.000	.000	.005	.598	.000	.004	.010	.147
Occupation status								
Housewife	60.50 (20.14)	79.38 (21.63)	65.55 (38.25)	62.17 (41.46)	61.78 (24.42)	59.40 (20.13)	62.20 (21.02)	73.27 (23.77)
Employed	63.26 (19.21)	80.91 (20.64)	65.00 (39.32)	61.87 (41.15)	63.77 (24.92)	61.35 (20.35)	64.21 (19.59)	73.16 (24.84)
Student	69.49 (20.20)	87.27 (19.68)	75.25 (35.98)	58.33 (42.73)	67.60 (25.20)	65.15 (21.50)	65.81 (22.19)	73.71 (24.63)
ANOVA p	.000	.004	.057	.699	.098	.037	.225	.983

### Table 2: Respondents' characteristics in association with the mean scores of health dimensions through ANOVA

Table 2: Cond...

SF-36 subscales: mean (SD)								
	General Health	Physical Functioning	Role Physical	Role Emotional	Body Pain	Vitality	Mental Health	Social Functioning
Independent Variables			-					-
Sufficiency of income for								
expenses								
Absolutely not	58.22 (20.46)	77.29 (23.04)	62.53 (39.07)	55.98 (42.08)	56.99 (25.38)	56.55 (20.62)	59.65 (21.68)	68.88 (25.62)
To some extent	64.39 (18.46)	82.71 (19.63)	69.43 (37.56)	65.11 (40.38)	67.23 (23.83)	63.11 (19.39)	64.14 (19.60)	75.91 (22.29)
Completely	67.01 (20.24)	84.45 (19.17)	70.18 (37.70)	67.02 (40.89)	68.46 (22.20)	64.91 (20.23)	68.06 (19.91)	77.93 (22.09)
ANOVA p	.000	.000	.036	.004	.000	.000	.000	.000
Crowding index								
Low	64.76 (19.92)	83.40 (20.96)	67.26 (39.56)	60.44 (43.25)	63.87 (24.02)	59.60 (20.88)	61.85 (21.40)	75.26 (23.96)
Averag	62.17 (20.53)	80.98 (20.36)	67.14 (37.30)	61.97 (41.13)	63.30 (25.28)	61.16 (21.44)	63.66 (21.48)	73.24 (24.70)
High	61.79 (20.11)	78.27 (22.70)	65.87 (38.70)	65.54 (39.72)	62.30 (23.54)	60.73 (19.00)	64.88 (19.91)	72.79 (23.02)
ANOVA p	.291	.096	.933	.519	.848	.705	.415	.575
Family Size								
Less than four	63.93 (20.11)	82.77 (20.17)	66.40 (38.08	61.21 (41.33)	63.61 (23.87)	59.69 (20.48)	63.05 (20.72)	73.27 (24.17)
Four to five	61.00 (19.93)	78.88 (22.27)	68.00 (38.36)	62.54 (41.77)	62.71 (25.26)	61.02 (20.97)	63.21 (21.48)	73.61 (23.96)
More than five	64.61 (22.04)	83.57 (20.20)	58.96 (41.04)	61.00 (41.20)	63.92 (25.54)	64.80 (15.93)	65.22 (17.87)	74.50 (24.36)
ANOVA p	.122	.051	.272	.904	.872	.230	.794	.940
Total score	62.36 (20.13	80.83 (21.26)	66.67 (38.33)	61.59 (41.52)	63.01 (24.69)	60.62 (20.43)	63.16 (20.84)	73.30 (24.12)

This Table shows only variables, which were remained as significant factors after adjusting other variables in regression analysis with dimensions of health. As can be seen from Table 3, significant associations were found between the followings dimensions; general health, physical functioning, role limitations due to physical problems, bodily pain, and vitality with age; mental health with marital status; all dimensions of health, except role limitations due to emotional problems and social functioning with women's education level; and all dimensions of health except role limitations due to physical problems with sufficiency of income for expenses.

T-1-1-2.T.	<b></b>	1 C +1		C	11-1-	1	1	L	1	
I able 5: Li	near regression	analysis for th	e prediction	n of variance in	neaith	dimensions	scores	DV SOCIO-	demographic	variables
			- F							

Variables	В	Confidence interval	Р
Model 1: Physical functioning			
Age	- 53	-0.690.37	0.000
Educational level	100		0.000
Illiterate/Primary (years 1-5)	Reference	-	
University	4.23	1.11 - 7.36	0.008
Sufficiency of income for expenses			
Completely	Reference	-	
Absolutely not	-4.42	-7.441.40	0.004
Model 2: Role physical		0.04 0.27	0.000
Age	65	-0.940.36	0.000
Educational level	D.C.		
Hiterate/Primary (years 1–5)	( 35 ( 027)	- 0.70 12.00	0.027
Model 3: Bodily pain	0.33 (.027)	0.70 - 12.00	0.027
Age	32	0.51 - 0.13	0.001
Educational level			
Illiterate/Primary (years 1-5)	Reference	-	
University	4.11	0.46 - 7.76	0.027
Sufficiency of income for expenses			
Completely	Reference		
Absolutely not	-9.50	-13.035.97	0.000
Model 4: General health			
Age	33	-0.480.18	0.000
Educational level			
Illiterate/Primary (years 1–5)	Reference	-	0.000
University	6.43	3.50 - 9.36	0.000
Completely	Reference		
Absolutely not	-5.65	-8 472 82	0.000
Model 5: Vitality			0.000
Age	16	0.04 0.32	0.041
Educational level			
Illiterate/Primary (years 1–5)	Reference	-	
University	3.61	0.55 - 6.67	0.021
Sufficiency of income for expenses			
Completely	Reference	-	
Absolutely not	-6.47	-9.423.52	0.000
Model 6: Social functioning			
Sufficiency of income for expenses	D.C		
Absolutely pot	7 01	- 11 37 4 45	0.000
Model 7: Bole emotional	-7.91	-11.57+.+5	0.000
Sufficiency of income for expenses			
Completely	Reference		
	0.07	15.00 4.04	0.001
Model 8: Mental health	-9.97	-13.894.04	0.001
Model 6. Mental health Marital status			
Married	Reference		
Divorced/Widowed	-9.02	-16.711.32	0.022
Educational level	2.02		0.022
Illiterate/Primary (years 1–5)	Reference	<u>-</u>	
University	4.08	0.99 - 7.16	0.010
Sufficiency of income for expenses			
Completely	Reference	-	
Absolutely not	-5.22	-8.262.18	0.001

### Discussion

In this study, the highest and the lowest mean scores were found in dimensions of physical functioning and vitality, respectively. In Iraqi immigrant women settled in Malavsia, the highest and the lowest scores were in physical functioning (73.5) and vitality (55.0) (27); on Arab-American women, the highest score was in physical functioning (83.75) and the lowest in role limitations due to emotional problems (46.94) (28); on U.S. women, the highest score was in physical functioning (84.2) and the lowest in vitality (60.9) (29). In line with the objectives of the study, dimensions of general health, physical functioning, and role limitations due to physical problems, bodily pain, and vitality significantly reduced with increasing age of the women. That is, mean score in these dimensions, with every one year's aging, reduced from 0.16 (vitality domain) to 0.65 (role limitation due to physical problems), which was in agreement with the findings of other studies (30, 31). This can be due to the active lifestyle of the younger age group compared to the older ones (31). In the present study, dimensions of social functioning, role limitations due to emotional problems, and mental health did not have a significant association with age; while in other studies a significant and positive association was observed between age and these dimensions, which is indicative of better mental status of older women (30, 32). Positive mental adjustment with aging could be due to re-assessment of values in women (30). In this study, single women were better in dimensions of general health, physical functioning, role limitations due to physical problems, bodily pain, vitality, and mental health than married women. However, these differences were insignificant in presence of other variables in regression analysis. Dimensions of vitality, social functioning, role limitations due to emotional problems, and mental health were significantly better in married women than in single women (27, 33). The impact of marare exposed to limited physical activity, unhealthy diets, and lack of attention to personal health (35). Married women in the Indian community, first play the role of the wife, the mother, and the caregiver in the family, and lastly attend to their own health needs (5). In the present study, divorced and widowed women had a significantly lower score in mental health compared to married women, which concurs with results of other studies (27, 30). The significant difference in mental health of married women may be due to the emotional support provided by the spouse (27).

In terms of educational level, all health dimensions except role limitations due to emotional problems and social functioning, were better in women with university education. Other studies also recognized university education as an influential factor in women's health (1, 36, 37). Education should be considered a central factor in explaining health inequalities. For a healthy lifestyle, education is a priority (37).

In respect to occupational status, student women were better in dimensions of general health, physical functioning, and vitality. However, in the multivariate regression analysis, no significant association was observed between occupational status and health dimensions. Findings of this study are in line with those of other studies conducted in Iran (16, 17). Although occupation is an important factor in women's health, and it is expected that women's occupational status would have a positive impact on women's empowerment, hence on their health, Lack of statistical difference between health dimensions in working women and housewives could be due to the balancing weights of positive and negative workrelated factors such as confidence and increased stress. Women with active jobs and high stress report more health problems (38). In European countries, the impact of occupational status on health was weaker compared to income or education, and its association with health was not the same across European countries. As in the central European countries, a positive association was observed, and in southern and eastern European countries a negative association. Negative association shows that in less developed countries, occupational status does not ensure better health, and health is associated with

ital status on health is different in various cultures

(34). Iranian married women, due to their huge

responsibilities in carrying out household duties

www.SID.ir

better income (37). Occupational structure such as salary level, workplace quality, and prestige of the job are effective in health (39), which was not measured in this study.

In terms of the variable of sufficiency of income for expenses, all health dimensions, except role limitations due to physical problems, had a significant association with this variable, that is, mean scores of health dimensions in women with affirmative answers to this question was significantly higher compared to women with negative answers. In other studies as well, all health dimensions in SF-36 had a significant association with people's wealth (4, 27). Women with low income reported poorer health dimensions (1, 30). Low-income women manage their own health less, which may be due to the expenses, travel, or receiving health information (30). Low income exposes people to such stressors as poorer life conditions, or economic pressures that predispose them to mental problems, where providing mental health care services can be helpful (33).

### Study strengths and limitations

Strength of this study is in the use of SF-36 to assess multiple health dimensions, which is a direct health measure, and as the "gold standard measure" has potential use in directing resources to health needs (40). Besides, according to being Tehran a capital city with mixed population, it reaches a maximum of the study's generalizability. Study limitation is that, the manner these variables impact health status was not studied. The causal pathways underlying women's social determinants of health in a lifespan are necessary to determine the right interventions and effective strategies to prevent health inequalities. Qualitative approach is suggested for the future researches to explain and interpret the findings and give reasons for connectivity or the lack of it.

# Conclusion

Most dimensions of health of women of reproductive age had a significant association with educational and income levels, and their reflection on negative experiences in women's health is alarming. Women's health is the basis for family and society's health, thus, efforts to make a healthy society as a platform for women's health are essential. In policymaking, in addition to the goal of reducing income disparities, attention must also be paid to women's education that actively contributes to maintaining their health. Making necessary arrangements for education and employment of women, while maintaining respect for their roles as mothers and wives, appear a necessity. Interaction between social determinants of health and health status must be considered in policymaking, and there is a need for policies that would enhance health of women in the low education and income brackets.

# **Ethical considerations**

The authors have completely observed Ethical issues (Including Informed Consent, confidentiality for information, plagiarism, double publication and/or submission, redundancy, etc.).

# Acknowledgement

This work was supported by Tehran University of Medical Sciences [grant no. 91-04-62-20184]. The authors would like to thank all women who participated in the present study. The authors declare that they have no competing interests.

### References

- Leipert BD, Reutter L (2005). Women's health in northern British Columbia: the role of geography and gender. <u>Can J Rural Med.</u>, 10(4): 241-53.
- World Bank (2008).International Economics Department.World Bank Publications: Development Data Group.In World Development Indicators.
- 3. Berke EM (2010). Geographic information systems (GIS): recognizing the importance of place in primary care research and practice.*J Am Board Fam Med*, 23: 9-12.
- 4. CDC, 2014. Social Determinants of Health. http://www.cdc.gov/socialdeterminants/Definitions.html.
- 5. D'Souza MS, Karkada SN, Somayaji G (2013). Factors associated with health-related quality

of life among Indian women in mining and agriculture. *Health and Quality of Life Outcomes*, 11(9): 9-11.

- 6. World Health Organization (2008).Closing the gap in a generation: health equity through action on the social determinants of health. Final report of the Commission on Social Determinants of Health. Geneva: Commission on Social Determinants of Health.
- Hofman K, Blomstedt Y, Addei S, Kalage R, Maredza M, Sankoh O et al. (2013). Addressing research capacity for health equity and the social determinants of health in 3 African countries: the INTREC programme. *Glob Health Action*, 6: 19668.
- 8. World Health Organization (2012).Closing the Gap in a Generation: Health Equity through Action on the Social Determinants of Health. Accessed on 26 June 2012: http://www.who.int/social\_determinants/the commission/finalreport/en/index.html.
- Smedley BD (2012). The lived experience of race and its health consequences. *Am J Public Health*,102(5): 933-5.
- Morency P, Gauvin L, Plante C, Fournier M, Morency C (2012). Neighborhood social inequalities in road traffic injuries: the influence of traffic volume and road design. *Am J Public Health*,102(6): 1112-9.
- 11. Bradley L (2012). The Social Determinants of Health.JANAC, 23(3): 181-3.
- Todorova IL, Tucker KL, Jimenez MP, Lincoln AK, Arevalo S, Falcón LM (2013). Determinants of self-rated health and the role of acculturation: implications for health inequalities. *Ethn Health*, 18(6): 563–85.
- Broom DH, D'Souza RM, Strazdins L, Butterworth P, Parslow R, Rodgers B (2006). The lesser evil: bad jobs or unemployment? A survey of mid-aged Australians. *Soc Sci Med*, 63: 575-86.
- McFadden E, Luben R, Bingham S, Wareham N, Kinmonth AL, Khaw KT (2008). Social inequalities in self-rated health by age: crosssectional study of 22,457 middle-aged men and women. *BMC Public Health*, 8: 230.
- Marmot M, Friel S, Bell R, Houweling TA, Taylor S (2008). Closing the gap in a generation: health equity through action on the social determinants of health. *Lancet*, 372(9650): 1661-9.

- Ahmad-Nia S (2002). Women's work and health in Iran: a comparison of working and nonworking mothers. *Soc Sci Med*, 54(5): 753–65.
- Saravi FK, Navidian A, Rigi SN, Montazeri A (2012). Comparing health-related quality of life of employed women and housewives: a cross sectional study from southeast Iran. BMC Women's Health, 12: 41.
- Statistical Center of Iran: National Census of Population and Housing of Iran (2006). http://amar.sci.org.ir.
- Rim H, Kim H, Lee K, Chang S, Hovell MF, Kim YT, et al (2011). Validity of Self-reported Healthcare Utilization Data in the Community Health Survey in Korea. J Korean Med Sci, 26 (11): 1409-14.
- 20. Statistical center of Iran (2011). http://www.amar.org.ir/Portals/1/Iran/census-2.pdf.
- 21. Ranji UR, Salganicoff A (2011). Women's Health Care Chart book.Key Findings from the Kaiser Women's Health Survey, 1-40.
- 22. Baheiraei A, Bakouei F, Mohammadi E, Hosseini M (2014). Social capital in association with health status of women in reproductive age: study protocol for a sequential explanatory mixed methods study. *Reproductive Health*, 11:35.
- Hemingway H, Nicholson A, Stafford M, Roberts R, Marmot M (1997). The Impact of Socioeconomic Status on Health Functioning as Assessed by the SF-36 Questionnaire: The Whitehall II Study. *Am J Public Health*,87(9): 1484-90.
- 24. Ware JE, Brook RH, Davies-Avery A (1980).Model of Health and Methodology. Santa Monica, CA: RAND Corporation, R-1987/1-HEW.
- Montazeri A, Goshtasebi A, Vahdaninia M, Gandek B (2005). The Short Form Health Survey (SF-36): translation and validation study of the Iranian version. *Qual Life Res*, 14(3):875–82.
- Ware JE, Snow KK, Kosinski M, Gandek B (1993). SF-36 health survey: manual and interpretation guide: The Health Institute, New England Medical Center.
- 27. Daher AM, Ibrahim HS, Daher TM, Anbori AK (2011). Health related quality of life among Iraqi immigrants settled in Malaysia. *BMC Public Health*, 11: 407.

- Mukhaimer JA (2007). Assessment of health status of Arab-American women in Michigan A Preliminarystudy. http://www.equityhealthj.com/content/12/1/25.
- 29. Ware J.E.Jr (2006). Health survey update. Retrieved December5. http://www.sf-36.org.
- 30. Williams JS, Cunich M, Byles J (2013). The impact of socioeconomic status on changes in the general and mental health of women over time: evidence from a longitudinal study of Australian women. *Int J Equity Health*, 12:25.
- Hills M, Mullett J (2005). Primary health care: a preferred health service delivery option for women. *Health Care Women Int*, 26(4): 325-39.
- Byles J, Dobson A, Pachana NA, Tooth L, Loxton D, Berecki J, et al (2010). Women, health and ageing: findings from the Australian Longitudinal Study on Women's Health. Brisbane: ALSWH: In: Report prepared for Australian Government Department of Health & Ageing, 268.
- Murata C, Kondo K, Hirai H, Ichida Y, Ojima T (2008). Association between depression and socio-economic status among communitydwelling elderly in Japan: the Aichi Gerontological Evaluation Study (AGES). *Health Place*,14(3): 406-14.
- Artazcoz L, Cortès I, Borrell C, Escribà-Agüir V, Cascant L (2011). Social inequalities in the association between partner/marital status and

health among workers in Spain.*SocSci* Med,72(4): 600-607.

- Azizi F, Ainy E (2008). Women, occupation, marital status and non-communicable disease risk factors: findings from the Tehran Lipid and Glucose Study. *Payesh*, 7(1): 11-15.
- Szanton SL, Seplaki CL, Thorpe RJ Jr, Allen JK, Fried LP (2010). Socioeconomic status is associated with frailty: the Women's Health and Aging Studies. J Epidemiol Community Health,64(1): 63-7.
- Alvarez-Galvez J, Rodero-Cosano ML, Motrico E et al. (2013). The impact of socio-economic status on self-rated health: study of 29 countries using European social surveys (2002-2008). *Int J Environ Res Public Health*, 10(3): 747-61.
- 38. Slopen N, Glynn RJ, Buring JE, Lewis TT, Williams DR, Albert MA (2012). Job strain, job insecurity, and incident cardiovascular disease in the Women's Health Study: results from a 10-year prospective study. *PLaS One*, 7(7): e40512.
- 39. Burke RJ (2002). Work stress and women's health: Occupational status effects. *Journal of Business Ethics*, 37(1): 91-102.
- Newbold K, Eyles J, Birch S, Spencer A (1998). Allocating resources in health care: alternative approaches to measuring needs in resource allocation formula in Ontario. *Health Place*, 4 (1): 79-89.