



Assessing Predictive Validity of Pressure Ulcer Risk Scales- A Systematic Review and Meta-Analysis

*Seong-Hi PARK¹, *Hea Shoon LEE²*

1. *Dept. of Nursing, College of Medicine, Soonchunhyang University, Cheonan, Korea*
2. *Dept. of Nursing, College of Life Science and Nano Technology, Hannam University, Daejeon, Korea*

***Corresponding Author:** Email: lhs7878@hanmail.net

(Received 23 Jul 2015; accepted 10 Sep 2015)

Abstract

Background: The purpose of this study was to present a scientific reason for pressure ulcer risk scales: Cubbin& Jackson modified Braden, Norton, and Waterlow, as a nursing diagnosis tool by utilizing predictive validity of pressure sores.

Methods: Articles published between 1966 and 2013 from periodicals indexed in the Ovid Medline, Embase, CINAHL, KoreaMed, NDSL, and other databases were selected using the key word “pressure ulcer”. QUADAS- II was applied for assessment for internal validity of the diagnostic studies. Selected studies were analyzed using meta-analysis with MetaDisc 1.4.

Results: Seventeen diagnostic studies with high methodological quality, involving 5,185 patients, were included. In the results of the meta-analysis, sROC AUC of Braden, Norton, and Waterlow scale was over 0.7, showing moderate predictive validity, but they have limited interpretation due to significant differences between studies. In addition, Waterlow scale is insufficient as a screening tool owing to low sensitivity compared with other scales.

Conclusion: The contemporary pressure ulcer risk scale is not suitable for uninformed practice on patients under standardized criteria. Therefore, in order to provide more effective nursing care for bedsores, a new or modified pressure ulcer risk scale should be developed upon strengths and weaknesses of existing tools.

Keywords: Pressure ulcer, Sensitivity, Specificity, Meta-analysis

Introduction

Pressure ulcers is a significant quality index of patient care and one of the most crucial challenges encountered in current health and medical treatment (1). Because incidence rate and prevalence rate of pressure ulcers is dependent on patients' characteristics, each study does not show same results, however, Prevalence rate of Pressure ulcers in Americans increased significantly, from 7% in 2000 (2) to 15% in 2009 (3). In Canada, 25% of patients in acute medical institutions and 30% in non-acute medical institutions had bed sore outbreaks (4). Lastly, 11% of patients (5) and 28.2% of nervous critical patients in Korea (6) had pressure ulcers. Millions of Pressure ulcers outbreaks occur annually worldwide and 60million patients die of complication caused by Pressure ulcers (7).

Pressure ulcers decrease quality of life, while increasing and medical expenses sharply as the pressure ulcers takes a serious turn (8). In addition, the duration of hospital stay will be extended; patient's function recovery will be delayed (9), which may cause problems such as the need of additional nurse workforce (10). Therefore, active prevention is important in reducing the occurrence of pressure ulcers (11). Ideal pressure ulcers prevention is possible, when we accurately select a few high-risk pressure ulcers occurrences among the total patient group by using qualified pressure ulcer risk scales. It also can be solved when we conduct education for patient/guardian and nursing management with concentrated prevention activity, thereby maximizing the efficiency (12).

Norton scale was developed and applied in the clinic for the first time. This scale was found to predict pressure ulcers of senile patients in England (13). In 1973, Gosnell revised the Norton scale for inpatients in long-term care facilities (14). In 1984, Shannon developed a scale made up of eight items (15), and in 1985, England, the Waterlow scale was developed in England for senile patients in acute medical institutions (16). In 1987, the Braden scale was developed for senile patients in long-term care facilities in the USA, lastly (17); also, in 1991, the Cubbin & Jackson scale was developed for ICU Patients (18). Other countries developed pressure ulcer risk scales that are suitable for their own patients (19-21). Beside from the Braden scale, predictive validity verification of tool application progress or repetitive studies was not performed so that development of a new scale is required (22-23).

In this study, we compared and analyzed the Cubbin & Jackson, modified Braden, Norton, and Waterlow pressure ulcer risk scales. We also aimed to provide scientific evidence of the accuracy of predictions of pressure sores through a diagnostic method oriented meta-analysis for the use of various pressure ulcer assessment tools in the clinic.

Methods

This study was conducted using the *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy* from the Cochrane Collaboration (24) and *Preferred reporting items for systematic reviews and meta-Analyses* from Preferred Reporting Items for Systematic Reviews and Meta-Analyses [PRISMA] (25).

Strategy for considering studies for this search

Criteria for considering studies for this review

By using the pressure ulcer risk scale: Cubbin & Jackson, modified Braden, Norton and Waterlow, we included studies that evaluated predictive validity of pressure ulcers outbreak in the proper patient group. Risk of pressure ulcers was judged upon each criterion, Based on a cut-off point suggested by each researcher

Pressure ulcer outbreak was verified with a rating scale of objective skin condition by professional groups and individual scholars: USA, National

Pressure Ulcer Advisory Panel [NPUAP], European Pressure Ulcer Advisory Panel [EPUAP], Agency for Health Care Policy and Research [AHCPR], Torrence Developmental Classification of Pressure Sore [TDCPS].

Search methods for identification of studies

Electronic database search was conducted on October 1st, 2013. We used electronic databases, including KoreaMed, NDSL, KERIS, Ovid-Medline, Embase, Cochrane Library, and CINAHL Plus with Full Text. We added our national medical/nursing academic journal search. Search words used were 'pressure ulcer', 'decubitus ulcer', 'skin ulcer', 'bed sore', and 'risk assessment'.

Risk of bias in included studies and data extraction

The quality of literature was evaluated independently by two authors using QUADAS-II (Quality Assessment of Diagnostic Accuracy Studies- II) (26). On data extraction, we filled out: type of study, location of study, number of research institutes, gender distribution of subjects, age, duration of hospital stay, and cut-off point of each scale, from selected literature after deciding on the basic form of evidence table; in addition, diagnosis result of pressure ulcer risk scale value of TP, FP, FN, TN was written. On the basis of it, we wrote a 2×2 table of specification and recalculated sensitivity, specificity, positive LR, negative LR, DOR, and 95% confidence interval [95% CI] using '2-way Contingency Table Analysis' program.

Statistical analysis

Studies were analyzed using meta-analysis with MetaDiSc 1.4. The general rule of the statistical model is to analyze with random effect estimation in order to reflect the differences between studies (24), and analyzed with total sensitivity, specificity, positive LR, negative LR, DOR, and sROC curve. In statistics of sROC curve, we recorded the examination accuracy through the Area under the Curve [AUC] and index Q* value. From AUC figure, AUC=0.5 is noninformative examination; 0.5<AUC≤0.7 is less accurate; 0.7<AUC≤0.9 is moderate; 0.9<AUC<1 is perfect examination (27), index Q* reflecting sensitivity and specificity on

ROC curve is judged on the basis of I^2 when it corrects 100% (28). Heterogeneity between studies was verified with sight while having a common part in the confidence interval and estimation of efficacy through picture of a forest. In addition, we investigated it using I^2 test, decreasing significance level to 5%; the standard of judgment of I^2 : $I^2 \leq 25\%$ has low heterogeneity; $25\% < I^2 \leq 75\%$ has moderate heterogeneity; $I^2 > 75\%$ has high heterogeneity (29).

Results

Finally, 17 documents were selected. The detailed selection method is shown in Figure 1. Seven documents (1, 23, 30-34) were about the Norton tool, 6 documents (30-31, 34-37) were about Waterlow, 6 documents (23, 32, 38-41) were about modified Braden, and the other 4 documents (6, 40, 42-43) were about Cubbin & Jackson (Table 1).

Risk of bias in included studies

None of the documents was estimated to have “high” risk of inequality after evaluating the quality of the 17 selected documents.

The Test of Validity and the Quality of Selected Documents of Pressure Ulcer risk scale Norton Scale

There were seven literature studies (1, 23, 30-34) reporting diagnostic accuracy of the Norton scale, involving 2,899 patients. All literature studies are prospective studies, except 1 (33) out of 7, involving inpatients who have no pressure ulcers. Regarding average age, there was one study about 50s (32), one study about 80s (1), three literature studies on 60s, and the two remaining literature studies (30, 34) that do not mention age. Based on standard reference, three literature studies (23, 30-21) use AHCPR, two of them (32-33) use NPUAP, EPUAP (1) and TDCPS (34) are used in one literature study separately. The three studies use cut-off point: 14points (1, 32, 24) and 16points (23, 31, 34), and one (30) utilizes 15points. Total sensitivity was 0.75 (95% CI 0.70, 0.79) (Figure 2-E), and total specificity was 0.57 (95% CI 0.55, 0.59). The heterogeneity between studies was high, as 88.9% ($X^2=54.29$, $P<.001$), 98.3% ($X^2=362.58$, $P<.001$) (Figure 2-F), and sROC AUC was 0.82 (SE=0.05) while Q^* value was 0.75 (SE=0.05)(Figure 3-C).

Table 1: Comparison of characteristics of pressure ulcer risk assessment tools

	Norton scale	Waterlow scale	Modified Braden scale	Cubbin & Jackson scale
Created year	1962 ¹⁾	1985 ²⁾	1987 (by Braden et al.) ³⁾ 1991 (by Choi & Song) ⁴⁾ 1998 (by Pang & Wong) ⁵⁾	1991 ⁶⁾
Specialty	Older people/generic	Orthopaedic/generic	Generic	Intensive care
Risk factors to assess pressure sore	Physical condition Mental status Activity Mobility* Continence*	Sex Age Build Appetite Nurses' visual assessment of skin condition Mobility* Continence* Factors contributing to tissue malnutrition Neurologic deficits Major surgery or trauma Medication	Sensory perception Moisture* Activity Mobility* Nutrition status Friction/shear by Choi & Song Temperature Use of medication by Pang & Wong Skin type Body build	Age Weight Skin condition Mental status Mobility* Nutrition Respiration Continence* Hygiene Hemodynamic status

Notes. 1) Norton, McLaren & Exton-Smith (1962); 2) Waterlow (1985); 3) Bergstrom, Braden, Laguzza, & Holman (1987); 4) Choi & Song (1991) 5) Pang & Wong (1998); 6) Cubbin & Jackson (1991); * Common risk factors to assess pressure sore




Fig. 1: Flow diagram of article selection

Through detailed analytical research, the total sensitivity of 60s was 0.69 (95% CI 0.61, 0.76) with a cut-off point of 94.2% ($X^2=34.75$, $P<.001$) In Standard reference AHCPR (23, 30-31), total sensitivity was 0.61 (95% CI 0.52, 0.70), and heterogeneity between literature studies was 94.5% ($X^2=36.17$, $P<.001$). The total sensitivity of NPUAP was 0.82 (95% CI 0.69, 0.91) and the heterogeneity between literature studies was 0.0% ($X^2=0.40$, $P=.53$). In all cases of total specificity, the heterogeneity between literature studies was over 90%.

Waterlow Scale

There were six literature studies (30-31, 34-37) applying the Waterlow scale, involving 1,268 inpatients. All literature studies were prospective studies in cases of no pressure ulcers of inpatients, and the subjects of study were elderly men over their 60s of average age. Standard reference was used in two literature studies with NPUAP (36-37) and AHCPR (30-31), and EPUAP (35), TDCPS (34) in one literature study. The cut-off point was ap-

plied variously; one literature study had 9 points (30), 15points (37), and 17points (36) and two literature studies had 16 points (31, 34). This cut-off point was not reported (35).

Total sensitivity was 0.55 (95% CI 0.49, 0.62) (Figure 2-G), and total specificity was 0.82 (95% CI 0.80, 0.85) (Figure 2-H). In addition, total sensitivity was 0.55 (95% CI 0.49, 0.62), between studies was high, as 90.0% ($X^2=49.76$, $P<.001$), 97.2% ($X^2=177.24$, $P<.001$), and sROC AUC was 0.82 (SE=0.03) while Q^* value was 0.75 (SE=0.03), regarding the heterogeneity, in the result of meta-analysis. Through detailed analytical research, in the case of standard reference NPUAP, total sensitivity was 0.68 (95%CI 0.43, 0.87) but heterogeneity between literature did not exist, as 0.0% ($X^2=0.05$, $P=0.83$), while heterogeneity between studies of total sensitivity and specificity of standard reference AHCPR was high, over 80%. Modified Braden scale was used in six studies (23, 32, 38-41), and the Cubbin & Jackson scale was applied in four studies (6, 40, 42-43).




Fig. 2: Diagnosis test accuracy of pressure ulcer risk assessment scales

Modified Braden Scale

Regarding the literature on the Modified Braden scale, four studies (23, 39-41) were reported in Korea and two studies (32, 38) in China. All literature studies are prospective studies aimed for patients who do not have pressure ulcers when they were in ward (38-39) and ICU (23, 40) which involve 1,314 patients. Three literature studies (32, 39-40) are about average age 50s, two literature studies about 60s, and one literature study about 70s (38). Three volumes are about IC; three volumes are about the study of standard reference AHCPR (23, 40-41); two volumes are about NPUAP (32, 38); one volume is about Bergstrom (39).

Total sensitivity was 0.95 (95% CI 0.91, 0.98) and heterogeneity between literature studies was moderate, as 49.9% ($X^2=9.99$, $P=0.08$)(Figure 2-C).

Total specificity was 0.70 (95% CI 0.68, 0.73) and heterogeneity between literature studies was high, as 96.4% ($X^2=138.48$, $P<.001$)(Figure 2-D). sROC AUC was 0.93 (SE=0.03); Q^* value was 0.87 (SE=0.03)(Figure3-B).

Through detailed analytical research, In the case of 50s, the total sensitivity was 0.92 (95% CI 0.83, 0.97), and heterogeneity between literature studies was 0.0% ($X^2=1.04$, $P=.60$). The total sensitivity of 60s was 1.00 (95% CI 0.95, 1.00) while heterogeneity between literature studies was 0.0% ($X^2=0.00$, $P=1.00$). The total sensitivity of standard reference was 0.98 (95% CI 0.94, 1.00), and heterogeneity between literature studies was 50.9% ($X^2=4.07$, $P=.13$) while NPUAP is only used in literature of China. Total specificity was over 90% in every case.




Fig. 3: Summary receiver operating characteristic curve

Cubbin & Jackson Scale

Literature on the Cubbin & Jackson Scale was four volumes (6, 40, 42-43), conducted in Korea with 662 ICU patients as a prospective study. Except

Jun exec (42), studies were conducted with patients who did not have pressure sores, and the average age of patients was in their 50s (40, 43) and 60s (6, 42). There were two literature studies (40,

42) with standard reference AHCPR, and one study (6) with NPUAP, one volume (43) on Low-thian. The cut-off point was used in two volumes (6, 42) in 24 points and 28 points (40) and 26 points (43).

The total sensitivity was 0.67 (95% CI 0.60, 0.74) (Fig. 2-A), and the total specificity was 0.75 (95% CI 0.70, 0.79) (Fig. 2-B). Each heterogeneity between literature studies was high, as 92.8% ($X^2=41.54$, $P<.001$), 82.0% ($X^2=16.70$, $P=.001$). sROC AUC was 0.82 (SE=0.06) and Q^* value was 0.75 (SE=0.05) (Fig.3-A).

Through detailed analytical research, despite division of 50s and 60s, the heterogeneity was high, over 90%; from detailed analysis of two literature studies that used standard reference AHCPR, the total sensitivity was 0.92(95% CI 0.83, 0.97) and the heterogeneity between literature studies was 5.4% ($X^2=1.06$, $P=.30$); the total specificity was 0.75 (95% CI 0.70, 0.81) and the heterogeneity between literature studies was 91.4% ($X^2=11.64$, $P=.001$).

Comparison of Norton scale and Waterlow scale

Two pressure ulcer risk scales were suggested in six literature studies out of 17 included in this study. Three literature studies are about comparison of Norton scale and Waterlow scale with the same subjects (30-31, 34). There were two studies (23, 32) comparing Norton scale and modified Braden, and only one study (40) comparing modified Braden and Cubbin & Jackson scale. In the results of detailed analysis, Norton scale had 0.53 (95% CI 0.43, 0.63) total sensitivity and 81.4% ($X^2=10.73$, $P=.005$) heterogeneity between literature studies, while total specificity was 0.88 (95% CI 0.83, 0.91) and heterogeneity between literature studies was 97.9% ($X^2=93.06$, $P<.001$); sROC AUC was 0.81 (SE=0.06) and Q^* value was 0.75 (SE=0.05). Waterlow scale had 0.74 (95% CI 0.64, 0.82) total sensitivity and 88.4% ($X^2=17.25$, $P=.001$) heterogeneity between literature studies while total specificity was 0.66 (95% CI 0.60, 0.72) and heterogeneity between literature studies was 95.1% ($X^2=40.54$, $P<.001$); sROC AUC was 0.82 (SE=0.03) and Q^* value was 0.76 (SE=0.03) (Table 2-3).

Discussion

In this experiment, we conducted a meta-analysis for selection of an appropriate pressure ulcer risk scale for patients; to examine the validity of Norton, Waterlow, Modified Braden, and Cubbin & Jackson tools, we analyzed 17 volumes and 5,185 diagnostic results of patients. Predictable validity is standard as we can see from an examination and evaluation tool that guides estimation of future results. The standard validity is comprised of sensitivity, specialty, positive and negative predictability (44). High sensitivity leads to low specialty, which is useful in measurement of the positive and negative predictability but not for precise meta-analysis because it influences the prevalence rate (24). To compensate for the defect, we can use the ROC analysis method for the calculation of general validity. ROC analysis method is a widely used method for management of not only a medical diagnosis and standard for decision-making but also a tool for development and standardization (45).

According to meta-analysis, the most sensitive tool is the modified Braden and is Waterlow, which is the tool with the greatest specialty. All pressure ulcer risk scales except the modified Braden, which have the same sROC AUC. Therefore, each tool has moderate accuracy for predictable validity of pressure sores of approximately $0.7 < AUC \leq 0.9$. Only the Waterlow tool was interpreted as insufficient because pressure ulcer assessment is a screening inspection that prefers a sensitive tool rather than a specialized tool. However, after a detailed analysis of the Norton scale and Waterlow scale, the Waterlow scale has a higher sensitivity (0.53 versus 0.74) and low specialty (0.88 versus 0.66) while they both have similar measured value in the ROC analysis method.

In order to analyze the problem in the studies, we classified them as average life expectancy, locations of hospitalization, and reference average standard. Four tools tend to lose difference between studies of combined sensitivity in common, depending on standard reference. To interpret the result of the research, using a skin assessment tool of consistent standardized factors of pressure ulcers in the clinic will increase the discrimination of pressure ulcer risk scale. Not all differences be-

tween studies were decreased, except the modified Braden's tool in patients' age. However, it is shown as a phenomenon of decreasing discrimination, because fewer studies at a given age stage might induce analysis in the higher proportion range of age, or no difference was shown from the late 50s to early 60s, despite the division of age. At places of hospitalization, only ICU is targeted in use of Cubbin & Jackson's tool, and modified Braden's tool can only be analyzed in depth, because Norton and Waterlow's tools are for inpatients. In the result of division: ICU and ward, we can see the decreasing aspect of difference between studies. To sum up, application in specific groups, which can well reflect the properties of each tool, will increase predictive validity, because pressure ulcer risk scales are easily affected by age, locations of hospitalization, and standard reference.

This study uses original cut-off points. Optimized cut-off point between studies does not show consistency in each instrument, and transition width is wide depending on age group. To interpret the result, it is misleading to determine and use the cut-off points of the current pressure ulcer risk scale, and applying with category would be more appropriate than specific threshold.

In other words, Norton, Waterlow, Modified Braden and Cubbin & Jackson scale cannot show consistent measured value, and they cannot determine the risk of pressure sore outbreak because of heterogeneity between studies, thus they are not suitable for use as general tools in the clinic toward inpatients. This statement is in agreement with studies of Anthony *et al.* (46) in that current pressure ulcer risk scales do not predict the outbreak of pressure ulcers, and some components include unnecessary items, it is apart from correlation of pressure sore outbreak. Only when we think about the alternative that can be applied instantly at this important time of prevention of pressure ulcers, use of the Norton scale first, which can be used more easily in the clinic, would be an effective method if the predictive validities of the four scales are similar, and then use the Waterlow scale with high specificity to increase the accuracy of the current pressure risk assessment.

Conclusion

In this research, predictive validity of Waterlow, Modified Braden, and Cubbin & Jackson's pressure ulcer risk scales was determined by meta-analysis based on 17 well-designed studies assessing diagnosis. As sROC AUC is over 0.7, all analyzed instruments show moderate predictive validity, but they have limited interpretation due to high differences between studies. In addition, Waterlow's instrument is insufficient for use as a screening tool owing to low sensitivity. Pressure ulcer risk scales in current use are not suitable for uniform practice on inpatients under standardized criteria. Therefore, in order to provide more effective nursing care of pressure ulcers, not only in-depth analysis reflecting patients' characteristics is required, but also a new or modified pressure ulcer risk scale should be developed with complementation of strength and weakness in existing tools.

Ethical Considerations

Ethical issues (Including plagiarism, Informed Consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc) have been completely observed by the authors.

Acknowledgments

This work was supported by the Soonchunhyang University Research Fund. Besides, it was supported by Hannam University, Academic Research Fund of 2016. The authors declare that there is no conflict of interests.

References

1. Defloor T, Grypdonck MFH (2005). Pressure ulcers: validation of two risk assessment scales. *J Clin Nurs*, 14(3): 373-82.
2. Whittington K, Patrick M, Roberts JL, Pieper B (2000). A national study of pressure ulcer prevalence and incidence in acute care hospitals. *J Wound Ostomy Continence Nurs*, 27(4): 209-15.
3. National Pressure Ulcer Advisory Panel and European Pressure Ulcer Advisory Panel

- (NPUAP & EPUAP). (2009). Prevention and treatment of pressure ulcers: clinical practice guideline. Washington DC: National Pressure Ulcer Advisory Panel and European Pressure Ulcer Advisory Panel.
4. Woodbury MG, Houghton PE (2004). Prevalence of pressure ulcers in Canadian health care settings. *Ostomy Wound Manage*, 50(10): 22-38.
 5. Cho MS, Park IS, Kim KH, Woo KS, Joo YH, Jeong EH, et al. (2004). Evaluation of predictive validity for the pressure ulcer risk Assessment tool in a medical ward inpatient: Using Braden Scale. *The Korean Nurses*, 43(2): 68-82.
 6. Im MJ, Park HS (2006). A study on the pressure ulcers in neurological patients in intensive care units. *J Korean Acad Fundam Nurs*, 13: 190-9.
 7. Wound Ostomy and Continence Nurses Society (2011). Wound, ostomy and continence nurses society. Overview white paper. Retrieved April 4, 2011, Available from: http://www.wocn.org/resource/resmgr/white_papers.pdf
 8. Bennett G, Dealey C, Posnett J (2004). The cost of pressure ulcers in the UK. *Age Ageing*, 33(3):230-5.
 9. Russo CA, Steiner C, Spector W (2008). Hospitalizations related to pressure ulcers among adults 18 years and older, 2006 (HCUP Statistical Brief #64). Rockville (MD): Agency for Healthcare Policy and Research (US).
 10. Bours GJ, Halfen RJ, Berger MP, Huijter A-Saad H, Grol RT (2003). Development of a model for case-mix adjustment of pressure ulcer prevalence rates. *Med Care*, 41(1):45-55.
 11. Choi Sj, Bae SY, Choi JY, Bang HJ (2005). Development and predictive validity of pressure ulcer predicting scale for patients with neurologic condition. *J Korean Clin Nurs Res*, 11(1):95-108.
 12. Perneger TV, Raè AC, Gaspoz JM, Borst F, Vitek O, Héliot C (2002). Screening for pressure ulcer risk in the acute care hospital: development of a brief bedside scale. *J Clin Epidemiol*, 55(5): 498-504.
 13. Norton D, McLaren R, Exton-Smith AN (1962). An investigation of geriatric nursing problems in hospital. London: National Corporation for the Care of Old People.
 14. Gosnell DJ (1973). An assessment tool to identify pressure sores. *Nurs Res*, 22(1): 55-9.
 15. Shannon ML (1984). Five famous fallacies about pressure sores. *Nursing*, 14(10):34-41.
 16. Waterlow J (1985). Pressure sores: a risk assessment card. *Nurs Times*, 81(48): 49-55.
 17. Bergstrom N, Braden BJ, Laguzza A, Holman V (1987). The Braden Scale for Predicting Pressure Sore Risk. *Nurs Res*, 36(4): 205-10.
 18. Cubbin B, Jackson C (1991). Trials a pressure ulcer risk calculator for intensive therapy patients. *Intensive Care Nurs*, 7(1): 40-44.
 19. Mertens E, Dassen T, Scheufele R, Halfens RJ, Tannen A (2010). Diagnostic validity of the care dependency scale as a screening tool for pressure ulcer and fall risks in Dutch and German hospitals. *Cent Eur J Med*, 5(5):577-87.
 20. Page KN, Barker AL, Kamer J (2011). Development and validation of pressure ulcer risk assessment tool for acute hospital patients. *Wound Rep Reg*, 19(1):31-7.
 21. Suriadi, Sanada H, Sugama J, Thigpen B, Subuh M (2008). Development of a new risk assessment scale for predicting pressure ulcers in an intensive care unit. *Nurs Crit Care*, 13(1):34-43.
 22. Schoonhoven L, Haalboom JRE, Bousema MT, Algra A, Grobbee DE, Grypdonck MH, et al.(2002). Prospective cohort study of routine use of risk assessment scales for prediction of pressure ulcers. *BMJ*, 325: 797-800.
 23. Lee Y, Jeong I, Jun S (2003). A comparative study on the validity among pressure ulcer risk assessment scales. *J Korean Acad Nurs*, 33(2): 162-9.
 24. Macaskill P, Gatsonis C, Deeks J, Harbord R, Takwoingi T (2010). *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy version 1.0*. The Cochrane Collaboration. Retrieved May 30, 2013, Available from: <http://www.cochranehandbook.org>.
 25. Moher D, Liberati A, Tetzlaff J, Altman DG, Antes G, Atkins D, et al. (2009). Preferred reporting items for systematic reviews and meta-Analyses: The PRISMA Statement. *Ann Intern Med*, 151(4):264-9
 26. Whiting PF, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, et al. (2011). QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med*, 155(8):529-536.
 27. Greiner M, Pfeiffer D, Smith RD (2000). Principles and practical application of the receiver-operating characteristic analysis for diagnostic tests. *Prev Vet Med*, 45(1-2):23-41.
 28. Walter SD (2002). Properties of the summary receiver operating characteristic (SROC) curve for diagnostic test data. *Stat Med*, 21(9):1237-56.

29. Higgins JP, Thompson SG (2002). Quantifying heterogeneity in a meta-analysis. *Stat Med*, 21(11):1539-1558.
30. Charlier C (2001). Prevalence, incidence and risk: a study of pressure ulcers at a rural base hospital. *Primary Intention: The Australian Journal of Wound Management*, 9: 12-21.
31. Jalali R, Rezaie M (2005). Predicting pressure ulcer risk: comparing the predictive validity of 4 scales. *Adv Skin Wound Care*, 18(2):92-7.
32. Kwong E, Pang S, Wong T, Ho J, Shao-ling X, Lijun T (2005). Predicting pressure ulcer risk with the modified Braden, Braden, and Norton scales in acute care hospitals in Mainland China. *Appl Nurs Res*, 18(2):122-8.
33. Marrie RA, Ross JB, Rockwood K (2003). Pressure Ulcers: Prevalence, Staging, and Assessment of Risk. *Geriatrics Today: Journal of the Canadian Geriatrics Society*, 6: 134-40.
34. Pang SM, Wong TK (1998). Predicting pressure sore risk with the Norton, Braden, and Waterlow scales in a Hong Kong rehabilitation hospital. *Nurs Res*, 47(3):147-53.
35. Compton F, Hoffmann F, Hortig T, Strauss M, Frey J, Zidek W, et al. (2008). Pressure ulcer predictors in ICU patients: nursing skin assessment versus objective parameters. *J Wound Care*, 17(10):417-24.
36. Serpa LF, de Gouveia Santos VLC, Gomboski G, Rosado SM (2009). Predictive validity of Waterlow Scale for pressure ulcer development risk in hospitalized patients. *J Wound Ostomy Continence Nurs*, 36(6):640-6.
37. Webster J, Gavin, N, Nicholas C, Coleman K, Gardner G (2007). Validity of the Waterlow scale and risk of pressure injury in acute care. *Br J Nurs*, 19(6):S14-S22.
38. Chan WS, Pang SMC, Kwong EWY (2009). Assessing predictive validity of the modified Braden scale for prediction of pressure ulcer risk of orthopaedic patients in an acute care setting. *J Clin Nurs*, 18(11):1565-73.
39. Choi KS, Song M (1991). Test of predictive validity for the new pressure risk assessment scale. *Korean J Adult Nurs*, 3(1):19-28.
40. Kim EK, Lee SM, Lee E, Eom MR (2009). Comparison of the predictive validity among pressure ulcer risk assessment scales for surgical ICU patients. *Aust J Adv Nurs*, 26(4): 87-94.
41. Kim SS, Choi KS (2004). Evaluating the predictive validity for the new pressure sores risk assessment scale. *Korean J Adult Nurs*, 16(2): 183-190.
42. Jun S, Jeong I, Lee Y (2004). Validity of pressure ulcer risk assessment scales; Cubbin and Jackson, Braden, and Douglas scale. *Int J Nurs Stud*, 41(2):199-204.
43. Kim YK (1997). Evaluation of a pressure ulcer risk assessment tool (I). *Korean J Adult Nurs*, 9(2):272-85.
44. Lyder CH, Ayello EA (2008). Pressure Ulcers: A Patient Safety Issue. In Hughes RG (ed). *Quality: An Evidence-Based Handbook for Nurses*. Rockville (MD): Agency for Healthcare Research and Quality (US). AHRQ Publication No. 08-0043.
45. Knottnerus JA (2008). The evidence base of clinical diagnosis. ed. Park SH, Kang CB, translator. Seoul: E-Public.
46. Anthony D, Papanikolaou P, Parboteeah S, Saleh M (2010). Do risk assessment scales for pressure ulcers work? *J Tissue Viability*, 19(4):132-6.

Table 2: Characteristics of selected studies

Year of publication	Authors	Location	Types of studies	Setting	PUs at adm.	Participants			Reference standards	Cut off point	2×2 Table				Value (95% Confidence interval)			DOR	
						Age (yr)	M:F (n)	Total (n)			TP	FP	FN	TN	SN	SP	PLR		NLR
Norton scale																			
2005	Defloor & Grypdonck	Belgium	Pro	LTCF	PUs (-)	84.6± 7.9	369:1,403	1,772	EPUAP (1999)	<14	152	710	35	875	0.81 (0.75-0.86)	0.55 (0.55-0.56)	1.82 (1.65-1.95)	0.34 (0.25-0.46)	5.35 (3.60-7.98)
2005	Jalali & Rezaie	Iran	Pro	ICU & wards	PUs (-)	60.0	100:130	230	AHCPR (1994)	<16	36	0	38	156	0.49 (0.43-0.49)	1.00 (0.89-1.00)	-	0.51 (0.51-0.58)	-
2005	Kwong et al.	China	Prospective	-	PUs (-)	54.1± 16.9	253:176	429	NPUAP (1989)	<14	8	164	1	256	0.89 (0.51-0.99)	0.61 (0.60-0.61)	2.28 (1.28-2.56)	0.18 (0.01-0.81)	12.49 (1.57-268.73)
2003	Lee at al.	Korea	Pro	ICU	PUs (-)	62.0	64:48	112	AHCPR (1994)	<16	34	55	1	22	0.97 (0.86-1.00)	0.29 (0.23-0.30)	1.36 (1.12-1.42)	0.10 (0.01-0.62)	13.60 (1.80-283.14)
2003	Marric et al.	Canada	Retro	-	-	61.0± 18.0	98:90	188	NPUAP (2002)	≤14	37	121	9	21	0.80 (0.69-0.90)	0.15 (0.11-0.18)	0.94 (0.78-1.09)	1.32 (0.59-2.78)	0.71 (0.28-1.85)
2001	Charlier	Australia	Prospective	Hosp.	PUs (-)	-	29:33	62	AHCPR (1995)	<15	3	1	8	50	0.27 (0.08-0.36)	0.98 (0.94-1.00)	13.91 (1.38-344.2)	0.74 (0.64-0.98)	18.75 (1.42-536.60)
1998	Pang & Wong	Hong Kong	Pro	Hosp.	PUs (-)	45-92	52:54	106	TDCPS (1983)	<16	17	35	4	50	0.81 (0.60-0.94)	0.59 (0.54-0.62)	1.97 (1.29-2.46)	0.32 (0.10-0.75)	6.07 (1.71-23.53)
Waterlow scale																			
2010	Webster et al.	Australia	Pro	Ward	PUs (-)	65.3± 7.7	137:137	274	NPUAP (2007)	<15	8	55	4	207	0.67 (0.36-0.89)	0.79 (0.78-0.80)	3.18 (1.61-4.43)	0.42 (0.14-0.82)	7.53 (1.96-31.04)
2009	Serpa et al.	Brazil	Pro	ICU & wards	PUs (-)	71.1± 5.5	49:49	98	NPUAP (2007)	<17	5	30	2	61	0.71 (0.31-0.95)	0.67 (0.64-0.69)	2.17 (0.86-3.04)	0.43 (0.08-1.08)	5.08 (0.80-40.52)
2008	Compton et al.	Germany	Pro	ICU	PUs (-)	66.0	392:306	698	EPUAP (1999)		45	31	76	546	0.37 (0.31-0.43)	0.95 (0.93-0.96)	6.9 (4.50-10.66)	0.66 (0.59-0.75)	10.4 (6.03-18.08)
2005	Jalali & Rezaie	Iran	Pro	ICU & wards	PUs (-)	60.0	100:130	230	AHCPR (1994)	<16	47	28	27	128	0.64 (0.54-0.72)	0.82 (0.78-0.86)	3.54 (2.41-5.13)	0.45 (0.33-0.59)	7.96 (4.07-15.67)
2001	Charlier	Australia	Prospective	Hosp.	PUs (-)	-	29:33	62	AHCPR (1995)	<9	11	23	0	28	1.00 (0.71-1.00)	0.55 (0.49-0.55)	2.22 (1.37-2.22)	0.00 (0.00-0.60)	-
1998	Pang & Wong	Hong Kong	Pro	Hosp.	PUs (-)	45-92	52:54	106	TDCPS (1983)	<16	20	48	1	37	0.95 (0.76-1.00)	0.44 (0.39-0.45)	1.69 (1.24-1.80)	0.11 (0.01-0.62)	15.42 (2.02-322.34)
Modified Braden scale																			
<i>By Choi & Song</i>																			
2009	Kim et al.	Korea	Prospective	SICU	PUs (-)	58.1± 1.2	145:74	219	AHCPR (1994)	<21	38	55	2	124	0.95 (0.83-0.99)	0.69 (0.67-0.70)	3.09 (2.48-3.33)	0.07 (0.01-0.26)	42.84 (9.59-266.55)
2004	Kim & Choi	Korea	Prospective	ICU & ward	PUs (-)	60.7	-	211	AHCPR (1996)	<23	34	42	0	135	1.00 (0.88-1.00)	0.76 (0.74-0.76)	4.21 (3.39-4.21)	0.00 (0.00-0.16)	-
2003	Lee at al.	Korea	Prospective	ICU	PUs (-)	62.0	64:48	112	AHCPR (1996)	<24	35	63	0	14	1.00 (0.90-1.00)	0.18 (0.14-0.18)	1.22 (1.04-1.22)	0.00 (0.00-0.73)	-
1991	Choi & Song	Korea	Prospective	Wards	PUs (-)	54.1	89:57	146	Bergstrom (1987)	<24	14	11	2	119	0.88 (0.64-0.98)	0.92 (0.89-0.93)	10.34 (5.57-13.58)	0.14 (0.02-0.41)	75.73 (13.54-559.83)
<i>By Pang & Wong</i>																			
2009	Chan et al.	Hong Kong	Prospective	OS ward	PUs (-)	79.4± 10.9	30:167	197	NPUAP (2007)	<19	16	68	2	111	0.89 (0.65-0.98)	0.62 (0.60-0.63)	2.34 (1.61-2.65)	0.18 (0.03-0.59)	13.06 (2.75-84.99)
2005	Kwong et al.	China	Prospective	-	PUs (-)	54.1± 16.9	253:176	429	NPUAP (1989)	<14	8	105	1	315	0.89 (0.51-0.99)	0.75 (0.74-0.75)	3.56 (1.98-4.01)	0.15 (0.01-0.66)	24.00 (3.01-517.73)
Cubbin & Jackson scale																			
2009	Kim et al.	Korea	Prospective	SICU	PUs (-)	58.1± 1.2	145:74	219	AHCPR (1994)	<28	38	33	2	146	0.95 (0.83-0.99)	0.82 (0.79-0.83)	5.15 (3.94-5.66)	0.06 (0.01-0.21)	84.06 (18.39-531.62)

Table 2: Cond...

Year	Author	Country	Study Design	Setting	PU Type	Mean PU	PU Rate	Sample Size	Scale	Cut-off	TP	FP	TN	FN	PLR	NLR	DOR	AUC	Q*
2006	Im & Park	Korea	Prospective	ICU	PU (-)	62.1± 1.8	43:35	78	NPUAP (1989)	<24	9	8	13	48	0.41 (0.24-0.57)	0.86 (0.79-0.92)	2.86 (1.13-7.07)	0.69 (0.47-0.97)	4.15 (1.17-15.06)
2003	Jun et al.	Korea	Prospective	ICU	PU (±)	62.0	64:48	112	AHCPR (1994)	<24	31	30	4	47	0.89 (0.75-0.96)	0.61 (0.55-0.65)	2.27 (1.65-2.71)	0.19 (0.06-0.46)	12.14 (3.58-45.36)
1997	Kim et al.	Korea	Prospective	ICU	PU (-)	56.2	162:91	253	Lowthian (1989)	<26	52	45	45	111	0.54 (0.46-0.61)	0.71 (0.66-0.76)	1.86 (1.34-2.55)	0.65 (0.51-0.83)	2.85 (1.63-5.01)

Notes: PU= Pressure ulcers; TP= True positive; FP= False positive; FN= False negative; TN= True negative; SN= Sensitivity; SP= Specificity; PLR= Positive likelihood ratio; NLR= Negative likelihood ratio; DOR= Diagnosis odds ratio; Retro= Retrospective study; Hosp.= Hospital; ICD-9= International Classification of Diseases version 9; Pro= Prospective study; LTCF= Long term care facilities; NPUAP= National Pressure Ulcer Advisory Panel; EPUAP= European Pressure Ulcer Advisory Panel; AHCPR= Agency for Health Care Policy and Research; TDCPS= Torrence Developmental Classification of Pressure Sore.

Table 3: Summary results of Meta-analysis by four risk assessment tools

Scale	Study No.	Sensitivity (95% CI)	Specificity (95% CI)	Positive LR (95% CI)	Negative LR (95% CI)	DOR (95% CI)	AUC (SE)	Q* (SE)
Cubbin & Jackson	4	0.67 (0.60-0.74)	0.75 (0.71, 0.79)	2.80 (1.66-4.72)	0.34 (0.15, 0.76)	9.46 (2.41, 37.22)	0.82 (0.06)	0.75 (0.06)
Modified Braden	6	0.95 (0.91-0.98)	0.70 (0.68, 0.73)	3.23 (1.74-6.00)	0.11 (0.06, 0.21)	35.16 (16.62, 74.37)	0.93 (0.03)	0.87 (0.03)
by Choi & Song	4	0.97 (0.92, 0.99)	0.70 (0.66, 0.73)	3.47 (1.33, 9.06)	0.08 (0.04, 0.19)	56.56 (21.88, 146.21)	0.95 (0.02)	0.90 (0.03)
by Pang & Wong	2	0.89 (0.71, 0.98)	0.71 (0.67, 0.75)	2.87 (1.88, 4.38)	0.17 (0.06, 0.49)	16.06 (4.75, 54.35)	-	-
Norton	7	0.75 (0.70-0.79)	0.57 (0.55, 0.59)	1.77 (1.26, 2.50)	0.49 (0.32-0.76)	7.57 (2.53-22.64)	0.82 (0.05)	0.75 (0.04)
Waterlow	6	0.55 (0.49-0.62)	0.82 (0.80, 0.85)	2.89 (1.74, 4.79)	0.46 (0.31-0.70)	9.22 (6.43-13.23)	0.82 (0.03)	0.75 (0.03)
Sub-group Analysis	3							
Norton vs		0.53 (0.43, 0.63)	0.88 (0.83, 0.91)	14.87 (0.12, 1805.61)	0.56 (0.39,, 0.80)	25.67 (2.28, 289.21)	0.81 (0.06)	0.75 (0.05)
Waterlow		0.74 (0.64, 0.82)	0.66 (0.60, 0.72)	2.30 (1.37, 3.83)	0.23 (0.06, 0.87)	8.84 (4.92, 15.89)	0.82 (0.03)	0.76 (0.03)

Note: LR=Likelihood ration; DOR= Diagnostic odds ratio; AUC= Area under the curve; SE= Standard error.