



Awareness of Patients and Relatives about Breast Cancer in Turkey

Serdar Gökay TERZIOĞLU, *Murat Özgür KILIÇ, Gül DAĞLAR

Dept. of General Surgery, Numune Training and Research Hospital, Ankara, Turkey

***Corresponding Author:** Email: murat05ozgur@hotmail.com

(Received 10 May 2017; accepted 24 May 2017)

Dear Editor-in-Chief

Increasing the awareness and knowledge of breast cancer (BC) is the most direct way to improve women's behavior for participating prevention programs.

In this cross-sectional study, the awareness of BC was investigated in patients and relatives in a Breast-Endocrine Clinic, Ankara Numune Training and Research Hospital, Turkey between May and July 2016. The survey form was modified from a study on BC awareness of hospital staffs (1).

The participants were informed about the purpose of the study and verbal informed consent was obtained from all.

Among 164 participants, 75% of the participants were female, and 54.8% were under 45. Educational status was primary/secondary school (53.7%) and \geq high school (46.3%). The majority of participants (82.9%) had no family history of BC. Age, family history, hormone replacement treatment (HRT), and early menarche were identified as a risk factor of BC by approximately 75%, 60%, 50%, and 35% of the participants, respectively. The overall success of general belief section was 80%. The comparison of answers according to age, gender, and educational status is presented in Table 1.

The overall awareness about BC risk factors in our study was 50%, similar to the literature (2). Age and family history were the best-known risk factors. In general, belief section, correct answer

rate was higher than that in other sections. The overall awareness was better than normal population in our study. In our opinion, the overall high correct answer rate of participants pointed out that exposure to the disease forced responders to do more research on BC.

Awareness of BC tends to decrease with increasing age (3). In our study, subgroup analysis about age failed to reveal a decrease, probably based on increased awareness of older responders due to the reason for doing more research BC. Gender is also a determining factor for awareness of cancers. In a study, male participants had similar knowledge about "uni-sex" cancers such as lung, but for BC, the level of awareness was determined significantly lower when age and education adjustments were made (4). In our study, questions about HRT and early menarche had a correct answer rate in females. This can be explained by the opinion that risk factors defined as "complex" seem to be more familiar with women. Educational status is also related to awareness. In our study, well-educated responders had a correct response rate of 80% higher than the literature (5). BC symptoms were often well known by our participants. As expected, the best-known symptoms were mass and change in breast shape. The weight of the correct answer from female participants also draws attention in this section.

Table 1: Comparison of answers about BC

	<i>Correct n (%)</i>	<i>Wrong n (%)</i>	<i>P</i>	<i>Correct n (%)</i>	<i>Wrong n (%)</i>	<i>P</i>
	<i>Does age increase BC risk?</i>			<i>Is BC a contagious disease?</i>		
Gender						
Male	26 (76.4)	8 (13.6)	0.748	32 (100)	0 (0)	
Female	90 (72.5)	34 (26.5)		86 (74.1)	30 (25.9)	0.031
Education						
Primary/Secondary	62 (73.8)	22 (26.2)	0.933	58 (70.7)	24 (29.6)	
≥High school	54 (84.4)	10 (15.6)		60 (90.9)	6 (9.1)	0.032
Age						
>45	53 (63.1)	31 (36.9)	0.048	54 (73)	20 (27)	0.248
≤45	62 (83.8)	12 (16.2)		59 (82)	13 (18)	
	<i>Does HRT increase BC risk?</i>			<i>Is BC a fatal disease?</i>		
Gender						
Male	10 (29.4)	24 (70.6)		22 (64.7)	12 (35.3)	
Female	66 (56.9)	50 (43.1)	0.046	66 (56.4)	51 (43.6)	0.688
Education						
Primary/Secondary	44 (53.6)	38 (46.4)		50 (61)	32 (39)	
≥High school	36 (53)	32 (47)	0.569	36 (54.5)	30 (45.5)	0.577
Age						
>45	44 (57.9)	32 (42.1)		40 (54.1)	34 (45.9)	
≤45	32 (42.1)	42 (57.9)	0.204	46 (62.1)	28 (37.9)	0.480
	<i>Does family history of BC increase BC risk?</i>			<i>Do only females have BC?</i>		
Gender						
Male	18 (56.2)	14 (43.7)	0.768	24 (75)	8 (25)	
Female	72 (60)	48 (40)		68 (58.6)	48 (41.4)	0.232
Education						
Primary/Secondary	48 (58.5)	34 (41.5)	0.897	42 (48.8)	44 (51.2)	
≥High school	42(60)	28 (40)		52 (78.8)	14 (21.2)	0.008
Age						
>45	46 (58.9)	32 (41.1)	0.966	44 (59.4)	30 (40.6)	
≤45	44 (61.1)	28 (38.9)		48 (64.9)	26 (35.1)	0.632
	<i>Does early menarche increase BC risk?</i>					
Gender						
Male	4 (12.5)	28 (87.5)	0.024			
Female	50 (43.1)	66 (56.9)				
Education						
Primary/Secondary	32 (39)	50 (61)	0.613			
≥High school	22 (33.3)	44 (66.7)				
Age						
>45	30 (40.5)	44 (59.5)	0.469			
≤45	24 (32.4)	50 (67.6)				

*Chi-square/Fisher exact tests were used for statistical analyses.

**P<0.05 (significance level)

Well-educated patients with a breast disease or their relatives can be a correct source for education of general population. Initiation of the

education programs for this group may be a feasible option to realize this idea.

Conflict of Interests

The authors declare that there is no conflict of interests.

References

1. Terzioğlu G, Özgü E, Kılıç MÖ et al (2017). Evaluation of breast cancer knowledge and awareness among hospital staff in a women health hospital in Turkey. *J Cancer Educ*, 32(1):59-64.
2. Sherman SM, Lane EL (2015). Awareness of risk Factors for Breast, Lung and Cervical Cancer in a UK Student Population. *J Cancer Educ*, 30(4):660-3.
3. Mandelblatt JS, Schechter CB, Yabroff KR et al (2005). Toward optimal screening strategies for older women. Costs, benefits, and harms of breast cancer screening by age, biology, and health status. *J Gen Intern Med*, 20(6):487-96.
4. Costa AR, Silva S, Moura-Ferreira P et al (2017). Cancer screening in Portugal: sex differences in prevalence, awareness of organized programmes and perception of benefits and adverse effects. *Health Expect*, 20(2):211-220.
5. Hadi MA, Hassali MA, Shafie AA, Awaisu A (2010). Evaluation of breast cancer awareness among female university students in Malaysia. *Pharm Pract (Granada)*, 8(1):29-34.

Archive of SID