



## The Frequency of Resistance Genes in *Salmonella enteritidis* Strains Isolated from Cattle

\*Reza RANJBAR<sup>1</sup>, Farhad SAFARPOOR DEHKORDI<sup>2</sup>, Mohammad HEIAT<sup>3</sup>

1. Molecular Biology Research Center, Systems Biology and Poisonings Institute, Baqiyatallah University of Medical Sciences, Tehran, Iran
2. Halal Research Center of IRI, FDA, Tehran, Iran
3. Baqiyatallah Research Center for Gastroenterology and Liver Disease, Baqiyatallah University of Medical Sciences, Tehran, Iran

\*Corresponding Author: Email: ranjbarre@gmail.com

(Received 10 Dec 2018; accepted 21 Feb 2019)

### Abstract

**Background:** *Salmonella enteritidis* causes infections in humans and animals. Antibiotics are used to eliminate bacterial infections, which become resistant to antibiotics after a while. This study aimed to isolate *S. enteritidis* from cattle feces samples and also to evaluate the frequency of genes associated with multi-drug resistance (MDR).

**Methods:** One hundred ten fecal samples of cattle were collected from Jul to Dec, 2017 in Khuzestan Province, southern Iran. Bacterial culture and molecular methods were used to isolate and identify *S. enteritidis*. Disk diffusion method was used to assess antibiotic susceptibility. Then Polymerase chain reaction (PCR), assay was used for definitive diagnosis of *S. enteritidis* and resistance genes.

**Results:** Overall, 101 (91.81%) samples were detected to be contaminated with *Salmonella* genus and 86 samples (85.14%) were identified as *S. enteritidis*. The highest and lowest antibiotic resistance were belonged to gentamicin (n=70, 81.39%), and tetracycline (n=6, 6.97%). Besides, 64 samples (74.42%) had 2-10 drugs resistance patterns. Moreover, the highest and the least resistance were related to *bla*<sub>IMP-1</sub> (n=73, 84.88%) and *tet*(B) (n=49, 56.97%) genes respectively.

**Conclusion:** The drug-induced genes in *S. enteritidis* have a high frequency. Therefore, antibiotic resistance and high MDR to antibiotics can be due to the incorrect use of antibiotics and the lack of health monitoring in Cattle farms.

**Keywords:** *Salmonella enteritidis*; Resistance gene; Multi-drug resistance; Cattle

## Introduction

*Salmonella* is one of the most important members of the Enterobacteriaceae family. This bacterium is a facultative inter-cellular, gram-negative, rod-shaped, bacilli-shaped, spore-free, aerobic and anaerobic pathogen (1). Most of the pathogenic *Salmonella* species that cause human disease belong to the *S. enterica* species, which live on the host gastrointestinal tract. *S. enteritidis* (*S. enteritidis*) is

among the most important disease-causing serotypes. So far, more than 2,500 *Salmonella* serovars have been identified based on flagellar and somatic antigens, most of which are pathogenic to humans and animals (2,3).

*Salmonella* in humans causes diseases such as salmonellosis, intestinal or typhoid fever, septicemia and gastroenteritis. In this regard, *S. enteritidis* plays

a significant role in the development of these infections (4,5). Diseases transmitted through food are the most important health issues. *Salmonella* is often found in food, meat and dairy products, related to the livestock industry (6,7).

The use of antibiotics for humans and animals causes major health problems, including the presence of antibiotic residues in body tissues and livestock products and the resistance of pathogens to antibiotics, leading to the selection of pathogenic bacteria resistant to multiple drugs (4,8). Multi-drug resistance (MDR) in bacteria is caused by various mechanisms, including reactive changes of the target protein and the enzymatic inactivation of the drug. Enzymatic inactivation is a common resistance mechanism for natural antibiotics such as aminoglycosides and  $\beta$ -lactams (9-11).

Due to increasing rate of MDR bacterium, the purpose of this study was to use PCR as a precise method for detecting *S. enteritidis* and also to determine the frequency of resistance genes of this bacterium in cattle samples in Iran.

## Materials and Methods

### Sampling

The present study was conducted from Jul to Dec 2017 in Khuzestan Province, southern Iran. Overall, 110 cattle feces samples were collected randomly and used for detection of *Salmonella*.

### Isolation and Culture of *Salmonella*

The feces samples were transferred to concentrated peptone water and incubated at 37 °C for 24 h. Then, the samples were transferred to Rappaport Vassiliadis (RV) and incubated again at 43 °C for 24 h. The isolated bacteria were cultured on *Salmonella Shigella* Agar (SSA). Black and gray colonies were considered as suspected *Salmonella*. Complementary assays and differential microbial tests (i.e., IMViC and urease) and culturing on Triple Sugar Iron Agar (TSI) according to Bergey's manual were used to detect *Salmonella* strain. The lysine decarboxylation was performed on isolated bacteria by culturing in lysine decarboxylase broth

to identify the *Salmonella* family. Sulfide Indole Motility medium (SIM) was used for direct detection of *Salmonella* spp. by sulfide production, indole formation, and motility.

### Gene Amplification

Genomic DNA was extracted using QIAamp mini kit (Qiagen, GmbH, Germany). PCR method using specific oligonucleotide primers for *16S rRNA* gene was performed for the direct detection of *Salmonella* family. In the next step, *Salmonella*-encoded fimbria (*sef-A*) gene was targeted for specific detection of *S. enteritidis* from other *Salmonella* strains. The primer sequences used in the present study are shown in Table 1. PCR reactions were performed in a total volume of 25  $\mu$ L in 0.2 ml tubes containing 2  $\mu$ L of template DNA, 1  $\mu$ M of each primer, 2 mM MgCl<sub>2</sub>, 5  $\mu$ L of 10X PCR buffer AMS, 200  $\mu$ M dNTPs, and 1 unit of *Taq* DNA polymerase (CinnaClon Co, Iran). The PCR assay was performed at 95 °C for 5 min and then for 32 cycles of 94 °C for 1 min, annealing temp according to Table 1 for 40 sec, 72 °C for 40 sec, and a final extension at 72 °C for 5 min, with a final hold at 10 °C in a thermal cycler (Mastercycler gradient, Eppendorf, Germany). The PCR amplified products were detected in 2% EtBr stained agarose gel electrophoresis.

### Antimicrobial drug susceptibility tests

The colonies were cultured on Nutrient Agar (Merck, Darmstadt, Germany) at 37 °C for 24 h after detection of positive *S. enteritidis* samples from other *Salmonella* strains. Then, the colonies were sub-cultured into Nutrient Broth (Merck, Germany) and incubated at 37 °C in a shaking incubator for 14 h. McFarland 0.5 turbidity standards were used for bacterial turbidity. After standardization, 100  $\mu$ L of bacterial suspensions were cultured on Mueller-Hinton agar medium (Merck, Darmstadt, Germany). Drug resistance testing was performed using standard Bauer-Kirby disk diffusion method for all samples (positive culture) according to CLSI protocols for determination of antibiotic susceptibility patterns of *S.*

**Table 1:** Primer sequence of resistance genes and related information in *S. enteritidis*

<b>Resistance Genes</b>	<b>Nucleotide sequence (5'to 3')</b>	<b>Annealing Temperature</b>	<b>Size (bp)</b>	<b>Reference</b>
<i>aadA1</i>	F:CTCCGCAGTGGATGGCGG R:GATCTGCGCGGAGGCCA	65	311	3
<i>aadA2</i>	F:CATTGAGCGCCATCTGGAAT R:ACATTTTCGCTCATCGCCGGC	65	432	3
<i>aadB</i>	F:CTAGCTGCGGCAGATGAGC R:CTCAGCCGCCTCTGGGCA	62	219	3
<i>aadD</i>	F:TATATCCGTGTCTCTCTGTCCA R:CTCTATTTTGCCGATTTATGATTC	55	419	12
<i>strA</i>	F:TGGCAGGAGGAACAGGAGG R:AGGTCGATCAGACCCGTGC	62	608	3
<i>strB</i>	F:GCGGACACCTTTTCCAGCCT R:TCCGCCATCTGTGCAATGCG	65	256	3
<i>bla<sub>TEM-1</sub></i>	F:CAGCGGTAAGATCCTTGAGA R:ACTCCCCGTCTGTAGATAA	55	643	13
<i>bla<sub>CMY-2</sub></i>	F:TGGCCGTGCGCTTATCTAC R:CCCGTTTTATGCACCCATGA	55	870	13
<i>bla<sub>IMP-1</sub></i>	F:TGAGGCTTACCTAATTGACA R:TCAGGCAACCAACCCTAC	55	324	13
<i>bla<sub>CTX-M1</sub></i>	F:AACCGTCACGCTGTTGTTAG R:TTGAGGCTGGGTGAAGTAAG	55	766	13
<i>bla<sub>OXA-1</sub></i>	F:AATGGCACCAGATTCAACTT R:CTTGGCTTTTATGCTTGATG	55	595	13
<i>bla<sub>PSE-1</sub></i>	F:TGCTTCGCAACTATGACTAC R:AGCCGTGTGTTTGAGCTAGAT	55	438	13
<i>sul1</i>	F:TTTCCTGACCCTGCGCTCTAT R:GTGCGGACGTAGTCAGCGCCA	55	793	14
<i>sul2</i>	F:CCTGTTTCGTCCGACACAGA R:GAAGCGCAGCCGCAATTCAT	55	667	14
<i>tet(A)</i>	F:TTGGCATCTGCAATCACTC R:GTATAGCTTGCCGGAAGTCG	55	494	14
<i>tet(B)</i>	F:CAGTGCTGTTGTGTCATTA R:GCTTGGAATACTGAGTGTA	55	571	14
<i>tet(G)</i>	F:GCTCGGTGGTATCTCTGCTC R:CAAAGCCCCTTGCTTGTTAC	55	550	14
<i>cat1</i>	F:AACCAGACCGTTCAGCTGGAT R:CCTGCCACTCATCGCAGTAC	55	549	14
<i>cat2</i>	F:AACGGCATGAACCTGAA R:ATCCCAATGGCATCGTAAAG	55	547	14
<i>floR</i>	F:ATGACCACCACACGCCCCG R:AGACGACTGGCGACTTCTTCG	55	198	14
<i>cmlA</i>	F:GGCCTCGCTCTTACGTATC R:GCGACACCAATACCCACTAGC	55	662	14
<i>sefA</i>	F:TGCTATTTTGCCCTGTACTGC R:TTCCGGGGGAGACTATACCTACAG	58	214	15
<i>16s rRNA</i>	F:AACCGACTCACTCTGGCAG R:TAACGCGATAGCGCTTC	58	214	15

*enteritidis* isolated from feces specimens and the cultures were incubated at 37 °C for 24 h. The antibiotic disks were used for determination of antibiotic susceptibility patterns of *S. enteritidis* isolated from feces specimens.

#### Detection of antibiotic resistance genes

The PCR was performed for identification of resistance genes. The primers were blasted at the NCBI using the experimental GENINFO BLAST Network Service to assess degree of homology between these primers and other reported sequences and at the end were obtained from CinnaClon Co, Iran. The specific oligonucleotide primers for gene amplification of antimicrobial resistance genes in isolated *S. enteritidis* and the sequences of primers are available in Table 1. The PCR was run in final reaction volumes of 25 µL containing the following reagents: 0.2 ml tubes containing 2 µL of template DNA, 1 µM of each primer, 2 mM MgCl<sub>2</sub>, 5 µL of 10X PCR buffer AMS, 200 µM dNTPs, and 1 unit of *Taq* DNA polymerase (CinnaClon Co, Iran). Reactions were initiated at 95 °C for 5 min, followed by 32 cycles of 94 °C for 1 min, annealing temp according to Table 1 for 40 sec, 72 °C for 40 sec and a final extension step at 72 °C for 5 min, with a final hold at 10 °C in a thermal cycler (Mas-

tercycler gradient, Eppendorf, Germany). Detection of PCR products was performed by the method mentioned earlier.

#### Statistical Analysis

All statistical tests were performed using Graph Pad Prism statistical software, ver. 7.00 (Graph Pad, San Diego, CA, USA). For all tests, *P*-value <0.05 was considered statistically significant.

## Results

#### Identification of bacteria

Out of 110 samples, 101 (91.81%) samples were positive for *Salmonella* by detecting 258 bp band of 16S *rRNA* gene. Furthermore, the numbers of 86 (85.14%) specimen were *sefA* gene positive (488 bp).

#### Antibiotic susceptibility

The highest antibiotic resistance was observed to gentamicin (n=70, 81.39%) and the lowest antibiotic resistance was observed in tetracycline (n=6, 6.97%). Moreover, while the lowest antibiotic susceptibility belonged to sulfamethoxazole (n=3, 3.48%) (Fig.1). According to the results, 64 samples had 2-10 drugs resistance patterns (Table 2) in cattle (74.42%).




Fig. 1: The frequency of antibiotic resistance among the *S. enteritidis* under the study

Table 2: The prevalence of MDR in cattle specimens

NO. of MDR	Antibiotics										NO. of Animal (n=86)	Per-cent
	S	GM	AM	C	SMX	TE	AN	P	CRO	IPM		
0	S	GM	AM	C	SMX	TE	AN	P	CRO	IPM	16	18.60
1	GM										6	6.98
2	GM	SMX									7	8.14
3	GM	SMX	CRO								5	5.81
4	S	GM	SMX	CRO							3	3.49
5	S	GM	SMX	AN	CRO						1	1.16
6	S	GM	SMX	AN	CRO	IPM					33	38.37
7	S	GM	C	SMX	AN	CRO	IPM				5	5.81
9	S	GM	AM	C	SMX	AN	P	CRO	IPM		4	4.65
10	S	GM	AM	C	SMX	TE	AN	P	CRO	IPM	6	6.98

### Resistance genes

In the studied samples, the highest resistance belonged to the *bla*<sub>IMP-1</sub> genes (n=73, 84.88%) and the lowest resistance, on the other hand, was for *tet*(B) (n=49, 56/97%) gene. Detection of resistance genes in isolated *S. enteritidis* using PCR technique revealed fragments with specific length sizes in Table 1.

### Discussion

The results of present study revealed that 91.81% and 85.14% were detected for *Salmonella* genus and *S. enteritidis* respectively. *bla*<sub>IMP-1</sub> (84.88%) and *tet*(B) (56.97%) genes indicated the highest and the lowest frequency in comparison to other studied genes. However, the highest and lowest antibiotic resistances belonged to gentamicin (81.39%), and tetracycline (6.97%) antibiotics.

In Iran, a high prevalence of *S. enteritidis* (43%) was reported in comparison with *S. typhimurium*. In contrast, a low *Salmonella* (1.6%) prevalence was reported in Italy (16-18). The frequency of aminoglycoside resistance genes of *aadA1*, *aadA2*, *aadB*, *strA*, and *strB*, of 245 samples were investigated,

62 samples belonged to the *S. enteritidis* serotype and the frequency of genes was 45.6%, 34.7%, 31.1%, 37.6%, and 22.4%, respectively (3). By comparing the frequency of genes with the present study, the frequencies are consistent. In Egypt, a study was conducted on 1600 samples to evaluate the multi-drug resistance. The  $\beta$ -lactamases encoding genes were identified in 75.4% of isolates. The frequency of *bla*<sub>TEM-1</sub> gene (41.5%) was relatively high. The highest resistance belonged to ampicillin (95.7%), kanamycin (93.6%), spectinomycin (93.6%), streptomycin (91.5%) and sulfamethoxazole (91.5%) (19). In comparison, in the present study, the highest resistance belonged to gentamicin (81.39%). The frequency of aminoglycoside, tetracycline, sulfonamide, and chloramphenicol genes was investigated and showed the frequency of *sul1* (76.6%), *cat1* (43.3%), *tetA* (40%) and *aadA1* (36.7%) genes (14). Randall et al. conducted some studies on most resistance genes on 397 specimens by culture and molecular methods. They identified resistance to ampicillin (91 isolates), chloramphenicol (85 isolates), gentamicin (2 isolates), streptomycin (119 isolates), tetracycline (108 isolates) and 219 susceptible specimens to all

antibiotics (20). In our study, resistance to ampicillin (10 isolates), chloramphenicol (15 isolates), gentamicin (70 isolates), streptomycin (52 isolates) and tetracycline (6 isolates) were identified and 58 isolates were found sensitive to penicillin. Finally, by comparing the two studies, there was a significant relation between the presence of resistance genes and related resistance phenotypes ( $P < 0.001$ ).

The phenomenon of antimicrobial resistance has been an urgent global problem since the 1990s (21). The excessive and incorrect use of antimicrobial agents and lack of appropriate infection control are the main possible reasons for this alarming phenomenon and development of MDR in bacteria (22).

In a study, the rate of 52% MDR was shown for Cattle (23). Firoozeh et al. in an Iranian study investigated the prevalence of *S. enteritidis* and the MDR. They identified 77.7% prevalence of *S. enteritidis*, 81% drug resistance, and 69% two or more resistances. The highest resistance phenotypes were streptomycin (58.3%), ampicillin (21.4%), sulfamethoxazole (17.85%), kanamycin (14.28%) and chloramphenicol (14.2%) (24). However, in present study, the highest resistance phenotypes were streptomycin, gentamicin, sulfamethoxazole, amikacin, chloramphenicol and imipenem antibiotics. Due to the presence of streptomycin and sulfamethoxazole antibiotics and the creation of resistance phenotypes in both studies, care should be taken in their use.

## Conclusion

The comparison of the results of present study with others suggests that both the prevalence and the antimicrobial resistance pattern for *Salmonella* is different based on geographical zones. This study determined the widely spread prevalence of *S. enteritidis* and MDR in cattle from Khuzestan Province. Efforts that include further implementation of hazard analysis of critical control in livestock production are needed to reduce the incidence of *Salmonella*.

## Ethical considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

## Acknowledgements

We would like to thank from the “Clinical Research Development Center of Baqiyatallah hospital” for their kindly cooperation.

## Funding

Financial support for the study and its publication were provided by the Clinical Research Development Center of Baqiyatallah Hospital, Tehran, Iran (2017).

## Conflict of interest

The authors declare that there is no conflicts of interest.

## References

1. Ranjbar R, Mortazavi SM, Mehrabi Tavana A et al (2017). Simultaneous Molecular Detection of *Salmonella enterica* Serovars Typhi, Enteritidis, Infantis, and Typhimurium. *Iran J Public Health*, 46(1):103-111.
2. Ranjbar R, Naghoni A, Afshar D et al (2016). Rapid Molecular Approach for Simultaneous Detection of *Salmonella* spp., *Shigella* spp., and *Vibrio cholera*. *Osong Public Health Res Perspect*, 7(6):373-377.
3. Doosti A, Mahmoudi E, Jami MS et al (2016). Prevalence of *aadA1*, *aadA2*, *aadB*, *strA* and *strB* genes and their associations with multi-drug resistance phenotype in *Salmonella* Typhimurium isolated from poultry carcasses. *Thai J Vet Med*, 46 (4):691-697.
4. Crum Cianflone N F (2008). Salmonellosis and the GI Tract: More than Just Peanut Butter. *Curr Gastroenterol Rep*, 10(4):424-431.

5. Bula-Rudas FJ, Rathore MH, Maraqa NF (2015). Salmonella Infections in Childhood. *Adv Pediatr*, 62(1):29-58.
6. Naghoni A, Ranjbar R, Tabaraie B et al (2010). High prevalence of integron-mediated resistance in clinical isolates of Salmonella enterica. *Jpn J Infect Dis*, 63(6):417-421.
7. Eng SK, Pusparajah P, Mutalib NSA et al (2015). Salmonella: A review on pathogenesis, epidemiology and antibiotic resistance. *Frontiers in Life Science*, 8(3):284-293.
8. Chen HM, Wang Y, Su LH et al (2013). Nontyphoid Salmonella infection: microbiology, clinical features, and antimicrobial therapy. *Pediatr Neonatol*, 54(3):147-152.
9. Alekshun MN, Levy SB (2007). Molecular mechanisms of antibacterial multidrug resistance. *Cell*, 128(6):1037-1050.
10. Ranjbar R, Giammanco GM, Aleo A et al (2010). Characterization of the first extended-spectrum beta-lactamase-producing nontyphoidal Salmonella strains isolated in Tehran, Iran. *Foodborne Pathog Dis*, 7(1):91-95.
11. Ranjbar R, Naghoni A, Yousefi S et al (2013). The Study of Genetic Relationship Among Third Generation Cephalosporin-resistant Salmonella enterica Strains by ERIC-PCR. *Open Microbiol J*, 7:142-145.
12. Ma M, Wang H, Yu Y et al (2007). Detection of antimicrobial resistance genes of pathogenic Salmonella from swine with DNA microarray. *J Vet Diagn Invest*, 19(2):161-167.
13. Chen S, Zhao S, White DG et al (2004). Characterization of multiple-antimicrobial-resistant Salmonella serovars isolated from retail meats. *Appl Environ Microbiol*, 70(1):1-7.
14. Adesiji YO, Deekshit VK, Karunasagar I (2014). Antimicrobial-resistant genes associated with Salmonella spp. isolated from human, poultry, and seafood sources. *Food Sci Nutr*, 2(4):436-442.
15. Doosti A, Doosti E, Rahimi E et al (2017). Frequency of Antimicrobial-Resistant Genes in Salmonella enteritidis Isolated from Traditional and Industrial Iranian White Cheeses. *Proc Natl Acad Sci India Sect B Biol Sci*, 87(1):73-80.
16. Afshari A, Baratpour A, Khanzade S et al (2018). Salmonella Enteritidis and Salmonella Typhimorium identification in poultry carcasses. *Iran J Microbiol*, 10(1):45-50.
17. Bonardi S, Bruini I, Magnani R et al (2017). Low Prevalence of Salmonella Enterica in Cull Dairy Cattle at Slaughter in Northern Italy. *Ital J Food Saf*, 6(1):6172.
18. Ghaderi R, Moradi Bidhendi S, Khaki P (2016). Occurrence of multidrug-resistant Salmonella enterica serovar Enteritidis isolates from poultry in Iran. *ARI*, 71(1):43-49.
19. Ahmed AM, Shimamoto T, Shimamoto T (2014). Characterization of integrons and resistance genes in multidrug-resistant Salmonella enterica isolated from meat and dairy products in Egypt. *Int J Food Microbiol*, 189:39-44.
20. Randall LP, Cooles SW, Osborn MK et al (2004). Antibiotic resistance genes, integrons and multiple antibiotic resistance in thirty-five serotypes of Salmonella enterica isolated from humans and animals in the UK. *J Antimicrob Chemother*, 53(2):208-216.
21. Jahandeh N, Ranjbar R, Behzadi P et al (2015). Uropathogenic Escherichia coli virulence genes: invaluable approaches for designing DNA microarray probes. *Cent European J Urol*, 68(4):452-458
22. Ranjbar R, Haghi-Ashtiani M, Jonaidi Jafari N et al (2009). The Prevalence and Antimicrobial Susceptibility of Bacterial Uropathogens Isolated from Pediatric Patients. *Iran J Public Health*, 38(2):134-138.
23. Kemal J, Sibhat B, Menkir S et al (2015). Antimicrobial resistance patterns of Salmonella in Ethiopia: A review. *Afr J Microbiol Res*, 9(46):2249-2256.
24. Firoozeh F, Zahraei-Salehi T, Shahcheraghi F et al (2012). Characterization of class I integrons among Salmonella enterica serovar Enteritidis isolated from humans and poultry. *FEMS Immunol Med Microbiol*, 64(2):237-243.